

**Table 1. Cancer incidence trends by joinpoint analyses\* for 1973 through 1999  
for the most common cancers by race, sex, and age, SEER†, 1973-1999.**

Site/Race/Sex	Age	Joinpoint Analyses* (1973-1999)							
		Trend 1		Trend 2		Trend 3		Trend 4	
		Range of Years	APC‡	Range of Years	APC‡	Range of Years	APC‡	Range of Years	APC‡
All Sites									
All Races - Both Sexes	All ages	1973-83	0.9 §	1983-92	1.8 §	1992-95	-1.9	1995-99	0.2
	<20	1973-91	1.0 §	1991-99	-0.2				
	20-49	1973-80	-0.6 §	1980-90	1.5 §	1990-99	-0.7 §		
	<50	1973-80	-0.5 §	1980-90	1.5 §	1990-99	-0.7 §		
	50-64	1973-99	0.9 §						
	65-74	1973-89	1.7 §	1989-92	4.8 §	1992-95	-3.3 §	1995-99	0.2
	75+	1973-92	1.5 §	1992-95	-3.3	1995-99	0.2		
All Races - Males	All ages	1973-89	1.4 §	1989-92	5.0 §	1992-95	-4.9 §	1995-99	-0.3
	<20	1973-97	0.8 §	1997-99	-4.7				
	20-49	1973-82	0.5 §	1982-87	3.5 §	1987-92	1.2 §	1992-99	-2.4 §
	<50	1973-82	0.4 §	1982-86	3.8 §	1986-92	1.3 §	1992-99	-2.2 §
	50-64	1973-85	1.3 §	1985-89	0.2	1989-92	4.6 §	1992-99	-0.1
	65-74	1973-89	1.6 §	1989-92	8.2 §	1992-95	-5.5 §	1995-99	-0.2
	75+	1973-89	1.3 §	1989-92	4.0	1992-95	-7.8 §	1995-99	-0.1
All Races - Females	All ages	1973-80	0.2	1980-87	1.6 §	1987-99	0.3 §		
	<20	1973-99	0.7 §						
	20-49	1973-80	-1.2 §	1980-86	1.2 §	1986-99	0.2		
	<50	1973-80	-1.1 §	1980-86	1.2 §	1986-99	0.2		
	50-64	1973-75	3.5	1975-78	-2.2	1978-99	0.7 §		
	65-74	1973-78	0.6	1978-87	2.4 §	1987-99	0.3 §		
	75+	1973-89	1.5 §	1989-99	0.1				
White - Both Sexes	All ages	1973-83	1.0 §	1983-92	1.9 §	1992-95	-2.1	1995-99	0.2
	<20	1973-99	0.7 §						
	20-49	1973-80	-0.5	1980-89	1.9 §	1989-99	-0.7 §		
	<50	1973-80	-0.4	1980-89	1.9 §	1989-99	-0.7 §		
	50-64	1973-82	0.4	1982-92	1.3 §	1992-99	0.5		
	65-74	1973-89	1.8 §	1989-92	4.8 §	1992-95	-3.5 §	1995-99	0.3
	75+	1973-92	1.5 §	1992-95	-3.5	1995-99	0.2		
White - Males	All ages	1973-89	1.4 §	1989-92	4.9 §	1992-95	-5.3 §	1995-99	-0.2
	<20	1973-99	0.6 §						
	20-49	1973-83	0.7 §	1983-86	5.8 §	1986-92	1.1 §	1992-99	-2.7 §
	<50	1973-82	0.6 §	1982-86	4.2 §	1986-91	1.7 §	1991-99	-2.3 §
	50-64	1973-89	1.1 §	1989-92	3.9 §	1992-99	-0.1		
	65-74	1973-88	1.5 §	1988-92	6.6 §	1992-95	-5.6 §	1995-99	-0.2
	75+	1973-89	1.4 §	1989-92	3.7	1992-95	-8.0 §	1995-99	-0.2
White - Females	All ages	1973-80	0.3	1980-87	1.7 §	1987-99	0.3 §		
	<20	1973-99	0.8 §						
	20-49	1973-80	-0.9 §	1980-90	1.1 §	1990-93	-1.4	1993-99	0.7 §
	<50	1973-80	-0.8 §	1980-87	1.2 §	1987-99	0.1		
	50-64	1973-75	3.7	1975-78	-2.3	1978-99	0.7 §		
	65-74	1973-78	0.8	1978-87	2.5 §	1987-99	0.3 §		
	75+	1973-89	1.6 §	1989-99	0.1				

\* Joinpoint analysis of trends allowed for up to three joinpoints based on rates per 100,000 that were age-adjusted to the 2000 U.S. standard population.

† SEER (Surveillance, Epidemiology, and End Results) registries covering 10% of the U.S. population: metropolitan areas of San Francisco, Detroit, Atlanta, and Seattle-Puget Sound; and States of Connecticut, Hawaii, Iowa, Utah, and New Mexico.

‡ APC is the Annual Percent Change based on rates age-adjusted to the 2000 U.S. standard population using the joinpoint regression program.

§ APC is statistically significantly different from zero (two-sided p<.05).

¶ Joinpoint results are not shown if less than 10 cases in any year.

**Table 1 (continued). Cancer incidence trends by joinpoint analyses\* for 1973 through 1999  
for the most common cancers by race, sex, and age, SEER†, 1973-1999.**

Site/Race/Sex	Age	Joinpoint Analyses* (1973-1999)							
		Trend 1		Trend 2		Trend 3		Trend 4	
		Range of Years	APC‡	Range of Years	APC‡	Range of Years	APC‡	Range of Years	APC‡
All Sites									
<b>Black - Both Sexes</b>	<b>All ages</b>	1973-84	1.5 §	1984-89	0.5	1989-92	4.4 §	1992-99	-1.4 §
	<b>&lt;20</b>	1973-99	0.6 §						
	<b>20-49</b>	1973-82	-0.9	1982-94	1.1 §	1994-99	-2.1 §		
	<b>&lt;50</b>	1973-82	-0.8	1982-94	1.1 §	1994-99	-2.1 §		
	<b>50-64</b>	1973-83	2.3 §	1983-90	-0.1	1990-93	4.5	1993-99	-0.5
	<b>65-74</b>	1973-89	1.8 §	1989-92	6.2	1992-99	-2.4 §		
	<b>75+</b>	1973-92	1.4 §	1992-99	-1.3 §				
<b>Black - Males</b>	<b>All ages</b>	1973-81	2.6 §	1981-89	0.6	1989-92	6.7 §	1992-99	-2.7 §
	<b>&lt;20</b>	1973-99	1.0 §						
	<b>20-49</b>	1973-85	0.3	1985-95	1.9 §	1995-99	-5.1 §		
	<b>&lt;50</b>	1973-86	0.1	1986-89	4.8	1989-95	0.7	1995-99	-4.1 §
	<b>50-64</b>	1973-82	3.2 §	1982-89	-1.0	1989-93	6.4 §	1993-99	-1.0
	<b>65-74</b>	1973-89	1.8 §	1989-92	9.5	1992-99	-4.1 §		
	<b>75+</b>	1973-89	1.6 §	1989-92	5.4	1992-95	-7.9	1995-99	-0.6
<b>Black - Females</b>	<b>All ages</b>	1973-91	1.1 §	1991-99	0.1				
	<b>&lt;20</b>	1973-99	0.3						
	<b>20-49</b>	1973-99	-0.1						
	<b>&lt;50</b>	1973-99	-0.1						
	<b>50-64</b>	1973-86	1.7 §	1986-99	0.3				
	<b>65-74</b>	1973-90	2.1 §	1990-99	0.0				
	<b>75+</b>	1973-99	1.0 §						

\* Joinpoint analysis of trends allowed for up to three joinpoints based on rates per 100,000 that were age-adjusted to the 2000 U.S. standard population.

† SEER (Surveillance, Epidemiology, and End Results) registries covering 10% of the U.S. population: metropolitan areas of San Francisco, Detroit, Atlanta, and Seattle-Puget Sound; and States of Connecticut, Hawaii, Iowa, Utah, and New Mexico.

‡ APC is the Annual Percent Change based on rates age-adjusted to the 2000 U.S. standard population using the joinpoint regression program.

§ APC is statistically significantly different from zero (two-sided p<.05).

¶ Joinpoint results are not shown if less than 10 cases in any year.