

Table XXI-4
OVARY CANCER (Invasive)

SEER INCIDENCE^a AND U.S. DEATH^b RATES, AGE-ADJUSTED AND AGE-SPECIFIC RATES, BY RACE

	Incidence ^a			US Death ^b		
	All Races, Females	White Females	Black Females	All Races, Females	White Females	Black Females
<u>AGE-ADJUSTED RATES, 2001-2005</u>						
AGE AT DIAGNOSIS/DEATH:						
All ages	13.3	14.1	10.1	8.8	9.2	7.5
Under 65	8.2	8.6	5.8	3.7	3.8	3.1
65 and over	48.9	51.7	39.6	44.7	46.2	37.7
All ages(IARC world std) ^c	9.5	10.1	7.1	5.6	5.9	4.8
<u>AGE-SPECIFIC RATES, 2001-2005</u>						
AGE AT DIAGNOSIS/DEATH:						
<1	-	-	-	-	-	-
1-4	-	-	-	-	-	-
5-9	0.3	0.3	-	-	-	-
10-14	0.7	0.8	-	-	-	-
15-19	1.5	1.6	1.1	0.1	0.1	-
20-24	1.7	1.7	1.0	0.2	0.2	0.2
25-29	2.1	2.1	1.6	0.3	0.4	0.4
30-34	3.1	3.0	2.9	0.6	0.6	0.6
35-39	5.1	5.1	5.0	1.2	1.3	1.4
40-44	8.9	9.2	6.0	2.8	2.8	2.4
45-49	15.8	16.4	11.0	5.8	6.1	4.9
50-54	22.2	23.5	15.0	10.4	10.9	8.4
55-59	28.8	30.8	18.5	16.0	16.7	13.4
60-64	37.2	40.0	27.9	24.1	25.1	19.9
65-69	41.3	44.1	32.7	32.3	33.6	28.0
70-74	47.4	50.6	36.4	40.2	41.6	35.5
75-79	53.3	56.8	37.6	50.2	52.0	41.5
80-84	57.8	59.8	52.3	59.0	61.0	45.7
85+	51.0	52.3	50.3	55.3	56.7	47.7

^a Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473. SEER 17 areas. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

^c Rates are per 100,000 and are age-adjusted to the IARC world standard population.

- Statistic not shown. Rate based on less than 16 cases for the time interval.