

Table XXV-4
TESTIS CANCER (Invasive)

SEER INCIDENCE^a AND U.S. DEATH^b RATES, AGE-ADJUSTED AND AGE-SPECIFIC RATES, BY RACE

	Incidence ^a			US Death ^b		
	All Races, Males	White Males	Black Males	All Races, Males	White Males	Black Males
<u>AGE-ADJUSTED RATES, 2001-2005</u>						
AGE AT DIAGNOSIS/DEATH:						
All ages	5.4	6.3	1.4	0.3	0.3	0.2
Under 65	6.0	7.1	1.5	0.2	0.3	0.1
65 and over	1.1	1.2	-	0.3	0.3	0.4
All ages(IARC world std) ^c	5.0	5.9	1.3	0.2	0.2	0.1
<u>AGE-SPECIFIC RATES, 2001-2005</u>						
AGE AT DIAGNOSIS/DEATH:						
<1	0.9	0.8	-	-	-	-
1-4	0.4	0.4	-	-	-	-
5-9	-	-	-	-	-	-
10-14	0.2	0.3	-	-	-	-
15-19	3.6	4.3	-	0.1	0.1	-
20-24	9.4	11.3	2.0	0.3	0.3	-
25-29	12.5	15.0	2.5	0.4	0.5	0.2
30-34	12.9	15.5	3.4	0.4	0.4	-
35-39	11.2	13.2	3.1	0.4	0.4	0.3
40-44	9.3	11.0	2.2	0.4	0.5	-
45-49	6.3	7.4	2.0	0.4	0.4	0.3
50-54	4.2	4.8	-	0.3	0.3	-
55-59	2.5	2.8	-	0.2	0.2	-
60-64	1.5	1.7	-	0.2	0.2	-
65-69	1.4	1.6	-	0.2	0.2	-
70-74	0.9	1.0	-	0.2	0.2	-
75-79	0.9	1.0	-	0.3	0.2	-
80-84	1.1	1.1	-	0.4	0.4	-
85+	1.3	1.4	-	0.6	0.7	-

^a SEER 17 areas. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

^c Rates are per 100,000 and are age-adjusted to the IARC world standard population.

- Statistic not shown. Rate based on less than 16 cases for the time interval.