

Table 34.1
Five-Year Cause-Specific Survival^a(%) for Malignant Cancers, 2001-2007
Both Sexes by Race/Ethnicity and Primary Site

Primary Site	White			Black			Asian/ Pacific Islander			American Indian/ Alaska Native ^b			Hispanic ^c		
	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE
All Sites	1,620,695	66.0	0.0	186,908	59.3	0.1	119,296	64.6	0.2	7,587	58.2	0.6	185,177	65.6	0.1
Oral Cavity & Pharynx	35,654	66.5	0.3	4,107	46.9	0.9	3,115	67.7	1.0	192	64.7	4.1	3,270	62.3	1.0
Esophagus	15,190	19.7	0.4	1,956	13.8	1.0	817	16.4	1.7	90	11.8	4.2	1,263	17.1	1.3
Stomach	23,048	26.3	0.3	4,043	28.7	0.8	4,699	36.5	0.8	252	25.6	3.1	5,506	27.3	0.7
Colon & Rectum	161,088	64.5	0.1	20,691	56.9	0.4	14,647	68.3	0.5	970	61.5	1.9	17,876	63.3	0.4
Liver & Intrahepatic Bile Duct	19,849	16.9	0.3	3,335	12.9	0.8	5,535	21.8	0.7	267	17.4	2.8	5,252	17.6	0.7
Pancreas	38,286	5.8	0.1	5,189	5.2	0.4	3,114	7.3	0.6	215	6.8	2.0	4,508	6.4	0.5
Larynx	11,941	67.0	0.5	2,132	58.1	1.3	474	67.7	2.5	40	42.8	10.0	1,175	66.6	1.6
Lung & Bronchus	203,664	17.6	0.1	25,239	14.5	0.3	13,276	18.7	0.4	884	12.1	1.3	13,257	16.2	0.4
Melanoma of the Skin	76,353	89.1	0.1	366	71.5	2.8	524	77.8	2.1	89	74.5	5.8	2,703	84.2	0.8
Breast	242,588	88.4	0.1	27,079	77.5	0.3	20,552	90.3	0.3	1,094	85.6	1.3	27,896	86.2	0.3
Cervix uteri	15,862	72.3	0.4	2,668	62.9	1.1	1,878	72.4	1.2	125	62.4	5.2	4,928	75.1	0.7
Corpus & Uterus, NOS	47,460	82.9	0.2	4,561	62.7	0.8	3,893	84.7	0.7	245	84.4	2.5	5,557	81.5	0.6
Ovary ^d	26,351	43.4	0.4	2,260	36.1	1.2	2,144	55.7	1.3	150	43.6	5.0	3,488	49.0	1.0
Prostate	262,340	93.7	0.1	41,502	91.6	0.2	15,481	94.1	0.2	789	89.9	1.3	28,242	92.6	0.2
Testis	12,851	95.9	0.2	377	88.1	1.8	447	92.0	1.6	94	98.3	1.7	2,647	93.1	0.5
Urinary Bladder	70,787	79.2	0.2	3,797	65.9	0.9	2,875	79.6	0.9	137	71.4	4.4	4,496	76.6	0.7
Kidney & Renal Pelvis	46,246	70.8	0.2	5,311	70.8	0.7	2,567	71.8	1.0	396	68.4	2.7	6,817	71.3	0.6
Brain & Nervous System	25,614	34.0	0.3	1,828	40.5	1.3	1,465	41.1	1.5	109	44.1	5.9	3,985	45.3	0.9
Thyroid	37,358	96.5	0.1	2,665	95.7	0.4	4,470	96.3	0.3	196	96.3	1.4	6,555	96.5	0.3
Hodgkin Lymphoma	11,394	86.2	0.4	1,418	83.0	1.2	615	85.4	1.7	35	89.2	6.2	2,045	84.9	0.9
Non-Hodgkin Lymphoma	69,770	68.7	0.2	6,106	63.6	0.7	4,993	65.7	0.8	271	68.5	3.0	9,017	65.9	0.6
Myeloma	18,031	44.0	0.5	4,148	46.0	1.0	1,195	45.8	1.9	112	47.1	5.8	2,622	44.6	1.3
Leukemia	45,108	56.7	0.3	3,848	52.1	0.9	2,863	53.2	1.1	209	57.0	4.0	6,969	58.6	0.7
Mesothelioma	3,763	7.7	0.6	185	15.1	3.3	121	13.3	3.5	-	-	-	419	10.3	2.0

Source: SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2001-2007 with follow-up through 2008.

^b Data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^c Hispanics are not mutually exclusive from whites, blacks, Asian/Pacific Islanders and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

^d Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

- Statistic could not be calculated due to less than 25 cases in the time interval.

Table 34.2
 Five-Year Cause-Specific Survival^a(%) for Malignant Cancers, 2001-2007
 Males by Race/Ethnicity and Primary Site

Primary Site	White			Black			Asian/ Pacific Islander			American Indian/ Alaska Native ^b			Hispanic ^c		
	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE
All Sites	837,722	66.1	0.1	100,266	61.6	0.2	57,272	60.4	0.2	3,631	55.3	0.9	91,532	64.6	0.2
Oral Cavity & Pharynx	24,905	66.3	0.4	2,841	42.2	1.1	1,971	65.8	1.3	132	62.0	5.3	2,208	59.8	1.2
Esophagus	11,852	19.6	0.4	1,376	12.3	1.1	624	15.6	1.9	64	8.6	4.6	1,014	16.5	1.5
Stomach	14,235	24.6	0.4	2,247	25.4	1.1	2,666	36.4	1.1	159	24.3	3.9	3,075	26.8	1.0
Colon & Rectum	82,187	65.1	0.2	9,616	55.5	0.6	7,627	67.8	0.6	473	59.6	2.8	9,502	62.9	0.6
Liver & Intrahepatic Bile Duct	14,124	17.4	0.4	2,435	11.5	0.9	3,869	22.2	0.9	190	19.1	3.5	3,742	17.5	0.8
Pancreas	18,958	5.5	0.2	2,302	4.1	0.6	1,540	6.6	0.9	104	6.4	2.6	2,131	5.1	0.6
Larynx	9,559	67.5	0.6	1,713	58.4	1.4	414	68.4	2.6	28	37.5	12.4	1,022	65.3	1.8
Lung & Bronchus	107,503	15.3	0.1	14,375	12.9	0.3	7,862	16.0	0.5	475	12.1	1.9	7,358	14.0	0.5
Melanoma of the Skin	42,440	86.6	0.2	152	62.8	4.5	255	73.4	3.3	45	59.1	8.9	1,081	77.1	1.5
Prostate	262,340	93.7	0.1	41,502	91.6	0.2	15,481	94.1	0.2	789	89.9	1.3	28,242	92.6	0.2
Testis	12,851	95.9	0.2	377	88.1	1.8	447	92.0	1.6	94	98.3	1.7	2,647	93.1	0.5
Urinary Bladder	53,369	80.8	0.2	2,425	70.6	1.1	2,176	81.3	1.0	108	74.6	4.9	3,315	79.3	0.8
Kidney & Renal Pelvis	28,450	70.4	0.3	3,173	70.3	0.9	1,600	72.2	1.3	231	65.6	3.7	4,053	69.8	0.8
Brain & Nervous System	14,297	32.7	0.4	937	35.7	1.8	791	38.2	2.0	60	47.6	7.8	2,144	44.1	1.2
Thyroid	8,825	93.6	0.3	498	93.1	1.3	867	94.1	0.9	37	86.4	5.7	1,185	92.6	0.9
Hodgkin Lymphoma	6,204	85.6	0.5	756	79.7	1.8	342	84.2	2.3	-	-	-	1,164	84.0	1.2
Non-Hodgkin Lymphoma	37,343	67.7	0.3	3,298	60.1	1.0	2,649	63.9	1.1	147	67.3	4.3	4,858	63.5	0.8
Myeloma	9,994	45.4	0.7	2,047	44.9	1.5	614	49.1	2.7	52	54.2	7.8	1,380	47.7	1.8
Leukemia	25,954	57.0	0.4	2,114	52.1	1.3	1,618	53.3	1.4	113	59.0	5.2	3,911	59.0	0.9
Mesothelioma	2,888	5.9	0.6	127	14.1	3.7	90	11.1	3.7	-	-	-	297	8.3	2.0

Source: SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2001-2007 with follow-up through 2008.

^b Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^c Hispanics are not mutually exclusive from whites, blacks, Asian/Pacific Islanders and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic could not be calculated due to less than 25 cases in the time interval.

Table 34.3
Five-Year Cause-Specific Survival^a(%) for Malignant Cancers, 2001-2007
Females by Race/Ethnicity and Primary Site

Primary Site	White			Black			Asian/ Pacific Islander			American Indian/ Alaska Native ^b			Hispanic ^c		
	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE
All Sites	782,973	65.9	0.1	86,642	56.5	0.2	62,024	68.4	0.2	3,956	60.8	0.9	93,645	66.7	0.2
Oral Cavity & Pharynx	10,749	67.0	0.5	1,266	57.4	1.6	1,144	71.1	1.6	60	70.3	6.3	1,062	67.6	1.8
Esophagus	3,338	19.8	0.8	580	17.5	1.9	193	18.5	3.4	26	-	-	249	19.5	3.0
Stomach	8,813	29.0	0.6	1,796	32.9	1.3	2,033	36.6	1.2	93	27.6	5.2	2,431	27.9	1.1
Colon & Rectum	78,901	63.9	0.2	11,075	58.1	0.6	7,020	68.9	0.7	497	63.1	2.5	8,374	63.7	0.6
Liver & Intrahepatic Bile Duct	5,725	15.8	0.6	900	16.3	1.6	1,666	21.0	1.3	77	13.6	4.6	1,510	17.9	1.3
Pancreas	19,328	6.0	0.2	2,887	6.1	0.6	1,574	8.0	0.8	111	7.6	2.9	2,377	7.5	0.7
Larynx	2,382	64.6	1.2	419	56.7	3.1	60	62.9	7.9	-	-	-	153	74.6	4.0
Lung & Bronchus	96,161	20.1	0.2	10,864	16.6	0.5	5,414	22.6	0.7	409	12.4	2.0	5,899	18.9	0.6
Melanoma of the Skin	33,913	92.0	0.2	214	77.8	3.4	269	81.9	2.7	44	92.7	4.1	1,622	88.7	0.9
Breast	240,911	88.4	0.1	26,827	77.5	0.3	20,449	90.3	0.3	1,088	85.6	1.3	27,761	86.1	0.3
Cervix uteri	15,862	72.3	0.4	2,668	62.9	1.1	1,878	72.4	1.2	125	62.4	5.2	4,928	75.1	0.7
Corpus & Uterus, NOS	47,460	82.9	0.2	4,561	62.7	0.8	3,893	84.7	0.7	245	84.4	2.5	5,557	81.5	0.6
Ovary ^d	26,351	43.4	0.4	2,260	36.1	1.2	2,144	55.7	1.3	150	43.6	5.0	3,488	49.0	1.0
Urinary Bladder	17,418	74.6	0.4	1,372	57.4	1.6	699	74.3	1.9	29	60.0	9.5	1,181	69.1	1.5
Kidney & Renal Pelvis	17,796	71.5	0.4	2,138	71.6	1.2	967	71.1	1.7	165	72.1	3.8	2,764	73.6	1.0
Brain & Nervous System	11,317	35.6	0.5	891	45.7	1.9	674	44.5	2.2	49	41.5	8.7	1,841	46.8	1.3
Thyroid	28,533	97.4	0.1	2,167	96.3	0.4	3,603	96.8	0.3	159	98.7	0.9	5,370	97.3	0.3
Hodgkin Lymphoma	5,190	87.0	0.5	662	86.5	1.5	273	87.0	2.4	-	-	-	881	86.1	1.3
Non-Hodgkin Lymphoma	32,427	69.8	0.3	2,808	67.8	1.0	2,344	67.7	1.1	124	69.7	4.4	4,159	68.7	0.8
Myeloma	8,037	42.2	0.7	2,101	47.1	1.4	581	42.7	2.7	60	42.1	8.1	1,242	41.2	1.8
Leukemia	19,154	56.3	0.4	1,734	52.1	1.4	1,245	53.1	1.6	96	54.7	6.2	3,058	58.0	1.0
Mesothelioma	875	13.8	1.5	58	17.9	6.9	31	20.1	7.9	-	-	-	122	14.4	4.8

Source: SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2001-2007 with follow-up through 2008.

^b Data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^c Hispanics are not mutually exclusive from whites, blacks, Asian/Pacific Islanders and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

^d Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

- Statistic could not be calculated due to less than 25 cases in the time interval.

Table 34.4

Cancer of the Female Breast

5-Year Cause-Specific Survival^a and Case Distribution by Race/Ethnicity
SEER 17 Registries, 2001-2007

Race/Ethnicity	All Ages				Ages <50				Ages 50+			
	Case Distribution		Survival		Case Distribution		Survival		Case Distribution		Survival	
	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE
All Races	291,619	100.0	87.5	0.1	73,512	100.0	88.7	0.1	218,107	100.0	87.1	0.1
White	240,911	82.6	88.4	0.1	56,699	77.1	90.0	0.2	184,212	84.5	87.9	0.1
White Non-Hispanic	214,131	73.4	88.6	0.1	46,628	63.4	90.8	0.2	167,503	76.8	88.0	0.1
White Hispanic	26,780	9.2	86.2	0.3	10,071	13.7	86.1	0.4	16,709	7.7	86.3	0.3
Black	26,827	9.2	77.5	0.3	8,696	11.8	77.8	0.5	18,131	8.3	77.4	0.4
American Indian/Alaska Native ^c	1,088	0.4	85.6	1.3	384	0.5	84.3	2.3	704	0.3	86.2	1.6
Asian	18,909	6.5	90.7	0.3	6,558	8.9	91.5	0.4	12,351	5.7	90.2	0.3
Asian Indian, Pakistani	1,439	0.5	89.3	1.0	561	0.8	91.0	1.5	878	0.4	88.1	1.4
Chinese	3,778	1.3	90.4	0.6	1,412	1.9	91.6	0.9	2,366	1.1	89.6	0.8
Filipino	5,248	1.8	89.0	0.5	1,653	2.2	90.0	0.9	3,595	1.6	88.6	0.7
Japanese	3,208	1.1	93.0	0.5	693	0.9	94.1	1.1	2,515	1.2	92.6	0.6
Korean	1,348	0.5	89.6	1.0	607	0.8	89.0	1.6	741	0.3	90.0	1.4
Vietnamese	1,150	0.4	91.1	1.1	508	0.7	90.8	1.6	642	0.3	91.3	1.4
Other Asian ^d	2,738	0.9	92.7	0.6	1,124	1.5	94.3	0.8	1,614	0.7	91.6	0.9
Pacific Islander	1,540	0.5	85.4	1.1	468	0.6	83.9	2.1	1,072	0.5	86.2	1.3
Hawaiian	1,048	0.4	88.2	1.2	291	0.4	88.6	2.2	757	0.3	88.1	1.4
Other Pacific Islander ^e	492	0.2	78.9	2.3	177	0.2	75.6	4.0	315	0.1	80.9	2.8
Hispanic ^f	27,761	9.5	86.1	0.3	10,428	14.2	86.0	0.4	17,333	7.9	86.3	0.3

Source: SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2001-2007 with follow-up through 2008.

^b Percent of cases do not sum to 100 because All Races includes unknown race and other race which are not shown separately.

^c Survival estimates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^d Other Asians include Laotian, Hmong, Kampuchean, Thai, and Asians, Not Otherwise Specified (NOS).

^e Other Pacific Islanders include Micronesian, Chamorroan, Guamanian, Polynesian, Tahitian, Samoan, Tongan, Melanesian, Fiji Islander, New Guinean, and Pacific Islander, NOS.

^f Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic not shown due to fewer than 25 cases during the time period.

Table 34.5

Cancer of the Colon and Rectum

5-Year Cause-Specific Survival^a and Case Distribution by Race/Ethnicity
SEER 17 Registries, 2001-2007

Race/Ethnicity	Both Sexes				Males				Females			
	Case Distribution		Survival		Case Distribution		Survival		Case Distribution		Survival	
	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE
All Races	199,078	100.0	64.2	0.1	100,833	100.0	64.5	0.2	98,245	100.0	63.8	0.2
White	161,088	80.9	64.5	0.1	82,187	81.5	65.1	0.2	78,901	80.3	63.9	0.2
White Non-Hispanic	143,825	72.2	64.6	0.1	72,976	72.4	65.4	0.2	70,849	72.1	63.9	0.2
White Hispanic	17,263	8.7	63.2	0.5	9,211	9.1	62.7	0.6	8,052	8.2	63.7	0.7
Black	20,691	10.4	56.9	0.4	9,616	9.5	55.5	0.6	11,075	11.3	58.1	0.6
American Indian/Alaska Native ^c	970	0.5	61.5	1.9	473	0.5	59.6	2.8	497	0.5	63.1	2.5
Asian	13,828	6.9	68.6	0.5	7,143	7.1	68.1	0.7	6,685	6.8	69.1	0.7
Asian Indian, Pakistani	611	0.3	71.3	2.2	345	0.3	73.4	2.9	266	0.3	68.5	3.5
Chinese	3,321	1.7	70.2	0.9	1,708	1.7	69.9	1.3	1,613	1.6	70.6	1.4
Filipino	2,926	1.5	65.6	1.1	1,548	1.5	65.2	1.5	1,378	1.4	66.2	1.5
Japanese	3,048	1.5	68.4	1.0	1,528	1.5	67.6	1.4	1,520	1.5	69.2	1.4
Korean	1,272	0.6	66.9	1.6	650	0.6	65.8	2.2	622	0.6	67.9	2.3
Vietnamese	1,051	0.5	65.3	1.8	551	0.5	62.7	2.6	500	0.5	68.3	2.5
Other Asian ^d	1,599	0.8	73.4	1.3	813	0.8	74.4	1.8	786	0.8	72.4	1.9
Pacific Islander	819	0.4	64.0	2.0	484	0.5	63.3	2.6	335	0.3	64.9	3.1
Hawaiian	562	0.3	65.0	2.4	344	0.3	62.7	3.1	218	0.2	68.2	3.7
Other Pacific Islander ^e	257	0.1	62.0	3.6	140	0.1	64.8	4.6	117	0.1	58.0	5.6
Hispanic ^f	17,876	9.0	63.3	0.4	9,502	9.4	62.9	0.6	8,374	8.5	63.7	0.6

Source: SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2001-2007 with follow-up through 2008.

^b Percent of cases do not sum to 100 because All Races includes unknown race and other race which are not shown separately.

^c Survival estimates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^d Other Asians include Laotian, Hmong, Kampuchean, Thai, and Asians, Not Otherwise Specified (NOS).

^e Other Pacific Islanders include Micronesian, Chamorro, Guamanian, Polynesian, Tahitian, Samoan, Tongan, Melanesian, Fiji Islander, New Guinean, and Pacific Islander, NOS.

^f Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic not shown due to fewer than 25 cases during the time period.

Table 34.6

Cancer of the Liver and Intrahepatic Bile Duct

5-Year Cause-Specific Survival^a and Case Distribution by Race/Ethnicity
SEER 17 Registries, 2001-2007

Race/Ethnicity	Both Sexes				Males				Females			
	Case Distribution		Survival		Case Distribution		Survival		Case Distribution		Survival	
	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE
All Races	29,188	100.0	17.5	0.3	20,757	100.0	17.7	0.3	8,431	100.0	16.9	0.5
White	19,849	68.0	16.9	0.3	14,124	68.0	17.4	0.4	5,725	67.9	15.8	0.6
White Non-Hispanic	14,751	50.5	16.8	0.4	10,482	50.5	17.4	0.5	4,269	50.6	15.4	0.7
White Hispanic	5,098	17.5	17.5	0.7	3,642	17.5	17.6	0.9	1,456	17.3	17.3	1.3
Black	3,335	11.4	12.9	0.8	2,435	11.7	11.5	0.9	900	10.7	16.3	1.6
American Indian/Alaska Native ^c	267	0.9	17.4	2.8	190	0.9	19.1	3.5	77	0.9	13.6	4.6
Asian	5,269	18.1	22.3	0.8	3,666	17.7	22.8	0.9	1,603	19.0	21.2	1.4
Asian Indian, Pakistani	143	0.5	24.5	4.6	103	0.5	24.8	5.9	40	0.5	21.7	7.2
Chinese	1,348	4.6	24.8	1.5	953	4.6	25.7	1.8	395	4.7	22.4	2.7
Filipino	794	2.7	19.6	2.0	572	2.8	20.5	2.3	222	2.6	17.7	3.7
Japanese	539	1.8	22.0	2.3	258	1.2	26.0	3.3	281	3.3	18.9	3.0
Korean	700	2.4	21.9	2.1	469	2.3	19.1	2.4	231	2.7	28.8	3.7
Vietnamese	1,080	3.7	22.6	1.8	832	4.0	23.8	2.0	248	2.9	18.6	3.5
Other Asian ^d	665	2.3	21.2	1.9	479	2.3	20.8	2.2	186	2.2	21.9	3.9
Pacific Islander	266	0.9	11.9	2.7	203	1.0	10.9	3.0	63	0.7	14.7	6.0
Hawaiian	150	0.5	10.0	3.2	109	0.5	9.5	3.8	41	0.5	10.0	6.0
Other Pacific Islander ^e	116	0.4	14.8	4.7	94	0.5	14.3	4.2	-	-	-	-
Hispanic ^f	5,252	18.0	17.6	0.7	3,742	18.0	17.5	0.8	1,510	17.9	17.9	1.3

Source: SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2001-2007 with follow-up through 2008.

^b Percent of cases do not sum to 100 because All Races includes unknown race and other race which are not shown separately.

^c Survival estimates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^d Other Asians include Laotian, Hmong, Kampuchean, Thai, and Asians, Not Otherwise Specified (NOS).

^e Other Pacific Islanders include Micronesian, Chamorro, Guamanian, Polynesian, Tahitian, Samoan, Tongan, Melanesian, Fiji Islander, New Guinean, and Pacific Islander, NOS.

^f Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic not shown due to fewer than 25 cases during the time period.

Table 34.7

Cancer of the Lung and Bronchus

5-Year Cause-Specific Survival^a and Case Distribution by Race/Ethnicity
SEER 17 Registries, 2001-2007

Race/Ethnicity	Both Sexes				Males				Females			
	Case Distribution		Survival		Case Distribution		Survival		Case Distribution		Survival	
	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE
All Races	243,894	100.0	17.3	0.1	130,657	100.0	15.1	0.1	113,237	100.0	19.9	0.1
White	203,664	83.5	17.6	0.1	107,503	82.3	15.3	0.1	96,161	84.9	20.1	0.2
White Non-Hispanic	190,869	78.3	17.7	0.1	100,410	76.9	15.4	0.1	90,459	79.9	20.1	0.2
White Hispanic	12,795	5.2	16.2	0.4	7,093	5.4	14.0	0.5	5,702	5.0	19.0	0.7
Black	25,239	10.3	14.5	0.3	14,375	11.0	12.9	0.3	10,864	9.6	16.6	0.5
American Indian/Alaska Native ^c	884	0.4	12.1	1.3	475	0.4	12.1	1.9	409	0.4	12.4	2.0
Asian	12,141	5.0	18.9	0.5	7,208	5.5	16.0	0.6	4,933	4.4	23.0	0.8
Asian Indian, Pakistani	467	0.2	25.3	2.5	315	0.2	24.2	2.9	152	0.1	27.8	4.9
Chinese	3,159	1.3	19.5	0.9	1,829	1.4	16.5	1.1	1,330	1.2	23.6	1.5
Filipino	3,156	1.3	18.9	0.9	1,987	1.5	15.2	1.0	1,169	1.0	24.7	1.7
Japanese	2,019	0.8	16.6	1.0	1,088	0.8	14.2	1.3	931	0.8	19.4	1.6
Korean	931	0.4	16.1	1.5	520	0.4	14.0	1.9	411	0.4	18.5	2.5
Vietnamese	1,302	0.5	18.5	1.4	848	0.6	15.9	1.7	454	0.4	23.3	2.6
Other Asian ^d	1,107	0.5	22.2	1.6	621	0.5	18.1	2.2	486	0.4	26.7	2.5
Pacific Islander	1,135	0.5	17.0	1.4	654	0.5	15.7	1.8	481	0.4	18.7	2.1
Hawaiian	807	0.3	16.5	1.6	446	0.3	14.1	2.1	361	0.3	19.2	2.4
Other Pacific Islander ^e	328	0.1	18.7	2.8	208	0.2	20.0	3.4	120	0.1	16.3	4.7
Hispanic ^f	13,257	5.4	16.2	0.4	7,358	5.6	14.0	0.5	5,899	5.2	18.9	0.6

Source: SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2001-2007 with follow-up through 2008.

^b Percent of cases do not sum to 100 because All Races includes unknown race and other race which are not shown separately.

^c Survival estimates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^d Other Asians include Laotian, Hmong, Kampuchean, Thai, and Asians, Not Otherwise Specified (NOS).

^e Other Pacific Islanders include Micronesian, Chamorroan, Guamanian, Polynesian, Tahitian, Samoan, Tongan, Melanesian, Fiji Islander, New Guinean, and Pacific Islander, NOS.

^f Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic not shown due to fewer than 25 cases during the time period.

Table 34.8

Cancer of the Prostate

5-Year Cause-Specific Survival^a and Case Distribution by Race/Ethnicity
SEER 17 Registries, 2001-2007

Race/Ethnicity	All Ages				Ages <65				Ages 65+			
	Case Distribution		Survival		Case Distribution		Survival		Case Distribution		Survival	
	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE
All Races	330,673	100.0	93.5	0.1	132,507	100.0	96.7	0.1	198,166	100.0	91.4	0.1
White	262,340	79.3	93.7	0.1	103,126	77.8	96.9	0.1	159,214	80.3	91.6	0.1
White Non-Hispanic	235,493	71.2	93.8	0.1	92,588	69.9	97.2	0.1	142,905	72.1	91.6	0.1
White Hispanic	26,847	8.1	92.6	0.2	10,538	8.0	95.0	0.3	16,309	8.2	91.0	0.3
Black	41,502	12.6	91.6	0.2	20,944	15.8	95.0	0.2	20,558	10.4	88.0	0.3
American Indian/Alaska Native ^c	789	0.2	89.9	1.3	329	0.2	93.2	1.7	460	0.2	87.5	1.9
Asian	14,505	4.4	94.5	0.2	4,272	3.2	97.3	0.3	10,233	5.2	93.3	0.3
Asian Indian, Pakistani	1,219	0.4	95.1	0.8	490	0.4	97.8	0.9	729	0.4	93.2	1.1
Chinese	3,202	1.0	94.6	0.5	737	0.6	97.8	0.7	2,465	1.2	93.6	0.6
Filipino	4,101	1.2	93.5	0.5	1,370	1.0	96.2	0.6	2,731	1.4	92.1	0.6
Japanese	2,820	0.9	94.7	0.5	614	0.5	98.2	0.7	2,206	1.1	93.7	0.6
Korean	661	0.2	94.3	1.2	164	0.1	96.3	1.6	497	0.3	93.5	1.5
Vietnamese	775	0.2	94.0	1.1	259	0.2	96.7	1.5	516	0.3	92.6	1.4
Other Asian ^d	1,727	0.5	96.1	0.6	638	0.5	98.6	0.6	1,089	0.5	94.6	0.8
Pacific Islander	976	0.3	88.5	1.2	341	0.3	91.2	1.7	635	0.3	86.8	1.7
Hawaiian	608	0.2	87.9	1.6	206	0.2	91.6	2.1	402	0.2	85.7	2.2
Other Pacific Islander ^e	368	0.1	89.6	1.9	135	0.1	90.7	2.9	233	0.1	89.0	2.5
Hispanic ^f	28,242	8.5	92.6	0.2	11,047	8.3	95.0	0.3	17,195	8.7	91.1	0.3

Source: SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2001-2007 with follow-up through 2008.

^b Percent of cases do not sum to 100 because All Races includes unknown race and other race which are not shown separately.

^c Survival estimates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^d Other Asians include Laotian, Hmong, Kampuchean, Thai, and Asians, Not Otherwise Specified (NOS).

^e Other Pacific Islanders include Micronesian, Chamorroan, Guamanian, Polynesian, Tahitian, Samoan, Tongan, Melanesian, Fiji Islander, New Guinean, and Pacific Islander, NOS.

^f Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic not shown due to fewer than 25 cases during the time period.