

Table 1.21  
U.S. Prevalence Counts, Invasive Cancers Only, January 1, 2008<sup>a</sup>  
Using Different Tumor Inclusion Criteria<sup>b</sup>

Site	Sex	5-Year Limited Duration			33-year Limited Duration	
		1st Invasive Tumor Ever <sup>c</sup>	1st Per Site in Previous 33 Years <sup>d</sup>	1st Per Site in Previous 5 Years <sup>e</sup>	1st Invasive Tumor Ever <sup>c</sup>	1st Per Site in Previous 33 Years <sup>d</sup>
All Sites	Both Sexes	4,249,706	4,331,842	4,734,286	11,688,972	11,892,768
	Male	2,188,126	2,219,768	2,413,895	5,506,074	5,570,768
	Female	2,061,580	2,112,074	2,320,391	6,182,898	6,322,000
Oral Cavity & Pharynx	Both Sexes	94,344	107,847	111,563	245,098	269,441
	Male	65,141	73,776	76,117	159,807	174,074
	Female	29,203	34,071	35,446	85,291	95,367
Esophagus	Both Sexes	19,829	23,939	23,951	30,597	35,817
	Male	15,229	18,405	18,417	23,362	27,236
	Female	4,600	5,534	5,534	7,235	8,581
Stomach	Both Sexes	33,089	39,300	39,521	65,498	74,863
	Male	19,927	23,973	24,073	37,436	43,123
	Female	13,162	15,327	15,448	28,062	31,740
Colon & Rectum	Both Sexes	407,014	469,977	478,669	1,098,089	1,223,141
	Male	205,716	237,703	241,917	540,332	598,652
	Female	201,298	232,274	236,752	557,757	624,489
Liver & Intrahepatic Bile Duct	Both Sexes	23,568	26,629	26,633	31,341	34,942
	Male	17,221	19,339	19,343	21,890	24,282
	Female	6,347	7,290	7,290	9,451	10,660
Pancreas	Both Sexes	26,137	31,728	31,739	34,955	41,506
	Male	12,950	15,750	15,761	16,972	20,161
	Female	13,187	15,978	15,978	17,983	21,345
Larynx	Both Sexes	30,287	36,587	36,906	87,991	98,455
	Male	24,152	29,051	29,312	70,449	78,408
	Female	6,135	7,536	7,594	17,542	20,047
Lung & Bronchus	Both Sexes	220,060	276,893	283,557	371,171	450,808
	Male	103,405	130,714	133,454	171,338	207,978
	Female	116,655	146,179	150,103	199,833	242,830
Melanoma of the Skin	Both Sexes	252,206	283,584	294,513	796,296	855,750
	Male	133,991	153,602	160,698	394,107	428,125
	Female	118,215	129,982	133,815	402,189	427,625
Breast	Female	805,515	868,236	918,683	2,606,497	2,763,127
Cervix	Female	38,294	40,160	40,257	200,357	206,078
Corpus & Uterus, NOS	Female	154,529	174,300	174,371	541,737	589,947
Ovary <sup>f</sup>	Female	54,970	63,837	63,889	161,711	181,687

<sup>a</sup> U.S. 2008 cancer prevalence counts are based on 2008 cancer prevalence proportions from the SEER 9 registries and 1/1/2008 U.S. population estimates based on the average of 2007 and 2008 population estimates from the U.S. Bureau of the Census.

<sup>b</sup> Prevalence estimates are ambiguous for those with multiple cancers, unless the tumor inclusion criteria are understood. Depending on the application, different inclusion criteria may be appropriate. This table provides three different methods of tumor inclusion:

<sup>c</sup> (c) First invasive tumor ever

<sup>d</sup> (d) First invasive tumor for each cancer site diagnosed during the previous 33 years (1975-2007)

<sup>e</sup> (e) First invasive tumor for each cancer site diagnosed during the previous 5 years (2003-2007)

For definitions (d) and (e) all sites is treated as a separate cancer "site".

Consider a woman who had three invasive cancers: Melanoma in 1981; Breast cancer in 2003; Melanoma in 2004.

In method (c) the melanoma is the woman's first cancer, and is counted for the melanoma and all sites 33-year limited duration prevalence. For 5-year limited duration prevalence, the woman is not counted at all since her first cancer occurred more than 5 years prior to 1/1/2008.

In method (d) the 1981 melanoma is counted for the melanoma and all sites 33-year limited duration prevalence. The 2003 breast cancer is counted for the breast 5-year and 33-year limited duration prevalence.

In method (e) the 2003 breast cancer is counted for the breast cancer and all sites 5-year limited duration prevalence. The 2004 melanoma is counted for 5-year limited duration prevalence for melanoma.

<sup>f</sup> Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

Table 1.21 - continued  
 U.S. Prevalence Counts, Invasive Cancers Only, January 1, 2008<sup>a</sup>  
 Using Different Tumor Inclusion Criteria<sup>b</sup>

Site	Sex	5-Year Limited Duration			33-year Limited Duration	
		1st Invasive Tumor Ever <sup>c</sup>	1st Per Site in Previous 33 Years <sup>d</sup>	1st Per Site in Previous 5 Years <sup>e</sup>	1st Invasive Tumor Ever <sup>c</sup>	1st Per Site in Previous 33 Years <sup>d</sup>
Prostate	Male	970,359	1,048,069	1,048,105	2,396,871	2,555,936
Testis	Male	40,125	40,828	41,397	187,670	190,120
Urinary Bladder	Both Sexes	194,097	239,746	244,623	531,166	614,655
	Male	146,585	181,724	185,560	395,025	456,695
	Female	47,512	58,022	59,063	136,141	157,960
Kidney & Renal Pelvis	Both Sexes	128,486	155,791	157,332	290,688	337,899
	Male	77,495	95,049	96,135	171,540	201,021
	Female	50,991	60,742	61,197	119,148	136,878
Brain & Nervous System	Both Sexes	40,898	43,394	43,831	116,955	120,623
	Male	22,693	24,140	24,351	62,642	64,690
	Female	18,205	19,254	19,480	54,313	55,933
Thyroid	Both Sexes	139,961	153,689	154,081	425,181	450,805
	Male	31,088	35,530	35,602	93,227	100,569
	Female	108,873	118,159	118,479	331,954	350,236
Hodgkin Lymphoma	Both Sexes	36,835	39,099	39,099	155,837	159,846
	Male	19,688	20,808	20,808	80,709	82,717
	Female	17,147	18,291	18,291	75,128	77,129
Non-Hodgkin Lymphoma	Both Sexes	189,410	220,661	222,116	452,242	502,943
	Male	98,768	116,412	117,193	234,240	260,923
	Female	90,642	104,249	104,923	218,002	242,020
Myeloma	Both Sexes	42,726	49,682	49,741	65,682	74,814
	Male	23,046	27,105	27,164	36,054	41,454
	Female	19,680	22,577	22,577	29,628	33,360
Leukemia	Both Sexes	103,646	117,597	117,772	252,119	274,930
	Male	59,538	68,094	68,189	142,621	155,942
	Female	44,108	49,503	49,583	109,498	118,988
Acute Lymphocytic Leukemia	Both Sexes	15,402	15,740	15,740	59,985	60,486
	Male	8,796	8,940	8,940	33,127	33,297
	Female	6,606	6,800	6,800	26,858	27,189
Childhood (Ages 0-19)	Both Sexes	59,739	59,851	60,342	277,294	277,798
	Male	31,980	32,036	32,261	142,787	143,010
	Female	27,759	27,815	28,081	134,507	134,788
Kaposi Sarcoma	Both Sexes	6,661	7,109	7,109	22,566	23,720
	Male	6,221	6,561	6,561	21,226	22,190
	Female	440	548	548	1,340	1,530
Mesothelioma	Both Sexes	3,148	3,988	3,988	4,609	5,619
	Male	2,120	2,735	2,735	2,681	3,419
	Female	1,028	1,253	1,253	1,928	2,200

<sup>a</sup> U.S. 2008 cancer prevalence counts are based on 2008 cancer prevalence proportions from the SEER 9 registries and 1/1/2008 U.S. population estimates based on the average of 2007 and 2008 population estimates from the U.S. Bureau of the Census.

<sup>b</sup> Prevalence estimates are ambiguous for those with multiple cancers, unless the tumor inclusion criteria are understood. Depending on the application, different inclusion criteria may be appropriate. This table provides three different methods of tumor inclusion:

<sup>c</sup> (c) First invasive tumor ever

<sup>d</sup> (d) First invasive tumor for each cancer site diagnosed during the previous 33 years (1975-2007)

<sup>e</sup> (e) First invasive tumor for each cancer site diagnosed during the previous 5 years (2003-2007)

For definitions (d) and (e) all sites is treated as a separate cancer "site".

Consider a woman who had three invasive cancers: Melanoma in 1981; Breast cancer in 2003; Melanoma in 2004.

In method (c) the melanoma is the woman's first cancer, and is counted for the melanoma and all sites 33-year limited duration prevalence. For 5-year limited duration prevalence, the woman is not counted at all since her first cancer occurred more than 5 years prior to 1/1/2008.

In method (d) the 1981 melanoma is counted for the melanoma and all sites 33-year limited duration prevalence. The 2003 breast cancer is counted for the breast 5-year and 33-year limited duration prevalence.

In method (e) the 2003 breast cancer is counted for the breast cancer and all sites 5-year limited duration prevalence. The 2004 melanoma is counted for 5-year limited duration prevalence for melanoma.

Table 1.22  
U.S. Complete Prevalence Counts, Invasive Cancers Only, January 1, 2008<sup>a</sup>  
By Age at Prevalence

Site/Sex	Age at Prevalence								
	All Ages <sup>c</sup>	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70+
All Sites									
Males	5,505,862	17,369	39,603	76,578	144,636	338,114	771,402	1,362,003	2,756,156
Females	6,451,737	14,868	33,960	79,574	212,422	605,640	1,175,930	1,493,293	2,836,051
Oral Cavity & Pharynx									
Males	164,159	55	433	1,463	3,321	14,386	39,210	45,877	59,413
Females	89,006	126	505	1,305	3,672	8,101	16,557	20,190	38,550
Esophagus									
Males	23,026	0	0	14	129	1,213	4,278	7,743	9,649
Females	7,178	0	0	0	49	289	930	1,818	4,092
Stomach									
Males	37,739	0	5	76	609	2,419	6,015	9,360	19,256
Females	28,271	4	11	140	619	1,985	4,077	4,970	16,465
Colon & Rectum									
Males	542,127	11	23	1,058	5,201	23,913	72,884	124,471	314,566
Females	567,950	0	102	1,290	4,857	21,841	61,776	101,891	376,192
Liver & Intrahep									
Males	21,567	413	537	414	426	1,632	7,439	5,891	4,815
Females	9,608	441	402	368	362	742	2,030	2,163	3,101
Pancreas									
Males	16,811	0	11	78	320	1,324	3,510	5,094	6,474
Females	17,846	0	58	117	413	1,288	3,104	4,215	8,651
Larynx									
Males	71,273	0	0	44	196	2,385	10,173	20,066	38,410
Females	17,668	0	0	41	140	1,042	3,056	4,519	8,870
Lung & Bronchus									
Males	173,428	44	70	377	1,119	5,823	23,597	51,308	91,089
Females	200,061	22	31	382	1,355	7,904	25,460	55,404	109,504
Melanoma of the Skin									
Males	400,706	92	701	4,870	16,596	46,265	87,532	102,829	141,819
Females	422,064	71	1,048	10,567	32,200	70,227	99,340	89,638	118,973

<sup>a</sup> U.S. 2008 cancer prevalence counts are based on 2008 cancer prevalence proportions from the SEER 9 registries (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta) and 1/1/2008 U.S. population estimates based on the average of 2007 and 2008 population estimates from the U.S. Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b</sup> Cases diagnosed more than 33 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

<sup>c</sup> Due to rounding, the sum of the age specific estimates may not equal the all ages estimate.

Table 1.22 - continued  
 U.S. Complete Prevalence Counts, Invasive Cancers Only, January 1, 2008<sup>a</sup>  
 By Age at Prevalence

Site/Sex	Age at Prevalence								
	All Ages <sup>c</sup>	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70+
Breast									
Males	13,616	0	0	16	81	576	1,940	3,622	7,381
Females	2,632,005	0	84	2,326	34,635	207,154	507,534	689,283	1,190,989
Cervix									
Females	243,884	0	56	2,010	16,779	42,631	57,933	50,513	73,961
Corpus & Uterus, NOS									
Females	573,300	0	54	399	5,192	23,450	79,633	138,800	325,772
Ovary <sup>d</sup>									
Females	177,578	82	863	3,458	6,583	19,450	38,659	42,317	66,166
Prostate									
Males	2,355,464	50	68	77	361	18,125	205,191	635,592	1,496,001
Urinary Bladder									
Males	398,329	22	119	672	2,397	12,099	41,421	91,664	249,934
Females	139,099	44	44	236	1,063	4,381	12,953	28,380	91,998
Kidney & Renal Pelvis									
Males	174,350	1,412	2,368	2,440	4,392	15,279	34,772	48,233	65,453
Females	121,724	1,339	2,604	2,584	4,202	11,129	20,525	28,201	51,141
Hodgkin Lymphoma									
Males	86,218	192	2,181	8,725	15,958	22,961	19,083	10,919	6,197
Females	80,558	99	1,575	9,569	16,034	22,130	16,962	8,071	6,118
Non-Hodgkin Lymphoma									
Males	235,433	723	3,328	6,821	11,902	27,303	46,564	57,135	81,657
Females	218,945	407	1,726	3,835	7,914	18,988	38,082	49,684	98,310
Myeloma									
Males	35,445	0	5	60	526	2,371	7,193	11,144	14,146
Females	29,170	0	0	32	210	1,855	5,358	8,290	13,424
Leukemia									
Males	142,702	6,224	11,976	10,886	9,714	11,892	19,280	27,233	45,497
Females	110,648	5,101	9,898	9,739	8,174	8,819	13,008	17,422	38,488
Acute Lymphocytic Leuk									
Males	34,306	5,184	10,430	8,380	5,501	2,691	1,104	643	374
Females	27,887	4,358	8,254	7,158	4,534	2,031	736	520	296

<sup>a</sup> U.S. 2008 cancer prevalence counts are based on 2008 cancer prevalence proportions from the SEER 9 registries (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta) and 1/1/2008 U.S. population estimates based on the average of 2007 and 2008 population estimates from the U.S. Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b</sup> Cases diagnosed more than 33 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

<sup>c</sup> Due to rounding, the sum of the age specific estimates may not equal the all ages estimate.

Table 2.21  
All Cancer Sites (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2008  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <33 <sup>e</sup>	>=33 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	4,132,758	2,851,221	1,878,782	1,129,972	658,827	413,357	9,795,922	11,478,794	478,805	11,957,599
	Males	2,121,106	1,419,006	899,769	444,824	220,318	123,455	4,857,657	5,407,975	97,887	5,505,862
	Females	2,011,652	1,432,215	979,013	685,148	438,509	289,902	4,938,265	6,070,819	380,918	6,451,737
White <sup>b</sup>	Both Sexes	3,552,919	2,502,741	1,667,443	1,025,739	599,931	378,080	8,540,079	10,073,552	422,039	10,495,591
	Males	1,815,492	1,236,957	795,345	407,506	202,679	114,503	4,213,126	4,720,180	91,527	4,811,707
	Females	1,737,427	1,265,784	872,098	618,233	397,252	263,577	4,326,953	5,353,372	330,512	5,683,884
Black <sup>b</sup>	Both Sexes	390,394	234,454	143,332	68,326	37,830	22,314	840,720	936,612	26,988	963,600
	Males	210,085	129,069	74,942	25,196	10,920	5,197	447,658	474,747	2,566	477,313
	Females	180,309	105,385	68,390	43,130	26,910	17,117	393,062	461,865	24,422	486,287
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	117,499	71,722	42,716	+	+	+	248,042	+	+	+
	Males	51,757	29,223	16,203	+	+	+	102,531	+	+	+
	Females	65,742	42,499	26,513	+	+	+	145,511	+	+	+
Hispanic <sup>d</sup>	Both Sexes	264,703	162,108	96,892	+	+	+	563,739	+	+	+
	Males	127,349	75,681	42,745	+	+	+	260,853	+	+	+
	Females	137,354	86,427	54,147	+	+	+	302,886	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2008, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	2.7119%	0.0789%	0.1632%	0.2690%	0.6819%	1.6511%	3.8386%	8.5534%	13.6175%	14.3510%	2.7635%
	Males	2.7122%	0.0839%	0.1727%	0.2544%	0.5250%	1.1291%	3.1627%	9.4144%	18.1814%	21.0990%	3.1491%
	Females	2.7117%	0.0736%	0.1533%	0.2845%	0.8435%	2.1749%	4.4853%	7.7733%	10.0184%	10.6162%	2.5395%
White <sup>c</sup>	Both Sexes	2.9529%	0.0850%	0.1778%	0.2992%	0.7570%	1.7435%	3.9685%	8.8923%	14.2663%	14.7234%	2.8859%
	Males	2.9217%	0.0908%	0.1891%	0.2835%	0.5945%	1.2063%	3.2350%	9.5835%	18.6768%	21.2550%	3.2330%
	Females	2.9843%	0.0790%	0.1658%	0.3164%	0.9318%	2.3027%	4.6947%	8.2482%	10.7059%	11.1238%	2.6942%
Black <sup>c</sup>	Both Sexes	1.9972%	0.0564%	0.1099%	0.1712%	0.4640%	1.3397%	3.7423%	9.0015%	13.2571%	13.1242%	2.6219%
	Males	2.2429%	0.0589%	0.1099%	0.1568%	0.3173%	0.9417%	3.8131%	12.4108%	20.8943%	22.8004%	3.5762%
	Females	1.7742%	0.0539%	0.1099%	0.1852%	0.5924%	1.6875%	3.6834%	6.3764%	8.0232%	8.6223%	2.0336%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	1.7873%	0.0627%	0.1259%	0.1673%	0.4219%	1.2419%	2.8185%	5.2979%	8.6994%	10.4282%	1.8661%
	Males	1.5886%	0.0662%	0.1370%	0.1511%	0.2623%	0.6568%	1.7373%	4.9396%	11.0274%	15.2947%	1.9117%
	Females	1.9735%	0.0590%	0.1142%	0.1835%	0.5729%	1.7777%	3.7589%	5.6029%	6.9589%	7.3696%	1.8895%
Hispanic <sup>d</sup>	Both Sexes	-	-	-	-	-	-	-	-	-	-	-
	Males	-	-	-	-	-	-	-	-	-	-	-
	Females	-	-	-	-	-	-	-	-	-	-	-

<sup>a</sup> US 2008 cancer prevalence counts are based on 2008 cancer prevalence proportions from the SEER registries and 1/1/2008 US population estimates based on the average of 2007 and 2008 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 33 years for 1975-2008 SEER 9 data; 18 years for 1990-2008 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 33 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <33 and >=33. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 3.21  
Brain and Other Nervous System (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2008  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <33 <sup>e</sup>	>=33 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	39,945	22,967	18,150	13,988	8,607	5,950	91,505	114,738	13,455	128,193
	Males	21,969	11,886	10,031	7,279	4,647	2,992	49,475	61,462	6,162	67,624
	Females	17,976	11,081	8,119	6,709	3,960	2,958	42,030	53,276	7,293	60,569
White <sup>b</sup>	Both Sexes	35,337	20,312	15,919	12,382	7,711	5,297	80,458	101,299	5,867	107,166
	Males	19,829	10,567	8,830	6,372	4,138	2,671	43,955	54,688	3,394 <sup>i</sup>	58,082
	Females	15,508	9,745	7,089	6,010	3,573	2,626	36,503	46,611	2,473 <sup>i</sup>	49,084
Black <sup>b</sup>	Both Sexes	2,891	1,728	1,407	1,117	621	480	7,002	8,725	+	+
	Males	1,295	915	729	620	364	227	3,490	4,373	+	+
	Females	1,596	813	678	497	257	253	3,512	4,352	+	+
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	1,146	600	507	+	+	+	2,440	+	+	+
	Males	555	303	250	+	+	+	1,214	+	+	+
	Females	591	297	257	+	+	+	1,226	+	+	+
Hispanic <sup>d</sup>	Both Sexes	4,197	2,640	1,602	+	+	+	9,310	+	+	+
	Males	2,334	1,358	802	+	+	+	4,944	+	+	+
	Females	1,863	1,282	800	+	+	+	4,366	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2008, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0257%	0.0135%	0.0294%	0.0231%	0.0243%	0.0318%	0.0321%	0.0318%	0.0242%	0.0147%	0.0257%
	Males	0.0279%	0.0141%	0.0316%	0.0240%	0.0265%	0.0346%	0.0357%	0.0357%	0.0272%	0.0157%	0.0278%
	Females	0.0235%	0.0130%	0.0271%	0.0221%	0.0219%	0.0290%	0.0287%	0.0283%	0.0219%	0.0142%	0.0236%
White <sup>c</sup>	Both Sexes	0.0287%	0.0147%	0.0328%	0.0256%	0.0278%	0.0364%	0.0359%	0.0348%	0.0263%	0.0156%	0.0287%
	Males	0.0314%	0.0155%	0.0359%	0.0264%	0.0308%	0.0395%	0.0392%	0.0399%	0.0296%	0.0179%	0.0312%
	Females	0.0261%	0.0138%	0.0295%	0.0247%	0.0247%	0.0333%	0.0326%	0.0300%	0.0236%	0.0144%	0.0262%
Black <sup>c</sup>	Both Sexes	0.0163%	0.0101%	0.0193%	0.0174%	0.0135%	0.0169%	0.0191%	0.0207%	0.0175%	0.0150%	0.0164%
	Males	0.0164%	0.0099%	0.0193%	0.0175%	0.0140%	0.0150%	0.0227%	0.0223%	0.0181%	-	0.0162%
	Females	0.0162%	0.0103%	0.0193%	0.0174%	0.0130%	0.0185%	0.0162%	0.0195%	0.0171%	0.0204%	0.0163%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0151%	0.0097%	0.0186%	0.0129%	0.0135%	0.0172%	0.0188%	0.0200%	0.0152%	0.0071%	0.0153%
	Males	0.0153%	0.0098%	0.0176%	0.0136%	0.0125%	0.0200%	0.0214%	0.0158%	0.0147%	-	0.0154%
	Females	0.0149%	0.0096%	0.0196%	0.0122%	0.0143%	0.0147%	0.0165%	0.0236%	0.0156%	0.0072%	0.0151%
Hispanic <sup>d</sup>	Both Sexes	-	-	-	-	-	-	-	-	-	-	-
	Males	-	-	-	-	-	-	-	-	-	-	-
	Females	-	-	-	-	-	-	-	-	-	-	-

<sup>a</sup> US 2008 cancer prevalence counts are based on 2008 cancer prevalence proportions from the SEER registries and 1/1/2008 US population estimates based on the average of 2007 and 2008 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 33 years for 1975-2008 SEER 9 data; 18 years for 1990-2008 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 33 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <33 and >=33. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 4.25  
Cancer of the Breast (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2008  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <33 <sup>e</sup>	>=33 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	800,302	665,332	449,100	306,246	170,735	97,423	2,135,402	2,572,603	73,018	2,645,621
	Males	5,477	3,417	1,917	1,062	738	394	11,712	13,420	196	13,616
	Females	794,825	661,915	447,183	305,184	169,997	97,029	2,123,690	2,559,183	72,822	2,632,005
White <sup>b</sup>	Both Sexes	691,124	589,133	401,493	278,454	155,119	88,902	1,874,734	2,273,204	64,753	2,337,957
	Males	4,680	2,929	1,712	949	676	368	10,063	11,653	9 <sup>i</sup>	11,662
	Females	686,444	586,204	399,781	277,505	154,443	88,534	1,864,671	2,261,551	64,744	2,326,295
Black <sup>b</sup>	Both Sexes	74,195	49,574	32,572	19,149	11,047	6,010	173,643	200,823	+	+
	Males	630	354	130	110	46	26	1,262	1,369	+	+
	Females	73,565	49,220	32,442	19,039	11,001	5,984	172,381	199,454	5,158	204,612
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	27,344	20,106	11,917	+	+	+	64,172	+	+	+
	Males	119	69	50	+	+	+	252	+	+	+
	Females	27,225	20,037	11,867	+	+	+	63,920	+	+	+
Hispanic <sup>d</sup>	Both Sexes	50,835	35,260	21,053	+	+	+	116,395	+	+	+
	Males	258	117	40	+	+	+	431	+	+	+
	Females	50,577	35,143	21,013	+	+	+	115,964	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2008, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.5908%	-	0.0002%	0.0055%	0.0835%	0.4464%	1.1167%	1.9788%	2.4655%	2.8427%	0.5895%
	Males	0.0066%	-	-	-	0.0004%	0.0027%	0.0087%	0.0244%	0.0430%	0.0569%	0.0077%
	Females	1.1652%	-	0.0004%	0.0112%	0.1691%	0.8917%	2.1769%	3.7494%	4.3758%	4.3846%	1.0815%
White <sup>c</sup>	Both Sexes	0.6442%	-	0.0001%	0.0050%	0.0834%	0.4462%	1.1456%	2.0863%	2.6395%	3.0204%	0.6163%
	Males	0.0071%	-	-	-	0.0003%	0.0025%	0.0082%	0.0240%	0.0454%	0.0610%	0.0078%
	Females	1.2832%	-	0.0003%	0.0103%	0.1726%	0.9079%	2.2718%	4.0083%	4.7337%	4.6515%	1.1434%
Black <sup>c</sup>	Both Sexes	0.4146%	-	-	0.0084%	0.0911%	0.4484%	1.0108%	1.7045%	1.9968%	2.4437%	0.5201%
	Males	0.0065%	-	-	-	-	0.0033%	0.0148%	0.0441%	0.0408%	0.0565%	0.0098%
	Females	0.7850%	-	-	0.0166%	0.1705%	0.8375%	1.8396%	2.9829%	3.3373%	3.5544%	0.8934%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.4480%	-	-	0.0060%	0.0777%	0.4391%	1.0148%	1.4717%	1.6779%	1.6701%	0.4541%
	Males	0.0038%	-	-	-	-	0.0019%	0.0058%	0.0120%	0.0273%	0.0262%	0.0045%
	Females	0.8642%	-	-	0.0115%	0.1506%	0.8395%	1.8924%	2.7140%	2.9119%	2.7033%	0.8220%
Hispanic <sup>d</sup>	Both Sexes	-	-	-	-	-	-	-	-	-	-	-
	Males	-	-	-	-	-	-	-	-	-	-	-
	Females	-	-	-	-	-	-	-	-	-	-	-

<sup>a</sup> US 2008 cancer prevalence counts are based on 2008 cancer prevalence proportions from the SEER registries and 1/1/2008 US population estimates based on the average of 2007 and 2008 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 33 years for 1975-2008 SEER 9 data; 18 years for 1990-2008 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 33 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <33 and >=33. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 5.15  
Cancer of the Cervix Uteri (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2008  
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <33 <sup>e</sup>	>=33 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>										
All Races <sup>b</sup>	38,079	36,825	35,079	30,973	24,009	20,597	128,098	196,451	47,433	243,884
White <sup>b</sup>	30,023	29,234	28,677	25,738	19,903	16,898	103,167	159,918	39,949	199,867
Black <sup>b</sup>	5,465	4,995	3,914	3,481	2,981	2,717	16,157	24,517	+	+
Asian/Pacific Islander <sup>c</sup>	1,988	1,723	1,677	+	+	+	6,128	+	+	+
Hispanic <sup>d</sup>	7,993	7,279	5,946	+	+	+	24,497	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2008, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>											
All Races <sup>c</sup>	0.0794%	-	0.0002%	0.0089%	0.0806%	0.1584%	0.1673%	0.1445%	0.1285%	0.0774%	0.0772%
White <sup>c</sup>	0.0807%	-	0.0003%	0.0099%	0.0919%	0.1703%	0.1636%	0.1335%	0.1153%	0.0613%	0.0784%
Black <sup>c</sup>	0.0696%	-	-	0.0075%	0.0626%	0.1262%	0.1675%	0.1652%	0.1486%	0.1470%	0.0742%
Asian/Pacific Islander <sup>c</sup>	0.0764%	-	-	0.0022%	0.0341%	0.1098%	0.1790%	0.1912%	0.1883%	0.1530%	0.0724%
Hispanic <sup>d</sup>	-	-	-	-	-	-	-	-	-	-	-

<sup>a</sup> US 2008 cancer prevalence counts are based on 2008 cancer prevalence proportions from the SEER registries and 1/1/2008 US population estimates based on the average of 2007 and 2008 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 33 years for 1975-2008 SEER 9 data; 18 years for 1990-2008 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 33 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <33 and >=33. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.



Table 6.28  
Cancer of the Colon and Rectum (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2008  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <33 <sup>e</sup>	>=33 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	405,303	270,741	166,502	114,834	70,032	38,374	915,005	1,078,730	31,347	1,110,077
	Males	203,619	135,867	81,515	56,282	32,296	15,683	457,112	531,058	11,069	542,127
	Females	201,684	134,874	84,987	58,552	37,736	22,691	457,893	547,672	20,278	567,950
White <sup>b</sup>	Both Sexes	341,574	235,193	145,609	101,365	62,575	34,312	782,775	928,164	27,784	955,948
	Males	174,185	119,126	71,669	50,297	28,945	14,085	395,488	461,862	9,942	471,804
	Females	167,389	116,067	73,940	51,068	33,630	20,227	387,287	466,302	17,842	484,144
Black <sup>b</sup>	Both Sexes	42,521	23,539	13,788	9,403	5,297	2,589	88,112	101,010	2,243	103,253
	Males	18,470	10,253	6,026	3,919	2,151	943	38,476	43,350	708	44,058
	Females	24,051	13,286	7,762	5,484	3,146	1,646	49,636	57,660	1,535	59,195
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	15,128	9,202	5,447	+	+	+	31,769	+	+	+
	Males	7,852	4,753	2,851	+	+	+	16,409	+	+	+
	Females	7,276	4,449	2,596	+	+	+	15,360	+	+	+
Hispanic <sup>d</sup>	Both Sexes	26,343	13,210	7,435	+	+	+	50,376	+	+	+
	Males	13,858	6,956	3,868	+	+	+	26,358	+	+	+
	Females	12,485	6,254	3,567	+	+	+	24,018	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2008, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.2568%	-	0.0002%	0.0049%	0.0238%	0.0965%	0.3139%	0.7469%	1.4385%	2.0905%	0.2632%
	Males	0.2609%	-	-	0.0046%	0.0242%	0.1014%	0.3489%	0.8839%	1.7230%	2.3760%	0.3054%
	Females	0.2527%	-	0.0004%	0.0051%	0.0235%	0.0916%	0.2805%	0.6228%	1.2141%	1.9325%	0.2303%
White <sup>c</sup>	Both Sexes	0.2675%	-	0.0003%	0.0049%	0.0234%	0.0931%	0.2990%	0.7276%	1.4408%	2.1100%	0.2602%
	Males	0.2716%	-	-	0.0047%	0.0238%	0.0963%	0.3371%	0.8681%	1.7268%	2.3951%	0.3029%
	Females	0.2633%	-	0.0004%	0.0052%	0.0229%	0.0897%	0.2612%	0.5966%	1.2099%	1.9528%	0.2260%
Black <sup>c</sup>	Both Sexes	0.2087%	-	-	0.0039%	0.0243%	0.1090%	0.3852%	0.9106%	1.4879%	2.0150%	0.2864%
	Males	0.1956%	-	-	0.0033%	0.0232%	0.1151%	0.4023%	1.0208%	1.6454%	2.0754%	0.3098%
	Females	0.2207%	-	-	0.0045%	0.0253%	0.1037%	0.3710%	0.8258%	1.3800%	1.9870%	0.2703%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.2381%	-	-	0.0038%	0.0244%	0.1000%	0.3259%	0.7174%	1.3527%	1.9526%	0.2532%
	Males	0.2530%	-	-	0.0041%	0.0252%	0.1137%	0.3530%	0.8544%	1.6866%	2.3414%	0.3017%
	Females	0.2241%	-	-	0.0036%	0.0237%	0.0874%	0.3022%	0.6008%	1.1030%	1.7082%	0.2167%
Hispanic <sup>d</sup>	Both Sexes	-	-	-	-	-	-	-	-	-	-	-
	Males	-	-	-	-	-	-	-	-	-	-	-
	Females	-	-	-	-	-	-	-	-	-	-	-

<sup>a</sup> US 2008 cancer prevalence counts are based on 2008 cancer prevalence proportions from the SEER registries and 1/1/2008 US population estimates based on the average of 2007 and 2008 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 33 years for 1975-2008 SEER 9 data; 18 years for 1990-2008 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 33 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <33 and >=33. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 7.15  
Cancer of the Corpus and Uterus, NOS (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2008  
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <33 <sup>e</sup>	>=33 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>										
All Races <sup>b</sup>	148,702	117,973	88,653	64,820	48,720	40,492	401,406	532,376	40,924	573,300
White <sup>b</sup>	130,665	107,199	81,673	60,487	46,122	38,387	361,167	484,757	35,491	520,248
Black <sup>b</sup>	10,430	5,474	3,701	2,465	1,393	942	22,098	25,952	999	26,951
Asian/Pacific Islander <sup>c</sup>	5,105	3,378	2,246	+	+	+	11,577	+	+	+
Hispanic <sup>d</sup>	10,082	6,144	4,013	+	+	+	22,241	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2008, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup> All Ages
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	
<u>Race</u>											
All Races <sup>c</sup>	0.2130%	-	-	0.0025%	0.0268%	0.1049%	0.3504%	0.7999%	0.9170%	0.7947%	0.1984%
White <sup>c</sup>	0.2410%	-	-	0.0022%	0.0265%	0.1050%	0.3699%	0.8742%	1.0234%	0.8741%	0.2149%
Black <sup>c</sup>	0.1013%	-	-	0.0022%	0.0218%	0.0647%	0.1745%	0.4926%	0.6049%	0.4628%	0.1205%
Asian/Pacific Islander <sup>c</sup>	0.1554%	-	-	0.0041%	0.0287%	0.1317%	0.3787%	0.5449%	0.4723%	0.3642%	0.1463%
Hispanic <sup>d</sup>	-	-	-	-	-	-	-	-	-	-	-

<sup>a</sup> US 2008 cancer prevalence counts are based on 2008 cancer prevalence proportions from the SEER registries and 1/1/2008 US population estimates based on the average of 2007 and 2008 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 33 years for 1975-2008 SEER 9 data; 18 years for 1990-2008 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 33 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <33 and >=33. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 8.21  
Cancer of the Esophagus (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2008  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <33 <sup>e</sup>	>=33 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	18,804	5,868	2,509	957	422	108	29,139	30,040	164	30,204
	Males	14,454	4,486	1,894	656	285	85	22,335	22,945	81	23,026
	Females	4,350	1,382	615	301	137	23	6,804	7,095	83	7,178
White <sup>b</sup>	Both Sexes	16,507	5,160	2,238	780	327	96	25,944	26,690	133	26,823
	Males	12,929	4,040	1,691	541	232	85	20,270	20,791	75	20,866
	Females	3,578	1,120	547	239	95	11	5,674	5,899	58	5,957
Black <sup>b</sup>	Both Sexes	1,769	521	209	139	91	11	2,378	2,515	38	2,553
	Males	1,119	308	153	90	53	0	1,448	1,525	10	1,535
	Females	650	213	56	49	38	11	930	990	28	1,018
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	339	117	45	+	+	+	525	+	+	+
	Males	278	76	30	+	+	+	403	+	+	+
	Females	61	41	15	+	+	+	122	+	+	+
Hispanic <sup>d</sup>	Both Sexes	726	170	93	+	+	+	1,052	+	+	+
	Males	607	143	68	+	+	+	881	+	+	+
	Females	119	27	25	+	+	+	171	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2008, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0078%	-	-	-	0.0004%	0.0028%	0.0112%	0.0315%	0.0465%	0.0345%	0.0080%
	Males	0.0120%	-	-	-	0.0006%	0.0045%	0.0188%	0.0535%	0.0771%	0.0633%	0.0135%
	Females	0.0036%	-	-	-	0.0002%	0.0011%	0.0039%	0.0116%	0.0224%	0.0185%	0.0034%
White <sup>c</sup>	Both Sexes	0.0088%	-	-	-	0.0005%	0.0032%	0.0122%	0.0340%	0.0504%	0.0364%	0.0086%
	Males	0.0137%	-	-	-	0.0008%	0.0052%	0.0205%	0.0585%	0.0841%	0.0673%	0.0147%
	Females	0.0039%	-	-	-	0.0002%	0.0011%	0.0040%	0.0112%	0.0232%	0.0193%	0.0035%
Black <sup>c</sup>	Both Sexes	0.0054%	-	-	-	-	0.0021%	0.0125%	0.0335%	0.0343%	0.0229%	0.0070%
	Males	0.0070%	-	-	-	-	0.0031%	0.0180%	0.0484%	0.0514%	0.0283%	0.0101%
	Females	0.0040%	-	-	-	-	0.0013%	0.0078%	0.0220%	0.0226%	0.0203%	0.0047%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0040%	-	-	-	-	0.0013%	0.0042%	0.0140%	0.0284%	0.0266%	0.0043%
	Males	0.0063%	-	-	-	-	0.0020%	0.0088%	0.0235%	0.0476%	0.0517%	0.0076%
	Females	0.0018%	-	-	-	-	-	-	0.0060%	0.0141%	0.0108%	0.0019%
Hispanic <sup>d</sup>	Both Sexes	-	-	-	-	-	-	-	-	-	-	-
	Males	-	-	-	-	-	-	-	-	-	-	-
	Females	-	-	-	-	-	-	-	-	-	-	-

<sup>a</sup> US 2008 cancer prevalence counts are based on 2008 cancer prevalence proportions from the SEER registries and 1/1/2008 US population estimates based on the average of 2007 and 2008 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 33 years for 1975-2008 SEER 9 data; 18 years for 1990-2008 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 33 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <33 and >=33. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 9.21  
Hodgkin LymphomaEstimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2008  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <33 <sup>e</sup>	>=33 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	35,728	29,609	25,629	22,995	17,817	12,409	106,629	152,859	13,917	166,776
	Males	18,621	15,264	13,210	11,656	9,396	6,452	55,639	79,172	7,046	86,218
	Females	17,107	14,345	12,419	11,339	8,421	5,957	50,990	73,687	6,871	80,558
White <sup>b</sup>	Both Sexes	30,306	25,807	22,847	21,234	16,492	11,715	92,811	136,078	15,515	151,593
	Males	15,902	13,273	11,844	10,858	8,542	6,006	48,651	70,429	8,060	78,489
	Females	14,404	12,534	11,003	10,376	7,950	5,709	44,160	65,649	7,455	73,104
Black <sup>b</sup>	Both Sexes	4,029	2,948	2,286	1,390	1,057	575	10,709	13,014	1,050	14,064
	Males	1,944	1,526	1,133	612	714	365	5,322	6,732	720	7,452
	Females	2,085	1,422	1,153	778	343	210	5,387	6,282	330	6,612
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	847	558	290	+	+	+	1,847	+	+	+
	Males	454	297	131	+	+	+	956	+	+	+
	Females	393	261	159	+	+	+	891	+	+	+
Hispanic <sup>d</sup>	Both Sexes	3,542	2,620	1,775	+	+	+	8,899	+	+	+
	Males	1,971	1,505	942	+	+	+	4,958	+	+	+
	Females	1,571	1,115	833	+	+	+	3,941	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2008, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0285%	0.0007%	0.0098%	0.0372%	0.0520%	0.0437%	0.0303%	0.0265%	0.0250%	0.0185%	0.0284%
	Males	0.0303%	0.0009%	0.0115%	0.0344%	0.0505%	0.0480%	0.0361%	0.0325%	0.0289%	0.0234%	0.0302%
	Females	0.0268%	0.0004%	0.0081%	0.0402%	0.0537%	0.0395%	0.0248%	0.0211%	0.0219%	0.0158%	0.0268%
White <sup>c</sup>	Both Sexes	0.0319%	0.0007%	0.0109%	0.0415%	0.0597%	0.0489%	0.0335%	0.0291%	0.0281%	0.0204%	0.0319%
	Males	0.0337%	0.0009%	0.0125%	0.0377%	0.0579%	0.0529%	0.0396%	0.0352%	0.0320%	0.0252%	0.0335%
	Females	0.0301%	0.0005%	0.0092%	0.0457%	0.0615%	0.0448%	0.0274%	0.0235%	0.0248%	0.0177%	0.0305%
Black <sup>c</sup>	Both Sexes	0.0252%	-	0.0075%	0.0295%	0.0479%	0.0414%	0.0300%	0.0278%	0.0214%	0.0146%	0.0258%
	Males	0.0265%	-	0.0084%	0.0278%	0.0430%	0.0483%	0.0375%	0.0343%	0.0280%	0.0247%	0.0281%
	Females	0.0241%	-	0.0066%	0.0312%	0.0522%	0.0353%	0.0238%	0.0227%	0.0169%	0.0099%	0.0240%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0113%	-	0.0053%	0.0201%	0.0178%	0.0142%	0.0103%	0.0079%	0.0085%	0.0067%	0.0108%
	Males	0.0121%	-	0.0078%	0.0210%	0.0164%	0.0153%	0.0103%	0.0121%	0.0094%	0.0119%	0.0119%
	Females	0.0105%	-	0.0027%	0.0192%	0.0190%	0.0132%	0.0104%	0.0042%	0.0078%	-	0.0099%
Hispanic <sup>d</sup>	Both Sexes	-	-	-	-	-	-	-	-	-	-	-
	Males	-	-	-	-	-	-	-	-	-	-	-
	Females	-	-	-	-	-	-	-	-	-	-	-

<sup>a</sup> US 2008 cancer prevalence counts are based on 2008 cancer prevalence proportions from the SEER registries and 1/1/2008 US population estimates based on the average of 2007 and 2008 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 33 years for 1975-2008 SEER 9 data; 18 years for 1990-2008 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 33 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <33 and >=33. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 10.15  
Kaposi SarcomaEstimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2008  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <33 <sup>e</sup>	>=33 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	6,212	5,026	7,147	1,933	617	204	20,361	22,104	+	+
	Males	5,793	4,683	6,912	1,791	584	134	19,173	20,791	+	+
	Females	419	343	235	142	33	70	1,188	1,313	+	+
White <sup>b</sup>	Both Sexes	4,509	3,628	5,908	1,600	509	171	15,633	17,087	+	+
	Males	4,186	3,381	5,741	1,490	487	101	14,731	16,071	+	+
	Females	323	247	167	110	22	70	902	1,016	+	+
Black <sup>b</sup>	Both Sexes	1,440	1,077	843	154	64	10	3,669	3,805	+	+
	Males	1,362	1,014	786	144	64	10	3,456	3,592	+	+
	Females	78	63	57	10	0	0	213	213	+	+
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	93	83	83	+	+	+	267	+	+	+
	Males	88	83	83	+	+	+	262	+	+	+
	Females	5	0	0	+	+	+	5	+	+	+
Hispanic <sup>d</sup>	Both Sexes	1,228	970	1,193	+	+	+	3,659	+	+	+
	Males	1,168	901	1,168	+	+	+	3,481	+	+	+
	Females	60	69	25	+	+	+	178	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2008, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0072%	-	-	0.0009%	0.0068%	0.0196%	0.0142%	0.0081%	0.0071%	0.0130%	0.0072%
	Males	0.0140%	-	-	0.0018%	0.0131%	0.0385%	0.0286%	0.0159%	0.0139%	0.0247%	0.0141%
	Females	0.0007%	-	-	-	0.0002%	0.0007%	0.0004%	0.0010%	0.0017%	0.0065%	0.0006%
White <sup>c</sup>	Both Sexes	0.0073%	-	-	0.0006%	0.0063%	0.0195%	0.0146%	0.0083%	0.0078%	0.0132%	0.0072%
	Males	0.0140%	-	-	0.0012%	0.0119%	0.0377%	0.0292%	0.0161%	0.0149%	0.0251%	0.0139%
	Females	0.0007%	-	-	-	0.0002%	0.0006%	-	0.0011%	0.0021%	0.0067%	0.0006%
Black <sup>c</sup>	Both Sexes	0.0098%	-	-	0.0029%	0.0140%	0.0293%	0.0178%	0.0094%	0.0043%	0.0101%	0.0103%
	Males	0.0196%	-	-	0.0055%	0.0296%	0.0615%	0.0371%	0.0200%	0.0091%	-	0.0211%
	Females	0.0008%	-	-	-	-	-	0.0017%	-	-	0.0115%	0.0010%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0018%	-	-	-	0.0015%	0.0041%	0.0041%	0.0016%	0.0029%	0.0059%	0.0018%
	Males	0.0037%	-	-	-	0.0031%	0.0082%	0.0089%	0.0035%	0.0068%	0.0136%	0.0039%
	Females	-	-	-	-	-	-	-	-	-	-	-
Hispanic <sup>d</sup>	Both Sexes	-	-	-	-	-	-	-	-	-	-	-
	Males	-	-	-	-	-	-	-	-	-	-	-
	Females	-	-	-	-	-	-	-	-	-	-	-

<sup>a</sup> US 2008 cancer prevalence counts are based on 2008 cancer prevalence proportions from the SEER registries and 1/1/2008 US population estimates based on the average of 2007 and 2008 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 33 years for 1975-2008 SEER 9 data; 18 years for 1990-2008 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 33 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <33 and >=33. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 11.21  
Cancer of the Kidney And Renal Pelvis (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2008  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <33 <sup>e</sup>	>=33 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	118,656	64,398	40,460	24,125	13,726	7,577	252,095	285,286	10,788	296,074
	Males	71,212	37,694	23,736	13,729	8,074	3,970	150,012	168,405	5,945	174,350
	Females	47,444	26,704	16,724	10,396	5,652	3,607	102,083	116,881	4,843	121,724
White <sup>b</sup>	Both Sexes	101,614	55,395	34,968	21,570	12,294	6,652	216,342	246,063	12,354	258,417
	Males	61,102	32,877	20,939	12,429	7,375	3,585	129,807	146,640	5,518	152,158
	Females	40,512	22,518	14,029	9,141	4,919	3,067	86,535	99,423	6,836	106,259
Black <sup>b</sup>	Both Sexes	11,788	6,679	4,183	2,010	1,160	766	25,594	28,438	1,862	30,300
	Males	6,962	3,407	2,063	1,013	523	313	14,111	15,327	638	15,965
	Females	4,826	3,272	2,120	997	637	453	11,483	13,111	1,224	14,335
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	2,946	1,421	698	+	+	+	5,295	+	+	+
	Males	1,841	859	440	+	+	+	3,267	+	+	+
	Females	1,105	562	258	+	+	+	2,028	+	+	+
Hispanic <sup>d</sup>	Both Sexes	10,670	5,375	2,726	+	+	+	19,898	+	+	+
	Males	6,216	3,119	1,483	+	+	+	11,396	+	+	+
	Females	4,454	2,256	1,243	+	+	+	8,502	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2008, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0702%	0.0064%	0.0089%	0.0028%	0.0114%	0.0455%	0.1183%	0.2426%	0.3260%	0.2829%	0.0709%
	Males	0.0854%	0.0062%	0.0075%	0.0027%	0.0120%	0.0557%	0.1559%	0.3309%	0.4391%	0.4031%	0.0936%
	Females	0.0552%	0.0065%	0.0104%	0.0029%	0.0108%	0.0353%	0.0824%	0.1626%	0.2369%	0.2164%	0.0521%
White <sup>c</sup>	Both Sexes	0.0765%	0.0067%	0.0095%	0.0025%	0.0120%	0.0481%	0.1231%	0.2560%	0.3415%	0.2957%	0.0743%
	Males	0.0930%	0.0065%	0.0084%	0.0023%	0.0125%	0.0573%	0.1611%	0.3457%	0.4619%	0.4263%	0.0978%
	Females	0.0600%	0.0069%	0.0106%	0.0028%	0.0115%	0.0386%	0.0854%	0.1723%	0.2443%	0.2237%	0.0545%
Black <sup>c</sup>	Both Sexes	0.0591%	0.0072%	0.0104%	0.0044%	0.0128%	0.0455%	0.1357%	0.2509%	0.3261%	0.2667%	0.0737%
	Males	0.0698%	0.0073%	0.0066%	0.0047%	0.0128%	0.0601%	0.1836%	0.3580%	0.4078%	0.3356%	0.0957%
	Females	0.0495%	0.0072%	0.0144%	0.0041%	0.0128%	0.0328%	0.0959%	0.1684%	0.2702%	0.2347%	0.0574%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0388%	0.0032%	0.0027%	0.0016%	0.0052%	0.0255%	0.0655%	0.1305%	0.2037%	0.1776%	0.0405%
	Males	0.0494%	0.0029%	0.0021%	0.0022%	0.0051%	0.0360%	0.0901%	0.1857%	0.2799%	0.2598%	0.0559%
	Females	0.0289%	0.0036%	0.0033%	-	0.0053%	0.0159%	0.0442%	0.0835%	0.1467%	0.1260%	0.0282%
Hispanic <sup>d</sup>	Both Sexes	-	-	-	-	-	-	-	-	-	-	-
	Males	-	-	-	-	-	-	-	-	-	-	-
	Females	-	-	-	-	-	-	-	-	-	-	-

<sup>a</sup> US 2008 cancer prevalence counts are based on 2008 cancer prevalence proportions from the SEER registries and 1/1/2008 US population estimates based on the average of 2007 and 2008 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 33 years for 1975-2008 SEER 9 data; 18 years for 1990-2008 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 33 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <33 and >=33. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 12.21  
Cancer of the Larynx (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2008  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <33 <sup>e</sup>	>=33 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	30,315	21,190	14,645	10,283	6,490	3,956	71,378	86,346	2,595	88,941
	Males	24,076	17,064	11,688	8,151	5,238	3,194	57,017	69,149	2,124	71,273
	Females	6,239	4,126	2,957	2,132	1,252	762	14,361	17,197	471	17,668
White <sup>b</sup>	Both Sexes	25,306	18,174	12,638	8,793	5,790	3,567	60,604	73,726	2,067	75,793
	Males	20,046	14,681	10,147	7,050	4,744	2,879	48,511	59,298	1,687	60,985
	Females	5,260	3,493	2,491	1,743	1,046	688	12,093	14,428	380	14,808
Black <sup>b</sup>	Both Sexes	4,229	2,449	1,679	1,234	598	280	8,965	10,470	180	10,650
	Males	3,351	1,885	1,292	928	419	223	6,975	8,093	152	8,245
	Females	878	564	387	306	179	57	1,990	2,377	28	2,405
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	468	328	217	+	+	+	1,113	+	+	+
	Males	412	280	188	+	+	+	969	+	+	+
	Females	56	48	29	+	+	+	144	+	+	+
Hispanic <sup>d</sup>	Both Sexes	1,605	1,049	613	+	+	+	3,593	+	+	+
	Males	1,371	870	520	+	+	+	3,023	+	+	+
	Females	234	179	93	+	+	+	570	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2008, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0185%	-	-	0.0002%	0.0007%	0.0060%	0.0273%	0.0741%	0.1124%	0.0829%	0.0190%
	Males	0.0303%	-	-	0.0002%	0.0008%	0.0086%	0.0442%	0.1301%	0.2060%	0.1941%	0.0346%
	Females	0.0070%	-	-	0.0002%	0.0006%	0.0034%	0.0112%	0.0234%	0.0385%	0.0213%	0.0066%
White <sup>c</sup>	Both Sexes	0.0200%	-	-	0.0003%	0.0009%	0.0065%	0.0282%	0.0761%	0.1163%	0.0849%	0.0196%
	Males	0.0324%	-	-	0.0002%	0.0009%	0.0091%	0.0456%	0.1311%	0.2104%	0.1978%	0.0353%
	Females	0.0076%	-	-	0.0003%	0.0009%	0.0037%	0.0111%	0.0250%	0.0404%	0.0227%	0.0070%
Black <sup>c</sup>	Both Sexes	0.0201%	-	-	-	0.0008%	0.0069%	0.0436%	0.1163%	0.1486%	0.0948%	0.0266%
	Males	0.0332%	-	-	-	-	0.0095%	0.0697%	0.2228%	0.2868%	0.2490%	0.0512%
	Females	0.0083%	-	-	-	-	0.0047%	0.0220%	0.0343%	0.0539%	0.0230%	0.0097%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0081%	-	-	-	-	0.0025%	0.0080%	0.0277%	0.0593%	0.0596%	0.0088%
	Males	0.0146%	-	-	-	-	0.0047%	0.0134%	0.0545%	0.1187%	0.1374%	0.0178%
	Females	0.0021%	-	-	-	-	-	0.0033%	0.0049%	0.0148%	0.0108%	0.0020%
Hispanic <sup>d</sup>	Both Sexes	-	-	-	-	-	-	-	-	-	-	-
	Males	-	-	-	-	-	-	-	-	-	-	-
	Females	-	-	-	-	-	-	-	-	-	-	-

<sup>a</sup> US 2008 cancer prevalence counts are based on 2008 cancer prevalence proportions from the SEER registries and 1/1/2008 US population estimates based on the average of 2007 and 2008 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 33 years for 1975-2008 SEER 9 data; 18 years for 1990-2008 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 33 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <33 and >=33. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 13.29  
LeukemiaEstimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2008  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <33 <sup>e</sup>	>=33 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	99,438	56,503	36,126	21,981	13,694	7,934	213,896	247,589	5,761	253,350
	Males	57,215	32,396	20,454	12,125	7,581	3,818	121,857	140,084	2,618	142,702
	Females	42,223	24,107	15,672	9,856	6,113	4,116	92,039	107,505	3,143	110,648
White <sup>b</sup>	Both Sexes	87,330	50,473	32,506	19,859	12,577	7,309	190,066	220,792	4,883	225,675
	Males	50,635	28,985	18,472	11,159	6,990	3,553	108,731	125,532	1,890	127,422
	Females	36,695	21,488	14,034	8,700	5,587	3,756	81,335	95,260	2,993	98,253
Black <sup>b</sup>	Both Sexes	7,025	3,371	1,916	1,158	551	350	13,360	14,947	662	15,609
	Males	3,780	1,694	980	508	313	119	6,993	7,734	305	8,039
	Females	3,245	1,677	936	650	238	231	6,367	7,213	357	7,570
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	2,487	1,457	957	+	+	+	5,338	+	+	+
	Males	1,347	846	519	+	+	+	2,951	+	+	+
	Females	1,140	611	438	+	+	+	2,387	+	+	+
Hispanic <sup>d</sup>	Both Sexes	9,123	5,565	3,707	+	+	+	19,980	+	+	+
	Males	5,427	2,943	2,145	+	+	+	11,457	+	+	+
	Females	3,696	2,622	1,562	+	+	+	8,523	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2008, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0611%	0.0292%	0.0529%	0.0221%	0.0214%	0.0339%	0.0664%	0.1394%	0.2226%	0.2483%	0.0623%
	Males	0.0706%	0.0314%	0.0578%	0.0235%	0.0244%	0.0401%	0.0820%	0.1815%	0.2936%	0.3416%	0.0771%
	Females	0.0517%	0.0269%	0.0477%	0.0207%	0.0182%	0.0276%	0.0515%	0.1012%	0.1666%	0.1967%	0.0501%
White <sup>c</sup>	Both Sexes	0.0688%	0.0322%	0.0593%	0.0245%	0.0235%	0.0354%	0.0718%	0.1532%	0.2467%	0.2725%	0.0684%
	Males	0.0792%	0.0347%	0.0644%	0.0259%	0.0266%	0.0422%	0.0878%	0.1986%	0.3247%	0.3744%	0.0843%
	Females	0.0584%	0.0296%	0.0540%	0.0229%	0.0202%	0.0283%	0.0560%	0.1108%	0.1838%	0.2163%	0.0549%
Black <sup>c</sup>	Both Sexes	0.0321%	0.0137%	0.0236%	0.0153%	0.0129%	0.0250%	0.0466%	0.0835%	0.1507%	0.1319%	0.0378%
	Males	0.0360%	0.0165%	0.0246%	0.0161%	0.0139%	0.0294%	0.0605%	0.1075%	0.1725%	0.1934%	0.0459%
	Females	0.0285%	0.0109%	0.0225%	0.0146%	0.0120%	0.0210%	0.0350%	0.0651%	0.1358%	0.1033%	0.0320%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0341%	0.0270%	0.0470%	0.0154%	0.0175%	0.0282%	0.0366%	0.0606%	0.0760%	0.0833%	0.0357%
	Males	0.0390%	0.0277%	0.0552%	0.0161%	0.0213%	0.0312%	0.0425%	0.0687%	0.1003%	0.1198%	0.0421%
	Females	0.0295%	0.0263%	0.0382%	0.0146%	0.0138%	0.0254%	0.0315%	0.0538%	0.0579%	0.0604%	0.0303%
Hispanic <sup>d</sup>	Both Sexes	-	-	-	-	-	-	-	-	-	-	-
	Males	-	-	-	-	-	-	-	-	-	-	-
	Females	-	-	-	-	-	-	-	-	-	-	-

<sup>a</sup> US 2008 cancer prevalence counts are based on 2008 cancer prevalence proportions from the SEER registries and 1/1/2008 US population estimates based on the average of 2007 and 2008 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 33 years for 1975-2008 SEER 9 data; 18 years for 1990-2008 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 33 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <33 and >=33. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.



Table 13.30  
Acute Lymphocytic Leukemia

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2008  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <33 <sup>e</sup>	>=33 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	14,970	11,377	9,740	8,456	6,603	4,787	42,240	58,854	3,339	62,193
	Males	8,566	6,204	5,463	4,628	3,618	2,365	23,656	32,504	1,802	34,306
	Females	6,404	5,173	4,277	3,828	2,985	2,422	18,584	26,350	1,537	27,887
White <sup>b</sup>	Both Sexes	12,749	9,996	8,426	7,435	5,982	4,419	36,484	51,483	+	+
	Males	7,338	5,453	4,794	4,144	3,298	2,220	20,549	28,617	+	+
	Females	5,411	4,543	3,632	3,291	2,684	2,199	15,935	22,866	+	+
Black <sup>b</sup>	Both Sexes	1,061	721	637	536	291	172	2,878	3,671	+	+
	Males	624	356	296	237	169	55	1,545	1,930	+	+
	Females	437	365	341	299	122	117	1,333	1,741	+	+
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	768	529	407	+	+	+	1,940	+	+	+
	Males	406	322	232	+	+	+	1,073	+	+	+
	Females	362	207	175	+	+	+	867	+	+	+
Hispanic <sup>d</sup>	Both Sexes	4,281	2,834	2,234	+	+	+	10,319	+	+	+
	Males	2,571	1,475	1,321	+	+	+	5,927	+	+	+
	Females	1,710	1,359	913	+	+	+	4,392	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2008, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0136%	0.0248%	0.0453%	0.0131%	0.0050%	0.0036%	0.0029%	0.0032%	0.0021%	0.0017%	0.0140%
	Males	0.0155%	0.0264%	0.0500%	0.0147%	0.0062%	0.0042%	0.0035%	0.0036%	0.0025%	0.0024%	0.0155%
	Females	0.0117%	0.0232%	0.0403%	0.0114%	0.0037%	0.0029%	0.0024%	0.0029%	0.0018%	0.0013%	0.0124%
White <sup>c</sup>	Both Sexes	0.0151%	0.0275%	0.0516%	0.0152%	0.0057%	0.0039%	0.0032%	0.0035%	0.0018%	0.0016%	0.0158%
	Males	0.0171%	0.0294%	0.0565%	0.0170%	0.0070%	0.0046%	0.0040%	0.0040%	0.0022%	0.0022%	0.0175%
	Females	0.0131%	0.0256%	0.0464%	0.0133%	0.0043%	0.0031%	0.0025%	0.0031%	0.0015%	0.0012%	0.0140%
Black <sup>c</sup>	Both Sexes	0.0065%	0.0111%	0.0180%	0.0074%	0.0017%	0.0007%	0.0013%	-	-	-	0.0060%
	Males	0.0075%	0.0134%	0.0192%	0.0077%	0.0015%	-	-	-	-	-	0.0066%
	Females	0.0057%	0.0086%	0.0167%	0.0070%	0.0019%	-	0.0017%	-	-	-	0.0054%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0109%	0.0227%	0.0380%	0.0072%	0.0041%	0.0045%	0.0021%	0.0034%	0.0038%	-	0.0120%
	Males	0.0127%	0.0222%	0.0452%	0.0087%	0.0052%	0.0050%	0.0024%	0.0029%	-	-	0.0134%
	Females	0.0091%	0.0232%	0.0304%	0.0057%	0.0030%	0.0041%	0.0018%	0.0039%	0.0036%	-	0.0105%
Hispanic <sup>d</sup>	Both Sexes	-	-	-	-	-	-	-	-	-	-	-
	Males	-	-	-	-	-	-	-	-	-	-	-
	Females	-	-	-	-	-	-	-	-	-	-	-

<sup>a</sup> US 2008 cancer prevalence counts are based on 2008 cancer prevalence proportions from the SEER registries and 1/1/2008 US population estimates based on the average of 2007 and 2008 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 33 years for 1975-2008 SEER 9 data; 18 years for 1990-2008 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 33 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <33 and >=33. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 14.21  
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2008  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <33 <sup>e</sup>	>=33 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	20,771	3,797	1,318	790	558	199	29,617	30,771	404	31,175
	Males	14,807	2,413	767	351	246	65	21,032	21,496	71	21,567
	Females	5,964	1,384	551	439	312	134	8,585	9,275	333	9,608
White <sup>b</sup>	Both Sexes	14,962	2,779	984	624	444	131	21,383	22,289	343	22,632
	Males	10,897	1,728	558	252	189	35	15,285	15,619	169 <sup>i</sup>	15,788
	Females	4,065	1,051	426	372	255	96	6,098	6,670	174 <sup>i</sup>	6,844
Black <sup>b</sup>	Both Sexes	2,834	286	90	93	62	45	3,739	3,887	+	+
	Males	1,860	172	43	47	27	22	2,557	2,627	+	+
	Females	974	114	47	46	35	23	1,182	1,260	+	+
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	2,900	754	200	+	+	+	3,910	+	+	+
	Males	2,081	506	146	+	+	+	2,772	+	+	+
	Females	819	248	54	+	+	+	1,138	+	+	+
Hispanic <sup>d</sup>	Both Sexes	3,383	613	228	+	+	+	4,305	+	+	+
	Males	2,494	376	160	+	+	+	3,090	+	+	+
	Females	889	237	68	+	+	+	1,215	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2008, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0117%	0.0024%	0.0016%	0.0007%	0.0014%	0.0062%	0.0273%	0.0408%	0.0480%	0.0273%	0.0116%
	Males	0.0167%	0.0024%	0.0018%	0.0007%	0.0018%	0.0093%	0.0453%	0.0621%	0.0688%	0.0414%	0.0173%
	Females	0.0068%	0.0023%	0.0013%	0.0007%	0.0009%	0.0032%	0.0101%	0.0216%	0.0317%	0.0195%	0.0065%
White <sup>c</sup>	Both Sexes	0.0091%	0.0025%	0.0015%	0.0007%	0.0011%	0.0047%	0.0221%	0.0292%	0.0317%	0.0196%	0.0086%
	Males	0.0130%	0.0026%	0.0018%	0.0008%	0.0013%	0.0067%	0.0365%	0.0447%	0.0455%	0.0296%	0.0129%
	Females	0.0052%	0.0025%	0.0013%	0.0006%	0.0009%	0.0025%	0.0077%	0.0148%	0.0206%	0.0141%	0.0048%
Black <sup>c</sup>	Both Sexes	0.0093%	-	0.0009%	0.0014%	0.0012%	0.0039%	0.0334%	0.0431%	0.0400%	0.0156%	0.0108%
	Males	0.0135%	-	0.0013%	-	0.0016%	0.0049%	0.0549%	0.0734%	0.0583%	0.0283%	0.0170%
	Females	0.0054%	-	-	0.0019%	-	0.0031%	0.0155%	0.0197%	0.0274%	0.0096%	0.0061%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0292%	0.0031%	0.0021%	-	0.0026%	0.0179%	0.0532%	0.1136%	0.1562%	0.0888%	0.0304%
	Males	0.0423%	0.0029%	0.0022%	-	0.0046%	0.0297%	0.0927%	0.1731%	0.2297%	0.1280%	0.0468%
	Females	0.0169%	0.0034%	0.0020%	-	-	0.0070%	0.0188%	0.0630%	0.1013%	0.0641%	0.0168%
Hispanic <sup>d</sup>	Both Sexes	-	-	-	-	-	-	-	-	-	-	-
	Males	-	-	-	-	-	-	-	-	-	-	-
	Females	-	-	-	-	-	-	-	-	-	-	-

<sup>a</sup> US 2008 cancer prevalence counts are based on 2008 cancer prevalence proportions from the SEER registries and 1/1/2008 US population estimates based on the average of 2007 and 2008 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 33 years for 1975-2008 SEER 9 data; 18 years for 1990-2008 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 33 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <33 and >=33. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 15.27  
Cancer of the Lung And Bronchus (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2008  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <33 <sup>e</sup>	>=33 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	214,402	69,655	38,971	20,535	10,970	5,721	338,696	364,377	9,112	373,489
	Males	101,209	31,015	17,928	9,571	4,964	2,741	156,162	168,236	5,192	173,428
	Females	113,193	38,640	21,043	10,964	6,006	2,980	182,534	196,141	3,920	200,061
White <sup>b</sup>	Both Sexes	184,129	61,258	34,230	18,320	9,683	5,167	293,306	316,025	4,120	320,145
	Males	85,874	26,864	15,432	8,397	4,353	2,442	133,331	143,833	1,987	145,820
	Females	98,255	34,394	18,798	9,923	5,330	2,725	159,975	172,192	2,133	174,325
Black <sup>b</sup>	Both Sexes	22,189	6,066	3,553	1,696	1,003	421	33,267	35,569	378	35,947
	Males	11,030	2,881	1,816	920	454	218	16,363	17,595	206	17,801
	Females	11,159	3,185	1,737	776	549	203	16,904	17,974	172	18,146
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	6,278	1,603	864	+	+	+	8,991	+	+	+
	Males	3,349	789	492	+	+	+	4,760	+	+	+
	Females	2,929	814	372	+	+	+	4,231	+	+	+
Hispanic <sup>d</sup>	Both Sexes	7,371	2,130	1,008	+	+	+	10,947	+	+	+
	Males	3,796	1,063	484	+	+	+	5,551	+	+	+
	Females	3,575	1,067	524	+	+	+	5,396	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2008, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0925%	0.0001%	0.0002%	0.0016%	0.0052%	0.0256%	0.1065%	0.3424%	0.6190%	0.5245%	0.0962%
	Males	0.0871%	0.0002%	0.0002%	0.0015%	0.0048%	0.0226%	0.1056%	0.3512%	0.6537%	0.6307%	0.1017%
	Females	0.0978%	-	-	0.0016%	0.0056%	0.0285%	0.1073%	0.3345%	0.5917%	0.4657%	0.0927%
White <sup>c</sup>	Both Sexes	0.0992%	0.0002%	0.0002%	0.0017%	0.0053%	0.0251%	0.1035%	0.3510%	0.6419%	0.5372%	0.0983%
	Males	0.0905%	0.0002%	-	0.0015%	0.0048%	0.0219%	0.0986%	0.3474%	0.6577%	0.6270%	0.1007%
	Females	0.1080%	-	-	0.0019%	0.0060%	0.0284%	0.1084%	0.3544%	0.6291%	0.4876%	0.0974%
Black <sup>c</sup>	Both Sexes	0.0789%	-	-	0.0012%	0.0053%	0.0352%	0.1575%	0.4171%	0.5994%	0.4677%	0.1054%
	Males	0.0816%	-	-	-	0.0052%	0.0317%	0.1800%	0.4842%	0.7104%	0.6853%	0.1255%
	Females	0.0765%	-	-	0.0014%	0.0053%	0.0382%	0.1388%	0.3654%	0.5232%	0.3664%	0.0924%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0698%	-	-	0.0007%	0.0044%	0.0208%	0.0826%	0.2388%	0.4949%	0.4610%	0.0753%
	Males	0.0760%	-	-	-	0.0042%	0.0207%	0.0920%	0.2910%	0.5920%	0.6293%	0.0915%
	Females	0.0641%	-	-	-	0.0045%	0.0208%	0.0744%	0.1944%	0.4223%	0.3552%	0.0632%
Hispanic <sup>d</sup>	Both Sexes	-	-	-	-	-	-	-	-	-	-	-
	Males	-	-	-	-	-	-	-	-	-	-	-
	Females	-	-	-	-	-	-	-	-	-	-	-

<sup>a</sup> US 2008 cancer prevalence counts are based on 2008 cancer prevalence proportions from the SEER registries and 1/1/2008 US population estimates based on the average of 2007 and 2008 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 33 years for 1975-2008 SEER 9 data; 18 years for 1990-2008 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 33 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <33 and >=33. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 16.21  
Melanoma of the Skin (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2008  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <33 <sup>e</sup>	>=33 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	241,058	176,666	124,103	90,687	64,051	42,149	619,976	782,730	40,040	822,770
	Males	127,277	90,838	62,241	42,150	27,810	17,035	318,521	387,474	13,232	400,706
	Females	113,781	85,828	61,862	48,537	36,241	25,114	301,455	395,256	26,808	422,064
White <sup>b</sup>	Both Sexes	232,442	170,770	118,922	87,204	61,002	40,794	597,809	753,604	37,829	791,433
	Males	123,037	87,875	59,671	40,934	26,485	16,399	307,819	374,049	12,628	386,677
	Females	109,405	82,895	59,251	46,270	34,517	24,395	289,990	379,555	25,201	404,756
Black <sup>b</sup>	Both Sexes	830	677	390	350	138	57	2,165	2,657	108	2,765
	Males	306	321	212	122	32	10	942	1,054	40	1,094
	Females	524	356	178	228	106	47	1,223	1,603	68	1,671
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	630	442	264	+	+	+	1,425	+	+	+
	Males	304	171	132	+	+	+	627	+	+	+
	Females	326	271	132	+	+	+	798	+	+	+
Hispanic <sup>d</sup>	Both Sexes	4,441	3,288	2,240	+	+	+	10,994	+	+	+
	Males	1,783	1,147	814	+	+	+	4,091	+	+	+
	Females	2,658	2,141	1,426	+	+	+	6,903	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2008, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.1567%	0.0004%	0.0037%	0.0292%	0.0901%	0.1802%	0.2836%	0.4293%	0.5107%	0.5303%	0.1562%
	Males	0.1653%	0.0004%	0.0031%	0.0184%	0.0630%	0.1537%	0.3001%	0.5399%	0.7293%	0.8385%	0.1798%
	Females	0.1482%	0.0004%	0.0043%	0.0407%	0.1181%	0.2069%	0.2678%	0.3291%	0.3383%	0.3597%	0.1412%
White <sup>c</sup>	Both Sexes	0.1989%	0.0004%	0.0046%	0.0374%	0.1161%	0.2266%	0.3525%	0.5215%	0.6238%	0.6201%	0.1924%
	Males	0.2084%	0.0005%	0.0039%	0.0231%	0.0791%	0.1894%	0.3676%	0.6481%	0.8803%	0.9874%	0.2176%
	Females	0.1895%	0.0004%	0.0053%	0.0530%	0.1559%	0.2654%	0.3376%	0.4036%	0.4166%	0.4177%	0.1774%
Black <sup>c</sup>	Both Sexes	0.0050%	-	-	0.0015%	0.0037%	0.0050%	0.0067%	0.0198%	0.0278%	0.0270%	0.0063%
	Males	0.0045%	-	-	-	0.0021%	0.0046%	0.0065%	0.0207%	0.0400%	0.0353%	0.0069%
	Females	0.0054%	-	-	0.0024%	0.0051%	0.0053%	0.0068%	0.0190%	0.0194%	0.0231%	0.0061%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0098%	-	0.0011%	0.0021%	0.0067%	0.0120%	0.0144%	0.0278%	0.0244%	0.0440%	0.0099%
	Males	0.0093%	-	-	0.0025%	0.0043%	0.0091%	0.0133%	0.0305%	0.0335%	0.0566%	0.0102%
	Females	0.0103%	-	-	0.0016%	0.0089%	0.0147%	0.0153%	0.0255%	0.0176%	0.0361%	0.0098%
Hispanic <sup>d</sup>	Both Sexes	-	-	-	-	-	-	-	-	-	-	-
	Males	-	-	-	-	-	-	-	-	-	-	-
	Females	-	-	-	-	-	-	-	-	-	-	-

<sup>a</sup> US 2008 cancer prevalence counts are based on 2008 cancer prevalence proportions from the SEER registries and 1/1/2008 US population estimates based on the average of 2007 and 2008 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 33 years for 1975-2008 SEER 9 data; 18 years for 1990-2008 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 33 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <33 and >=33. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 17.13  
MesotheliomaEstimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2008  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <33 <sup>e</sup>	>=33 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	2,955	599	318	236	150	159	4,090	4,528	+	+
	Males	2,081	291	105	58	80	60	2,471	2,635	+	+
	Females	874	308	213	178	70	99	1,619	1,893	+	+
White <sup>b</sup>	Both Sexes	2,759	486	282	232	128	135	3,758	4,148	+	+
	Males	1,947	235	92	54	58	47	2,275	2,401	+	+
	Females	812	251	190	178	70	88	1,483	1,747	+	+
Black <sup>b</sup>	Both Sexes	96	69	24	0	23	23	209	254	+	+
	Males	60	34	13	0	23	13	127	162	+	+
	Females	36	35	11	0	0	10	82	92	+	+
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	51	4	7	+	+	+	61	+	+	+
	Males	36	1	7	+	+	+	43	+	+	+
	Females	15	3	0	+	+	+	18	+	+	+
Hispanic <sup>d</sup>	Both Sexes	233	63	10	+	+	+	312	+	+	+
	Males	108	47	0	+	+	+	156	+	+	+
	Females	125	16	10	+	+	+	156	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2008, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	Age-Adjusted <sup>f</sup> All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0012%	-	-	-	0.0002%	0.0006%	0.0014%	0.0038%	0.0066%	0.0075%	0.0012%
	Males	0.0015%	-	-	-	-	0.0005%	0.0015%	0.0055%	0.0102%	0.0143%	0.0017%
	Females	0.0009%	-	-	-	0.0004%	0.0007%	0.0012%	0.0024%	0.0037%	0.0037%	0.0008%
White <sup>c</sup>	Both Sexes	0.0014%	-	-	-	0.0002%	0.0006%	0.0016%	0.0043%	0.0076%	0.0085%	0.0013%
	Males	0.0017%	-	-	-	-	0.0004%	0.0017%	0.0061%	0.0113%	0.0157%	0.0019%
	Females	0.0011%	-	-	-	0.0004%	0.0008%	0.0015%	0.0027%	0.0045%	0.0045%	0.0010%
Black <sup>c</sup>	Both Sexes	0.0005%	-	-	-	-	-	0.0010%	0.0024%	0.0031%	-	0.0006%
	Males	0.0007%	-	-	-	-	-	-	-	-	-	0.0010%
	Females	0.0003%	-	-	-	-	-	-	-	-	-	0.0004%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0005%	-	-	-	-	-	-	0.0016%	0.0021%	-	0.0005%
	Males	0.0007%	-	-	-	-	-	-	0.0029%	0.0049%	-	0.0009%
	Females	0.0002%	-	-	-	-	-	-	-	-	-	0.0002%
Hispanic <sup>d</sup>	Both Sexes	-	-	-	-	-	-	-	-	-	-	-
	Males	-	-	-	-	-	-	-	-	-	-	-
	Females	-	-	-	-	-	-	-	-	-	-	-

<sup>a</sup> US 2008 cancer prevalence counts are based on 2008 cancer prevalence proportions from the SEER registries and 1/1/2008 US population estimates based on the average of 2007 and 2008 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 33 years for 1975-2008 SEER 9 data; 18 years for 1990-2008 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 33 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <33 and >=33. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 18.21  
Myeloma (Invasive)Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2008  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <33 <sup>e</sup>	>=33 <sup>g</sup>	Complete <sup>h</sup>	
Race	Sex											
All Races <sup>b</sup>	Both Sexes	40,832	12,683	4,969	2,065	608	277	62,683	64,351	264	64,615	
	Males	21,848	7,254	2,753	1,193	391	122	34,428	35,347	98	35,445	
	Females	18,984	5,429	2,216	872	217	155	28,255	29,004	166	29,170	
White <sup>b</sup>	Both Sexes	32,008	9,945	3,875	1,523	434	219	49,035	50,193	102	50,295	
	Males	17,823	5,940	2,194	910	283	97	27,966	28,601	0 <sup>i</sup>	28,601	
	Females	14,185	4,005	1,681	613	151	122	21,069	21,592	102 <sup>i</sup>	21,694	
Black <sup>b</sup>	Both Sexes	7,328	2,171	903	461	135	49	11,343	11,776	+	+	
	Males	3,307	1,012	482	256	85	25	5,336	5,595	+	+	
	Females	4,021	1,159	421	205	50	24	6,007	6,181	+	+	
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	885	317	112	+	+	+	1,358	+	+	+	
	Males	454	152	61	+	+	+	691	+	+	+	
	Females	431	165	51	+	+	+	667	+	+	+	
Hispanic <sup>d</sup>	Both Sexes	3,009	1,041	334	+	+	+	4,473	+	+	+	
	Males	1,694	568	210	+	+	+	2,532	+	+	+	
	Females	1,315	473	124	+	+	+	1,941	+	+	+	

Estimated prevalence percent<sup>a</sup> on January 1, 2008, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	All Ages	Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages	
Race	Sex											
All Races <sup>c</sup>	Both Sexes	0.0186%	-	-	0.0002%	0.0021%	0.0092%	0.0299%	0.0706%	0.0955%	0.0891%	0.0188%
	Males	0.0206%	-	-	0.0003%	0.0028%	0.0105%	0.0352%	0.0857%	0.1169%	0.1131%	0.0229%
	Females	0.0166%	-	-	-	0.0013%	0.0078%	0.0250%	0.0569%	0.0787%	0.0758%	0.0155%
White <sup>c</sup>	Both Sexes	0.0183%	-	-	0.0002%	0.0019%	0.0085%	0.0278%	0.0662%	0.0902%	0.0844%	0.0177%
	Males	0.0209%	-	-	0.0003%	0.0026%	0.0104%	0.0332%	0.0819%	0.1165%	0.1106%	0.0222%
	Females	0.0157%	-	-	-	0.0011%	0.0065%	0.0226%	0.0516%	0.0690%	0.0699%	0.0138%
Black <sup>c</sup>	Both Sexes	0.0285%	-	-	-	0.0043%	0.0191%	0.0614%	0.1419%	0.1815%	0.1677%	0.0373%
	Males	0.0284%	-	-	-	0.0055%	0.0180%	0.0667%	0.1786%	0.1867%	0.1825%	0.0415%
	Females	0.0286%	-	-	-	0.0032%	0.0200%	0.0569%	0.1137%	0.1779%	0.1609%	0.0342%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0104%	-	-	-	0.0008%	0.0040%	0.0144%	0.0394%	0.0580%	0.0685%	0.0109%
	Males	0.0107%	-	-	-	0.0015%	0.0043%	0.0193%	0.0410%	0.0598%	0.0831%	0.0124%
	Females	0.0100%	-	-	-	-	0.0038%	0.0101%	0.0380%	0.0566%	0.0593%	0.0098%
Hispanic <sup>d</sup>	Both Sexes	-	-	-	-	-	-	-	-	-	-	-
	Males	-	-	-	-	-	-	-	-	-	-	-
	Females	-	-	-	-	-	-	-	-	-	-	-

<sup>a</sup> US 2008 cancer prevalence counts are based on 2008 cancer prevalence proportions from the SEER registries and 1/1/2008 US population estimates based on the average of 2007 and 2008 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 33 years for 1975-2008 SEER 9 data; 18 years for 1990-2008 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 33 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <33 and >=33. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 19.25  
Non-Hodgkin LymphomaEstimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2008  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <33 <sup>e</sup>	>=33 <sup>g</sup>	Complete <sup>h</sup>	
Race	Sex											
All Races <sup>b</sup>	Both Sexes	182,673	110,737	63,458	35,762	20,400	10,933	393,921	444,120	10,258	454,378	
	Males	95,333	57,078	32,992	18,498	10,558	5,416	204,407	230,061	5,372	235,433	
	Females	87,340	53,659	30,466	17,264	9,842	5,517	189,514	214,059	4,886	218,945	
White <sup>b</sup>	Both Sexes	161,166	98,707	56,413	32,455	18,598	10,042	349,134	394,945	10,581	405,526	
	Males	84,263	50,994	29,349	16,844	9,647	4,979	181,805	205,270	5,920	211,190	
	Females	76,903	47,713	27,064	15,611	8,951	5,063	167,329	189,675	4,661	194,336	
Black <sup>b</sup>	Both Sexes	13,987	7,425	4,336	2,144	1,146	604	28,195	31,023	791	31,814	
	Males	7,116	3,722	2,197	1,050	568	264	13,949	15,355	337	15,692	
	Females	6,871	3,703	2,139	1,094	578	340	14,246	15,668	454	16,122	
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	4,770	2,916	1,713	+	+	+	9,961	+	+	+	
	Males	2,495	1,511	845	+	+	+	5,149	+	+	+	
	Females	2,275	1,405	868	+	+	+	4,812	+	+	+	
Hispanic <sup>d</sup>	Both Sexes	12,495	7,295	3,969	+	+	+	25,097	+	+	+	
	Males	6,471	3,853	2,119	+	+	+	13,077	+	+	+	
	Females	6,024	3,442	1,850	+	+	+	12,020	+	+	+	

Estimated prevalence percent<sup>a</sup> on January 1, 2008, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	All Ages	Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages	
Race	Sex											
All Races <sup>c</sup>	Both Sexes	0.1116%	0.0024%	0.0114%	0.0200%	0.0383%	0.0844%	0.1708%	0.3267%	0.4881%	0.5342%	0.1129%
	Males	0.1177%	0.0031%	0.0148%	0.0248%	0.0437%	0.1000%	0.1936%	0.3687%	0.5432%	0.6117%	0.1290%
	Females	0.1057%	0.0017%	0.0078%	0.0150%	0.0328%	0.0687%	0.1489%	0.2887%	0.4446%	0.4913%	0.0991%
White <sup>c</sup>	Both Sexes	0.1239%	0.0025%	0.0117%	0.0208%	0.0398%	0.0887%	0.1816%	0.3510%	0.5320%	0.5688%	0.1207%
	Males	0.1301%	0.0033%	0.0153%	0.0259%	0.0459%	0.1049%	0.2052%	0.3936%	0.5886%	0.6506%	0.1374%
	Females	0.1178%	0.0018%	0.0080%	0.0152%	0.0332%	0.0719%	0.1583%	0.3114%	0.4863%	0.5236%	0.1060%
Black <sup>c</sup>	Both Sexes	0.0679%	0.0023%	0.0101%	0.0194%	0.0398%	0.0786%	0.1439%	0.2399%	0.2755%	0.2603%	0.0808%
	Males	0.0720%	0.0024%	0.0135%	0.0235%	0.0475%	0.0946%	0.1625%	0.2823%	0.2696%	0.2805%	0.0911%
	Females	0.0642%	0.0022%	0.0066%	0.0155%	0.0330%	0.0646%	0.1283%	0.2072%	0.2794%	0.2509%	0.0723%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0714%	0.0015%	0.0102%	0.0158%	0.0252%	0.0563%	0.1085%	0.1925%	0.3208%	0.3842%	0.0741%
	Males	0.0754%	0.0023%	0.0125%	0.0201%	0.0247%	0.0652%	0.1209%	0.2106%	0.3783%	0.4577%	0.0851%
	Females	0.0676%	-	0.0076%	0.0115%	0.0257%	0.0481%	0.0977%	0.1771%	0.2778%	0.3380%	0.0654%
Hispanic <sup>d</sup>	Both Sexes	-	-	-	-	-	-	-	-	-	-	
	Males	-	-	-	-	-	-	-	-	-	-	
	Females	-	-	-	-	-	-	-	-	-	-	

<sup>a</sup> US 2008 cancer prevalence counts are based on 2008 cancer prevalence proportions from the SEER registries and 1/1/2008 US population estimates based on the average of 2007 and 2008 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 33 years for 1975-2008 SEER 9 data; 18 years for 1990-2008 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 33 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <33 and >=33. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 20.27  
Cancer of the Oral Cavity and Pharynx (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2008  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <33 <sup>e</sup>	>=33 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	91,329	55,615	36,520	23,518	15,736	10,003	202,684	240,722	12,443	253,165
	Males	62,730	36,405	23,300	14,212	9,267	5,866	134,834	156,987	7,172	164,159
	Females	28,599	19,210	13,220	9,306	6,469	4,137	67,850	83,735	5,271	89,006
White <sup>b</sup>	Both Sexes	78,234	48,573	31,799	21,004	14,028	8,815	175,746	209,525	10,587	220,112
	Males	54,395	32,127	20,614	12,905	8,482	5,352	118,404	138,541	6,142	144,683
	Females	23,839	16,446	11,185	8,099	5,546	3,463	57,342	70,984	4,445	75,429
Black <sup>b</sup>	Both Sexes	7,522	3,959	2,683	1,322	1,047	658	15,113	17,506	620	18,126
	Males	4,854	2,365	1,467	660	450	227	9,101	10,123	237	10,360
	Females	2,668	1,594	1,216	662	597	431	6,012	7,383	383	7,766
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	3,429	2,080	1,505	+	+	+	7,516	+	+	+
	Males	2,163	1,239	896	+	+	+	4,607	+	+	+
	Females	1,266	841	609	+	+	+	2,909	+	+	+
Hispanic <sup>d</sup>	Both Sexes	4,509	2,576	1,701	+	+	+	9,403	+	+	+
	Males	2,849	1,620	982	+	+	+	5,733	+	+	+
	Females	1,660	956	719	+	+	+	3,670	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2008, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)										Age-Adjusted <sup>f</sup>
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0571%	0.0004%	0.0019%	0.0056%	0.0147%	0.0437%	0.1162%	0.1948%	0.2169%	0.2043%	0.0566%
	Males	0.0765%	0.0003%	0.0018%	0.0056%	0.0141%	0.0573%	0.1712%	0.2958%	0.3169%	0.3071%	0.0813%
	Females	0.0380%	0.0005%	0.0020%	0.0056%	0.0153%	0.0301%	0.0636%	0.1033%	0.1380%	0.1475%	0.0357%
White <sup>c</sup>	Both Sexes	0.0608%	0.0005%	0.0017%	0.0053%	0.0133%	0.0414%	0.1189%	0.2035%	0.2305%	0.2139%	0.0580%
	Males	0.0823%	0.0004%	0.0012%	0.0050%	0.0131%	0.0555%	0.1773%	0.3081%	0.3386%	0.3236%	0.0840%
	Females	0.0392%	0.0005%	0.0022%	0.0058%	0.0134%	0.0268%	0.0611%	0.1060%	0.1433%	0.1535%	0.0353%
Black <sup>c</sup>	Both Sexes	0.0366%	0.0007%	0.0029%	0.0054%	0.0121%	0.0364%	0.0970%	0.1570%	0.1496%	0.1227%	0.0438%
	Males	0.0470%	-	0.0039%	0.0076%	0.0087%	0.0418%	0.1396%	0.2499%	0.2257%	0.1631%	0.0616%
	Females	0.0272%	-	0.0019%	0.0033%	0.0150%	0.0317%	0.0616%	0.0854%	0.0974%	0.1039%	0.0303%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0517%	-	0.0017%	0.0072%	0.0224%	0.0605%	0.1081%	0.1509%	0.1654%	0.1579%	0.0519%
	Males	0.0653%	-	0.0025%	0.0071%	0.0232%	0.0778%	0.1466%	0.2244%	0.2148%	0.2356%	0.0698%
	Females	0.0389%	-	-	0.0073%	0.0217%	0.0446%	0.0746%	0.0883%	0.1285%	0.1091%	0.0372%
Hispanic <sup>d</sup>	Both Sexes	-	-	-	-	-	-	-	-	-	-	-
	Males	-	-	-	-	-	-	-	-	-	-	-
	Females	-	-	-	-	-	-	-	-	-	-	-

<sup>a</sup> US 2008 cancer prevalence counts are based on 2008 cancer prevalence proportions from the SEER registries and 1/1/2008 US population estimates based on the average of 2007 and 2008 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 33 years for 1975-2008 SEER 9 data; 18 years for 1990-2008 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 33 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <33 and >=33. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.



Table 21.15  
Cancer of the Ovary (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2008  
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <33 <sup>e</sup>	>=33 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>										
All Races <sup>b</sup>	53,320	31,146	22,768	18,765	15,667	11,406	119,365	158,794	18,784	177,578
White <sup>b</sup>	46,813	27,505	20,077	17,160	14,031	10,237	104,980	140,439	16,156	156,595
Black <sup>b</sup>	3,659	1,918	1,552	971	1,033	778	7,976	10,563	1,697	12,260
Asian/Pacific Islander <sup>c</sup>	2,222	1,349	825	+	+	+	4,778	+	+	+
Hispanic <sup>d</sup>	4,195	2,522	1,479	+	+	+	9,008	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2008, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup> All Ages	
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+		
<u>Race</u>												
All Races <sup>c</sup>	0.0679%	0.0004%	0.0054%	0.0168%	0.0269%	0.0629%	0.1347%	0.1985%	0.2085%	0.1651%	0.0639%	
White <sup>c</sup>	0.0740%	0.0004%	0.0057%	0.0177%	0.0263%	0.0648%	0.1424%	0.2120%	0.2281%	0.1748%	0.0676%	
Black <sup>c</sup>	0.0371%	-	0.0041%	0.0109%	0.0230%	0.0414%	0.0757%	0.1221%	0.1278%	0.1254%	0.0411%	
Asian/Pacific Islander <sup>c</sup>	0.0619%	-	0.0049%	0.0172%	0.0318%	0.0696%	0.1379%	0.1717%	0.1434%	0.1105%	0.0584%	
Hispanic <sup>d</sup>	-	-	-	-	-	-	-	-	-	-	-	

- <sup>a</sup> Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.  
<sup>a</sup> US 2008 cancer prevalence counts are based on 2008 cancer prevalence proportions from the SEER registries and 1/1/2008 US population estimates based on the average of 2007 and 2008 population estimates from the US Bureau of the Census.  
<sup>b c d</sup> Prevalence was calculated using the First Malignant Primary Only for a person.  
<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.  
<sup>e</sup> Maximum limited-duration prevalence: 33 years for 1975-2008 SEER 9 data; 18 years for 1990-2008 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).  
<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.  
<sup>g h i</sup> (g) Cases diagnosed more than 33 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <33 and >=33. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.  
 - Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.  
 + Not available.

Table 22.21  
Cancer of the Pancreas (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2008  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <33 <sup>e</sup>	>=33 <sup>g</sup>	Complete <sup>h</sup>	
Race	Sex											
All Races <sup>b</sup>	Both Sexes	24,630	3,944	1,990	1,127	530	343	32,838	34,305	352	34,657	
	Males	12,189	1,979	854	543	176	158	16,086	16,665	146	16,811	
	Females	12,441	1,965	1,136	584	354	185	16,752	17,640	206	17,846	
White <sup>b</sup>	Both Sexes	21,106	3,222	1,566	874	401	293	27,872	28,983	294	29,277	
	Males	10,791	1,766	663	467	132	140	14,010	14,477	134	14,611	
	Females	10,315	1,456	903	407	269	153	13,862	14,506	160	14,666	
Black <sup>b</sup>	Both Sexes	2,359	467	259	205	91	42	3,373	3,662	129	3,791	
	Males	878	132	93	65	36	10	1,319	1,418	46	1,464	
	Females	1,481	335	166	140	55	32	2,054	2,244	83	2,327	
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	869	176	68	+	+	+	1,135	+	+	+	
	Males	430	61	30	+	+	+	529	+	+	+	
	Females	439	115	38	+	+	+	606	+	+	+	
Hispanic <sup>d</sup>	Both Sexes	1,704	266	179	+	+	+	2,223	+	+	+	
	Males	808	123	49	+	+	+	1,015	+	+	+	
	Females	896	143	130	+	+	+	1,208	+	+	+	

Estimated prevalence percent<sup>a</sup> on January 1, 2008, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	All Ages	Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages	
Race	Sex											
All Races <sup>c</sup>	Both Sexes	0.0098%	-	0.0002%	0.0005%	0.0016%	0.0056%	0.0154%	0.0331%	0.0515%	0.0478%	0.0100%
	Males	0.0097%	-	-	0.0004%	0.0012%	0.0059%	0.0167%	0.0381%	0.0556%	0.0510%	0.0108%
	Females	0.0099%	-	0.0003%	0.0006%	0.0019%	0.0053%	0.0142%	0.0286%	0.0482%	0.0461%	0.0093%
White <sup>c</sup>	Both Sexes	0.0102%	-	0.0002%	0.0005%	0.0016%	0.0052%	0.0154%	0.0340%	0.0514%	0.0477%	0.0100%
	Males	0.0104%	-	-	0.0004%	0.0013%	0.0059%	0.0167%	0.0395%	0.0569%	0.0516%	0.0110%
	Females	0.0101%	-	0.0003%	0.0006%	0.0019%	0.0046%	0.0142%	0.0288%	0.0469%	0.0455%	0.0091%
Black <sup>c</sup>	Both Sexes	0.0083%	-	-	-	0.0022%	0.0066%	0.0182%	0.0342%	0.0514%	0.0500%	0.0106%
	Males	0.0070%	-	-	-	0.0021%	0.0056%	0.0201%	0.0316%	0.0488%	0.0353%	0.0098%
	Females	0.0094%	-	-	-	0.0023%	0.0075%	0.0166%	0.0363%	0.0531%	0.0568%	0.0111%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0086%	-	-	-	0.0011%	0.0064%	0.0134%	0.0257%	0.0504%	0.0469%	0.0092%
	Males	0.0083%	-	-	-	-	0.0056%	0.0146%	0.0331%	0.0477%	0.0502%	0.0096%
	Females	0.0090%	-	-	-	0.0016%	0.0071%	0.0123%	0.0194%	0.0524%	0.0448%	0.0089%
Hispanic <sup>d</sup>	Both Sexes	-	-	-	-	-	-	-	-	-	-	-
	Males	-	-	-	-	-	-	-	-	-	-	-
	Females	-	-	-	-	-	-	-	-	-	-	-

<sup>a</sup> US 2008 cancer prevalence counts are based on 2008 cancer prevalence proportions from the SEER registries and 1/1/2008 US population estimates based on the average of 2007 and 2008 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 33 years for 1975-2008 SEER 9 data; 18 years for 1990-2008 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 33 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <33 and >=33. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 23.15  
Cancer of the Prostate (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2008  
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <33 <sup>e</sup>	>=33 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>										
All Races <sup>b</sup>	947,233	719,432	438,078	142,054	27,477	8,596	2,275,367	2,353,355	2,109	2,355,464
White <sup>b</sup>	780,290	608,406	378,709	129,392	25,198	7,935	1,910,974	1,982,239	2,364	1,984,603
Black <sup>b</sup>	124,196	87,139	48,310	10,111	1,967	553	278,480	284,018	190	284,208
Asian/Pacific Islander <sup>c</sup>	18,979	12,388	5,850	+	+	+	38,742	+	+	+
Hispanic <sup>d</sup>	52,180	35,973	18,196	+	+	+	110,871	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2008, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>											
All Races <sup>c</sup>	1.2583%	0.0002%	0.0002%	0.0003%	0.0013%	0.0755%	0.9874%	4.8815%	10.8493%	12.1247%	1.5298%
White <sup>c</sup>	1.3115%	-	-	-	0.0009%	0.0659%	0.9418%	4.8095%	10.8928%	11.8279%	1.5102%
Black <sup>c</sup>	1.3855%	-	-	-	0.0036%	0.1892%	1.9501%	8.4946%	15.6555%	16.9040%	2.3657%
Asian/Pacific Islander <sup>c</sup>	0.6427%	-	-	-	0.0014%	0.0195%	0.3296%	2.0396%	5.8479%	8.5821%	0.8257%
Hispanic <sup>d</sup>	-	-	-	-	-	-	-	-	-	-	-

<sup>a</sup> US 2008 cancer prevalence counts are based on 2008 cancer prevalence proportions from the SEER registries and 1/1/2008 US population estimates based on the average of 2007 and 2008 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 33 years for 1975-2008 SEER 9 data; 18 years for 1990-2008 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 33 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <33 and >=33. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 24.21  
Cancer of the Stomach (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2008  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <33 <sup>e</sup>	>=33 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	32,747	13,519	7,594	5,180	2,990	1,436	57,258	64,307	1,703	66,010
	Males	19,436	7,758	4,229	2,376	1,625	737	33,163	36,772	967	37,739
	Females	13,311	5,761	3,365	2,804	1,365	699	24,095	27,535	736	28,271
White <sup>b</sup>	Both Sexes	24,954	9,954	5,708	3,804	2,308	1,193	43,027	48,501	1,383	49,884
	Males	15,263	5,748	3,113	1,770	1,289	649	25,507	28,410	755	29,165
	Females	9,691	4,206	2,595	2,034	1,019	544	17,520	20,091	628	20,719
Black <sup>b</sup>	Both Sexes	4,832	1,977	993	715	401	99	8,233	9,093	217	9,310
	Males	2,434	1,047	597	312	208	40	4,234	4,634	122	4,756
	Females	2,398	930	396	403	193	59	3,999	4,459	95	4,554
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	3,024	1,398	889	+	+	+	5,684	+	+	+
	Males	1,719	777	492	+	+	+	3,200	+	+	+
	Females	1,305	621	397	+	+	+	2,484	+	+	+
Hispanic <sup>d</sup>	Both Sexes	4,509	1,708	946	+	+	+	7,549	+	+	+
	Males	2,473	919	439	+	+	+	3,998	+	+	+
	Females	2,036	789	507	+	+	+	3,551	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2008, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0207%	-	-	0.0006%	0.0031%	0.0113%	0.0273%	0.0577%	0.1124%	0.1493%	0.0212%
	Males	0.0238%	-	-	0.0005%	0.0031%	0.0123%	0.0329%	0.0784%	0.1519%	0.2030%	0.0276%
	Females	0.0176%	-	-	0.0008%	0.0031%	0.0102%	0.0219%	0.0390%	0.0812%	0.1196%	0.0162%
White <sup>c</sup>	Both Sexes	0.0172%	-	-	0.0006%	0.0028%	0.0094%	0.0221%	0.0475%	0.0875%	0.1106%	0.0167%
	Males	0.0202%	-	-	0.0005%	0.0029%	0.0100%	0.0273%	0.0674%	0.1203%	0.1500%	0.0222%
	Females	0.0141%	-	-	0.0008%	0.0027%	0.0086%	0.0169%	0.0290%	0.0610%	0.0889%	0.0124%
Black <sup>c</sup>	Both Sexes	0.0198%	-	-	-	0.0039%	0.0154%	0.0399%	0.0776%	0.1194%	0.1852%	0.0264%
	Males	0.0206%	-	-	-	0.0042%	0.0190%	0.0428%	0.0958%	0.1477%	0.2303%	0.0320%
	Females	0.0190%	-	-	-	0.0036%	0.0123%	0.0375%	0.0636%	0.1001%	0.1643%	0.0227%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0431%	-	-	0.0009%	0.0040%	0.0185%	0.0487%	0.1097%	0.2646%	0.4126%	0.0464%
	Males	0.0497%	-	-	-	0.0030%	0.0201%	0.0607%	0.1420%	0.3620%	0.5527%	0.0607%
	Females	0.0368%	-	-	-	0.0049%	0.0170%	0.0383%	0.0822%	0.1919%	0.3245%	0.0358%
Hispanic <sup>d</sup>	Both Sexes	-	-	-	-	-	-	-	-	-	-	-
	Males	-	-	-	-	-	-	-	-	-	-	-
	Females	-	-	-	-	-	-	-	-	-	-	-

<sup>a</sup> US 2008 cancer prevalence counts are based on 2008 cancer prevalence proportions from the SEER registries and 1/1/2008 US population estimates based on the average of 2007 and 2008 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 33 years for 1975-2008 SEER 9 data; 18 years for 1990-2008 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 33 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <33 and >=33. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 25.15  
Cancer of the Testis (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2008  
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <33 <sup>e</sup>	>=33 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>										
All Races <sup>b</sup>	39,225	36,242	31,402	28,539	21,720	15,555	126,678	184,341	16,546	200,887
White <sup>b</sup>	36,888	33,945	29,897	27,400	21,006	15,058	119,463	175,171	15,886	191,057
Black <sup>b</sup>	923	1,006	600	372	320	225	2,829	3,687	157	3,844
Asian/Pacific Islander <sup>c</sup>	687	615	477	+	+	+	2,034	+	+	+
Hispanic <sup>d</sup>	5,072	3,666	2,714	+	+	+	12,899	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2008, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>											
All Races <sup>c</sup>	0.0704%	0.0019%	0.0060%	0.0638%	0.1438%	0.1563%	0.0913%	0.0369%	0.0161%	0.0066%	0.0695%
White <sup>c</sup>	0.0852%	0.0019%	0.0069%	0.0780%	0.1783%	0.1884%	0.1069%	0.0433%	0.0188%	0.0079%	0.0842%
Black <sup>c</sup>	0.0152%	0.0013%	0.0022%	0.0120%	0.0271%	0.0367%	0.0261%	0.0080%	-	-	0.0157%
Asian/Pacific Islander <sup>c</sup>	0.0244%	0.0032%	0.0047%	0.0205%	0.0424%	0.0505%	0.0389%	0.0092%	0.0049%	-	0.0234%
Hispanic <sup>d</sup>	-	-	-	-	-	-	-	-	-	-	-

<sup>a</sup> US 2008 cancer prevalence counts are based on 2008 cancer prevalence proportions from the SEER registries and 1/1/2008 US population estimates based on the average of 2007 and 2008 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 33 years for 1975-2008 SEER 9 data; 18 years for 1990-2008 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 33 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <33 and >=33. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 26.21  
Cancer of the Thyroid (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2008  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <33 <sup>e</sup>	>=33 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	128,157	83,662	60,394	43,715	34,679	28,867	320,828	417,546	40,857	458,403
	Males	28,749	18,752	13,617	9,327	7,535	6,022	71,016	91,582	9,370	100,952
	Females	99,408	64,910	46,777	34,388	27,144	22,845	249,812	325,964	31,487	357,451
White <sup>b</sup>	Both Sexes	112,068	73,460	53,439	39,316	31,124	26,013	282,053	369,232	37,010	406,242
	Males	25,703	16,999	12,387	8,664	6,782	5,516	64,055	82,881	7,901	90,782
	Females	86,365	56,461	41,052	30,652	24,342	20,497	217,998	286,351	29,109	315,460
Black <sup>b</sup>	Both Sexes	7,823	4,611	3,218	1,958	1,609	1,339	18,272	22,598	1,690	24,288
	Males	1,489	760	600	340	307	177	3,281	3,991	188	4,179
	Females	6,334	3,851	2,618	1,618	1,302	1,162	14,991	18,607	1,502	20,109
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	6,296	4,245	2,761	+	+	+	14,548	+	+	+
	Males	1,112	811	446	+	+	+	2,574	+	+	+
	Females	5,184	3,434	2,315	+	+	+	11,974	+	+	+
Hispanic <sup>d</sup>	Both Sexes	13,241	8,258	5,457	+	+	+	29,246	+	+	+
	Males	2,232	1,447	875	+	+	+	4,937	+	+	+
	Females	11,009	6,811	4,582	+	+	+	24,309	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2008, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	All Ages	Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages	
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0935%	0.0001%	0.0044%	0.0328%	0.1045%	0.1611%	0.1836%	0.1870%	0.1575%	0.1073%	0.0924%
	Males	0.0407%	-	0.0019%	0.0091%	0.0333%	0.0611%	0.0853%	0.1060%	0.1053%	0.0730%	0.0418%
	Females	0.1455%	-	0.0071%	0.0579%	0.1779%	0.2614%	0.2777%	0.2604%	0.1987%	0.1263%	0.1415%
White <sup>c</sup>	Both Sexes	0.1008%	0.0002%	0.0048%	0.0359%	0.1156%	0.1760%	0.1928%	0.1918%	0.1603%	0.1047%	0.0983%
	Males	0.0452%	-	0.0021%	0.0099%	0.0366%	0.0681%	0.0930%	0.1122%	0.1119%	0.0721%	0.0451%
	Females	0.1566%	-	0.0077%	0.0643%	0.2005%	0.2884%	0.2917%	0.2659%	0.1994%	0.1227%	0.1519%
Black <sup>c</sup>	Both Sexes	0.0437%	-	0.0014%	0.0120%	0.0434%	0.0749%	0.0999%	0.1232%	0.1052%	0.0864%	0.0490%
	Males	0.0162%	-	0.0012%	0.0025%	0.0116%	0.0266%	0.0395%	0.0550%	0.0643%	0.0671%	0.0206%
	Females	0.0686%	-	0.0015%	0.0212%	0.0713%	0.1172%	0.1502%	0.1756%	0.1333%	0.0954%	0.0724%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0939%	-	0.0048%	0.0319%	0.0950%	0.1454%	0.1892%	0.1983%	0.1734%	0.1330%	0.0918%
	Males	0.0348%	-	-	0.0090%	0.0325%	0.0464%	0.0701%	0.0974%	0.0869%	0.0793%	0.0360%
	Females	0.1493%	-	0.0090%	0.0549%	0.1542%	0.2360%	0.2929%	0.2842%	0.2380%	0.1667%	0.1410%
Hispanic <sup>d</sup>	Both Sexes	-	-	-	-	-	-	-	-	-	-	-
	Males	-	-	-	-	-	-	-	-	-	-	-
	Females	-	-	-	-	-	-	-	-	-	-	-

<sup>a</sup> US 2008 cancer prevalence counts are based on 2008 cancer prevalence proportions from the SEER registries and 1/1/2008 US population estimates based on the average of 2007 and 2008 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 33 years for 1975-2008 SEER 9 data; 18 years for 1990-2008 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 33 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <33 and >=33. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 27.21  
Cancer of the Urinary Bladder (Invasive and *In Situ*)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2008  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <33 <sup>e</sup>	>=33 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	189,650	130,423	87,562	54,145	33,652	18,975	442,780	522,368	15,060	537,428
	Males	142,414	97,205	64,930	40,306	24,192	13,434	331,088	388,564	9,765	398,329
	Females	47,236	33,218	22,632	13,839	9,460	5,541	111,692	133,804	5,295	139,099
White <sup>b</sup>	Both Sexes	175,220	122,390	82,567	51,825	31,957	18,173	413,072	489,067	11,769	500,836
	Males	132,507	91,539	61,226	38,582	22,942	12,931	310,059	364,922	7,507	372,429
	Females	42,713	30,851	21,341	13,243	9,015	5,242	103,013	124,145	4,262	128,407
Black <sup>b</sup>	Both Sexes	7,948	4,594	2,988	1,472	1,025	414	16,636	18,804	239	19,043
	Males	5,031	3,008	2,223	1,091	748	246	11,107	12,704	193	12,897
	Females	2,917	1,586	765	381	277	168	5,529	6,100	46	6,146
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	3,223	1,686	1,141	+	+	+	6,470	+	+	+
	Males	2,514	1,277	828	+	+	+	4,935	+	+	+
	Females	709	409	313	+	+	+	1,535	+	+	+
Hispanic <sup>d</sup>	Both Sexes	6,795	3,952	2,510	+	+	+	14,302	+	+	+
	Males	5,130	2,937	1,881	+	+	+	10,712	+	+	+
	Females	1,665	1,015	629	+	+	+	3,590	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2008, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.1159%	0.0001%	0.0004%	0.0015%	0.0068%	0.0290%	0.1076%	0.3525%	0.7370%	0.9946%	0.1203%
	Males	0.1761%	-	0.0005%	0.0021%	0.0096%	0.0436%	0.1689%	0.5748%	1.2784%	2.0104%	0.2128%
	Females	0.0567%	-	-	0.0009%	0.0039%	0.0143%	0.0490%	0.1512%	0.3100%	0.4323%	0.0521%
White <sup>c</sup>	Both Sexes	0.1379%	0.0001%	0.0003%	0.0018%	0.0082%	0.0325%	0.1217%	0.4047%	0.8423%	1.0823%	0.1356%
	Males	0.2083%	-	0.0006%	0.0024%	0.0113%	0.0479%	0.1870%	0.6519%	1.4481%	2.2112%	0.2381%
	Females	0.0673%	-	-	0.0011%	0.0049%	0.0164%	0.0571%	0.1742%	0.3533%	0.4602%	0.0586%
Black <sup>c</sup>	Both Sexes	0.0406%	-	-	-	0.0033%	0.0178%	0.0609%	0.1717%	0.3201%	0.4777%	0.0577%
	Males	0.0579%	-	-	-	0.0048%	0.0271%	0.1047%	0.2857%	0.5202%	0.8495%	0.0974%
	Females	0.0250%	-	-	-	0.0019%	0.0096%	0.0244%	0.0840%	0.1830%	0.3047%	0.0319%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0502%	-	-	0.0007%	0.0024%	0.0141%	0.0449%	0.1254%	0.3164%	0.5744%	0.0547%
	Males	0.0787%	-	-	-	0.0032%	0.0231%	0.0771%	0.2114%	0.5732%	1.0735%	0.0983%
	Females	0.0235%	-	-	-	0.0016%	0.0059%	0.0169%	0.0522%	0.1243%	0.2607%	0.0228%
Hispanic <sup>d</sup>	Both Sexes	-	-	-	-	-	-	-	-	-	-	-
	Males	-	-	-	-	-	-	-	-	-	-	-
	Females	-	-	-	-	-	-	-	-	-	-	-

<sup>a</sup> US 2008 cancer prevalence counts are based on 2008 cancer prevalence proportions from the SEER registries and 1/1/2008 US population estimates based on the average of 2007 and 2008 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 33 years for 1975-2008 SEER 9 data; 18 years for 1990-2008 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 33 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <33 and >=33. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 28.12  
Childhood Cancer (Invasive, <20 Years old at diagnosis)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2008  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <33 <sup>e</sup>	>=33 <sup>f</sup>	Complete <sup>g</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	58,522	51,832	45,147	39,508	33,666	28,341	179,963	271,858	+	+
	Males	31,328	26,794	23,733	20,786	17,027	13,701	94,953	140,000	+	+
	Females	27,194	25,038	21,414	18,722	16,639	14,640	85,010	131,858	+	+
White <sup>b</sup>	Both Sexes	48,265	43,997	37,972	33,839	29,595	25,098	151,158	231,868	+	+
	Males	26,095	22,716	19,963	17,928	15,058	12,245	80,090	119,939	+	+
	Females	22,170	21,281	18,009	15,911	14,537	12,853	71,068	111,929	+	+
Black <sup>b</sup>	Both Sexes	6,456	4,944	4,472	3,658	2,628	2,280	18,093	25,644	+	+
	Males	3,304	2,517	2,266	1,858	1,242	976	9,231	12,654	+	+
	Females	3,152	2,427	2,206	1,800	1,386	1,304	8,862	12,990	+	+
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	2,280	1,849	1,570	+	+	+	6,519	+	+	+
	Males	1,202	996	846	+	+	+	3,439	+	+	+
	Females	1,078	853	724	+	+	+	3,080	+	+	+
Hispanic <sup>d</sup>	Both Sexes	11,317	8,598	6,535	+	+	+	29,844	+	+	+
	Males	6,408	4,539	3,531	+	+	+	16,402	+	+	+
	Females	4,909	4,059	3,004	+	+	+	13,442	+	+	+

<sup>a</sup> US 2008 cancer prevalence counts are based on 2008 cancer prevalence proportions from the SEER registries and 1/1/2008 US population estimates based on the average of 2007 and 2008 population estimates from the US Bureau of the Census.

<sup>b c d</sup> Prevalence was calculated using the First Malignant Primary Only for a person. Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 33 years for 1975-2008 SEER 9 data; 18 years for 1990-2008 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f g</sup> (f) Cases diagnosed more than 33 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (g) Complete prevalence is obtained by summing 0 to <33 and >=33.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.



Table 29.7  
U.S. Childhood Cancer Survivors at January 1, 2008  
Number of People Previously Diagnosed with Cancer as Children (Ages 0-19 Years) in the United States and Alive January 1, 2008  
By International Classification of Childhood Cancer (ICCC)<sup>a</sup> Selected Group and Subgroup, Age Group, and Sex

Site	Sex	Complete Prevalence Counts by Age at Prevalence							All Ages		
		0-19	20-29	30-39	40-49	50-59	60+	All Ages	Diagnosed between 1975-2007	Diagnosed before 1975 Count	% <sup>b</sup>
All Sites	Both	108,674	86,684	59,026	52,916	31,051	14,143	352,494	277,123	75,371	21
	Male	58,583	45,136	28,547	23,218	13,499	7,741	176,724	142,911	33,813	19
	Female <sup>c</sup>	50,091	41,548	30,479	29,698	17,552	6,402	175,770	134,212	41,558	24
I Leukemia	Both	33,969	17,955	9,704	3,714	1,061	91	66,494	62,215	4,279	6
	Male	18,631	9,597	4,911	1,896	510	42	35,587	33,619	1,968	6
	Female	15,338	8,358	4,793	1,818	551	49	30,907	28,596	2,311	7
Acute lymphocytic leukemia <sup>d</sup>	Both	28,843	14,901	8,472	2,890	218	0	55,324	52,032	3,292	6
	Male	15,958	8,014	4,425	1,627	157	0	30,181	28,421	1,760	6
	Female	12,885	6,887	4,047	1,263	61	0	25,143	23,611	1,532	6
I(b) Acute myeloid leukemia	Both	3,698	1,937	986	492	164	0	7,277	6,798	479	7
	Male	1,868	959	391	256	88	0	3,562	3,312	250	7
	Female	1,830	978	595	236	76	0	3,715	3,486	229	6
II(a) Hodgkin lymphoma	Both	4,156	8,887	7,851	7,587	3,882	742	33,105	27,400	5,705	17
	Male	2,438	4,326	3,921	3,689	1,953	402	16,729	13,657	3,072	18
	Female	1,718	4,561	3,930	3,898	1,929	340	16,376	13,743	2,633	16
II(b,c,e) Non-Hodgkin lymphoma	Both	6,387	6,365	4,059	2,732	1,578	907	22,028	18,356	3,672	17
	Male	4,182	4,270	2,808	1,564	799	544	14,167	12,179	1,988	14
	Female	2,205	2,095	1,251	1,168	779	363	7,861	6,177	1,684	21
III CNS and misc intracranial & intraspinal neoplasms	Both	19,603	13,781	8,499	8,242	3,877	1,030	55,032	43,583	11,449	21
	Male	10,917	7,684	4,466	3,962	1,731	461	29,221	23,957	5,264	18
	Female	8,686	6,097	4,033	4,280	2,146	569	25,811	19,626	6,185	24
IV Neuroblastoma & other peripheral nervous cell tumor	Both	9,215	3,368	1,895	1,382	1,385	511	17,756	13,902	3,854	22
	Male	4,787	1,609	816	508	351	141	8,212	7,025	1,187	14
	Female	4,428	1,759	1,079	874	1,034	370	9,544	6,877	2,667	28
VI Renal tumors	Both	7,794	4,327	3,527	3,911	1,865	510	21,934	14,435	7,499	34
	Male	3,831	2,018	1,642	2,090	1,043	303	10,927	6,810	4,117	38
	Female	3,963	2,309	1,885	1,821	822	207	11,007	7,625	3,382	31
VIII Malignant bone tumors	Both	3,371	4,067	2,898	2,470	2,016	597	15,419	11,755	3,664	24
	Male	1,844	2,335	1,597	1,141	906	336	8,159	6,480	1,679	21
	Female	1,527	1,732	1,301	1,329	1,110	261	7,260	5,275	1,985	27
VIII(a) Osteosarcoma	Both	1,661	2,220	1,627	1,230	1,072	668	8,478	6,258	2,220	26
	Male	884	1,248	906	549	425	362	4,374	3,383	991	23
	Female	777	972	721	681	647	306	4,104	2,875	1,229	30
VIII(c) Ewing tumor & related sarcomas of bone	Both	1,266	1,263	835	572	362	63	4,361	3,743	618	14
	Male	709	734	389	283	160	18	2,293	2,046	247	11
	Female	557	529	446	289	202	45	2,068	1,697	371	18
IX Soft tissue & other extraosseous sarcomas	Both	7,120	6,259	4,034	4,789	3,828	3,859	29,889	19,404	10,485	35
	Male	3,819	3,439	2,056	2,219	1,752	2,155	15,440	10,263	5,177	34
	Female	3,301	2,820	1,978	2,570	2,076	1,704	14,449	9,141	5,308	37
X Germ cell & trophoblastic tumors & neoplasms of gonads	Both	4,775	7,246	5,528	6,538	6,093	5,787	35,967	21,455	14,512	40
	Male	2,549	4,614	3,272	3,387	3,053	2,584	19,459	12,706	6,753	35
	Female	2,226	2,632	2,256	3,151	3,040	3,203	16,508	8,749	7,759	47

<sup>a</sup> International Classification of Childhood Cancer is based on ICD-O-3. Steliarova-Foucher E, Stiller C, Lacour B, Kaatsch P. International Classification of Childhood Cancer, Third Edition. Cancer. April 1, 2005; Vol 103, No. 7, pg 1457-1467.  
<sup>b</sup> Percent diagnosed before 1975 of all ages complete prevalence counts.  
<sup>c</sup> All sites females excludes breast cancer.  
<sup>d</sup> SEER site recode Acute Lymphocytic Leukemia was used instead of the ICCC classification lymphocytic leukemia (Ia) because of statistical modeling requirements.  
For more information regarding the modeling of the prevalence estimates, see Mariotto et al. Long-Term Survivors of Childhood Cancers in the United States. Cancer Epidemiol Biomarkers Prev 2009, page 1034.