

Table 33.1
Five-Year Cause-Specific Survival^a(%) for Malignant Cancers, 2004-2010
Both Sexes by Race/Ethnicity and Primary Site

Primary Site	White			Black			Asian/ Pacific Islander			American Indian/ Alaska Native ^b			Hispanic ^c		
	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE
All Sites	1,779,294	67.1	0.0	231,453	61.5	0.1	134,840	65.9	0.1	8,894	58.9	0.6	212,954	67.3	0.1
Oral Cavity & Pharynx	41,538	68.3	0.3	4,837	48.9	0.8	3,455	70.3	0.9	215	61.7	3.9	3,846	63.8	0.9
Esophagus	16,999	20.4	0.4	2,321	14.8	0.9	928	19.6	1.7	103	20.2	4.6	1,494	17.9	1.2
Stomach	24,504	28.5	0.3	4,809	30.1	0.8	4,980	38.8	0.8	267	21.7	3.0	6,106	29.0	0.7
Colon & Rectum	166,444	65.6	0.1	24,921	59.2	0.4	16,145	69.2	0.4	1,065	61.1	1.8	20,284	65.1	0.4
Liver & Intrahepatic Bile Duct	25,769	19.7	0.3	4,648	15.2	0.7	6,248	25.8	0.7	360	17.4	2.5	6,851	20.6	0.6
Pancreas	43,947	6.9	0.2	6,685	7.0	0.4	3,717	8.6	0.6	266	3.4	1.6	5,404	7.9	0.5
Larynx	13,096	66.8	0.5	2,454	58.9	1.2	488	71.4	2.5	60	37.1	9.7	1,267	63.8	1.7
Lung & Bronchus	219,621	19.3	0.1	29,675	16.1	0.3	14,800	20.5	0.4	971	12.8	1.4	14,528	18.6	0.4
Melanoma of the Skin	88,262	89.3	0.1	461	70.5	2.5	563	74.2	2.2	115	85.9	3.5	2,953	83.6	0.8
Breast	260,490	88.6	0.1	34,132	79.4	0.3	23,752	91.3	0.2	1,269	86.0	1.2	31,842	87.4	0.2
Cervix uteri	16,700	71.4	0.4	3,079	61.5	1.0	1,916	72.6	1.2	150	62.2	4.7	5,024	74.6	0.7
Corpus & Uterus, NOS	54,019	83.4	0.2	6,175	63.8	0.7	4,867	83.4	0.6	323	83.9	2.5	6,877	82.0	0.5
Ovary ^d	28,024	44.8	0.4	2,773	37.4	1.1	2,445	56.9	1.2	165	46.9	4.5	3,940	51.4	1.0
Prostate	281,715	94.0	0.1	51,820	92.4	0.1	16,472	94.2	0.2	922	89.5	1.2	31,662	93.0	0.2
Testis	13,939	95.7	0.2	431	91.8	1.4	534	94.5	1.1	103	94.0	2.7	3,244	93.0	0.5
Urinary Bladder	77,420	78.9	0.2	4,646	67.5	0.8	3,272	79.6	0.8	182	69.5	3.8	5,036	78.4	0.7
Kidney & Renal Pelvis	56,090	74.0	0.2	7,343	75.2	0.6	3,212	72.7	0.9	518	69.9	2.3	8,684	75.2	0.5
Brain & Nervous System	28,365	33.7	0.3	2,251	43.3	1.2	1,660	43.8	1.4	135	50.2	4.9	4,329	47.1	0.9
Thyroid	48,699	97.0	0.1	3,883	96.2	0.3	5,852	96.2	0.3	254	95.8	1.3	8,792	97.2	0.2
Hodgkin Lymphoma	12,211	87.5	0.3	1,803	84.2	1.0	693	87.4	1.4	44	94.0	4.2	2,367	85.7	0.8
Non-Hodgkin Lymphoma	77,126	70.9	0.2	7,427	65.5	0.6	5,664	68.2	0.7	326	68.6	3.0	10,618	68.5	0.5
Myeloma	20,706	49.1	0.5	5,464	51.6	0.9	1,497	49.7	1.7	116	56.0	5.4	3,090	48.6	1.2
Leukemia	50,086	61.0	0.2	4,653	55.9	0.9	3,267	56.0	1.0	240	61.7	3.5	8,082	62.4	0.6
Mesothelioma	3,968	8.7	0.6	230	14.3	3.0	119	6.4	3.0	-	-	-	432	12.6	2.1

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2004-2010 with follow-up through 2011.

^b Data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^c Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

^d Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

- Statistic could not be calculated due to less than 25 cases in the time interval.

Table 33.2
Five-Year Cause-Specific Survival^a(%) for Malignant Cancers, 2004-2010
Males by Race/Ethnicity and Primary Site

Primary Site	White			Black			Asian/ Pacific Islander			American Indian/ Alaska Native ^b			Hispanic ^c		
	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE
All Sites	923,039	67.1	0.1	123,664	63.6	0.2	63,210	61.6	0.2	4,251	56.2	0.9	105,520	65.8	0.2
Oral Cavity & Pharynx	29,518	68.3	0.3	3,337	44.5	1.0	2,236	67.3	1.2	149	60.6	4.9	2,599	60.4	1.1
Esophagus	13,544	20.6	0.4	1,610	13.5	1.1	702	19.8	1.9	69	17.5	5.7	1,201	17.4	1.4
Stomach	15,173	26.8	0.4	2,680	27.3	1.1	2,771	39.2	1.1	157	16.8	3.7	3,496	28.5	0.9
Colon & Rectum	86,058	66.2	0.2	11,820	57.6	0.6	8,292	69.4	0.6	535	61.9	2.6	10,996	64.9	0.6
Liver & Intrahepatic Bile Duct	18,804	19.9	0.4	3,434	13.7	0.9	4,353	26.3	0.8	259	20.5	3.0	4,971	20.2	0.8
Pancreas	22,032	6.6	0.2	3,085	6.7	0.6	1,795	8.2	0.8	134	0.0	0.0	2,644	6.7	0.6
Larynx	10,534	67.6	0.5	1,960	58.9	1.3	428	70.8	2.6	48	48.2	11.2	1,088	63.9	1.8
Lung & Bronchus	115,317	16.8	0.1	16,649	13.7	0.3	8,574	18.0	0.5	500	12.3	1.9	7,946	14.9	0.5
Melanoma of the Skin	49,500	87.0	0.2	195	60.3	4.1	279	68.2	3.4	57	78.7	6.1	1,250	77.8	1.4
Prostate	281,715	94.0	0.1	51,820	92.4	0.1	16,472	94.2	0.2	922	89.5	1.2	31,662	93.0	0.2
Testis	13,939	95.7	0.2	431	91.8	1.4	534	94.5	1.1	103	94.0	2.7	3,244	93.0	0.5
Urinary Bladder	58,633	80.3	0.2	3,037	72.1	1.0	2,482	80.9	0.9	146	72.4	4.3	3,741	80.5	0.8
Kidney & Renal Pelvis	34,726	73.5	0.3	4,364	74.3	0.8	2,013	72.5	1.2	298	69.6	3.0	5,191	73.7	0.7
Brain & Nervous System	15,910	32.9	0.4	1,123	40.4	1.7	894	42.1	2.0	70	52.9	6.4	2,414	46.2	1.2
Thyroid	11,454	94.4	0.2	699	93.1	1.1	1,144	94.6	0.8	55	92.2	3.8	1,560	94.5	0.7
Hodgkin Lymphoma	6,640	87.0	0.5	962	81.7	1.5	377	86.9	1.9	-	-	-	1,301	83.7	1.2
Non-Hodgkin Lymphoma	41,591	70.1	0.3	4,024	62.2	0.9	3,049	67.0	1.0	174	68.4	4.4	5,705	66.3	0.7
Myeloma	11,577	51.0	0.6	2,652	50.8	1.3	788	47.6	2.4	59	52.5	7.8	1,642	50.3	1.7
Leukemia	28,800	61.7	0.3	2,511	56.6	1.2	1,810	57.2	1.3	137	62.5	4.6	4,566	63.2	0.8
Mesothelioma	3,037	6.8	0.6	160	10.9	3.1	84	3.5	3.1	-	-	-	307	10.0	2.2

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2004-2010 with follow-up through 2011.

^b Data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^c Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic could not be calculated due to less than 25 cases in the time interval.

Table 33.3
Five-Year Cause-Specific Survival^a(%) for Malignant Cancers, 2004-2010
Females by Race/Ethnicity and Primary Site

Primary Site	White			Black			Asian/ Pacific Islander			American Indian/ Alaska Native ^b			Hispanic ^c		
	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE
All Sites	856,255	67.1	0.1	107,789	59.0	0.2	71,630	69.7	0.2	4,643	61.3	0.8	107,434	68.9	0.2
Oral Cavity & Pharynx	12,020	68.4	0.5	1,500	58.4	1.5	1,219	75.7	1.4	66	63.6	6.8	1,247	70.9	1.5
Esophagus	3,455	19.9	0.8	711	17.9	1.8	226	18.8	3.5	34	24.6	8.2	293	20.5	2.8
Stomach	9,331	31.4	0.6	2,129	33.5	1.2	2,209	38.2	1.2	110	29.5	4.8	2,610	29.7	1.1
Colon & Rectum	80,386	65.0	0.2	13,101	60.6	0.5	7,853	69.0	0.6	530	60.3	2.5	9,288	65.4	0.6
Liver & Intrahepatic Bile Duct	6,965	19.0	0.6	1,214	19.0	1.5	1,895	24.6	1.3	101	8.2	4.3	1,880	21.5	1.2
Pancreas	21,915	7.2	0.2	3,600	7.3	0.6	1,922	9.0	0.8	132	5.6	2.2	2,760	8.9	0.7
Larynx	2,562	63.6	1.1	494	59.3	2.6	60	76.0	6.4	-	-	-	179	63.1	4.4
Lung & Bronchus	104,304	22.0	0.2	13,026	19.0	0.4	6,226	23.8	0.7	471	13.5	2.0	6,582	23.1	0.6
Melanoma of the Skin	38,762	92.3	0.2	266	77.9	3.1	284	80.0	2.7	58	92.8	3.5	1,703	87.8	0.9
Breast	258,579	88.6	0.1	33,822	79.5	0.3	23,624	91.4	0.2	1,264	85.9	1.2	31,680	87.4	0.2
Cervix uteri	16,700	71.4	0.4	3,079	61.5	1.0	1,916	72.6	1.2	150	62.2	4.7	5,024	74.6	0.7
Corpus & Uterus, NOS	54,019	83.4	0.2	6,175	63.8	0.7	4,867	83.4	0.6	323	83.9	2.5	6,877	82.0	0.5
Ovary ^d	28,024	44.8	0.4	2,773	37.4	1.1	2,445	56.9	1.2	165	46.9	4.5	3,940	51.4	1.0
Urinary Bladder	18,787	74.8	0.4	1,609	58.8	1.4	790	75.5	1.8	36	57.8	8.3	1,295	72.3	1.4
Kidney & Renal Pelvis	21,364	74.8	0.3	2,979	76.5	0.9	1,199	73.1	1.5	220	70.5	3.4	3,493	77.3	0.8
Brain & Nervous System	12,455	34.7	0.5	1,128	46.2	1.7	766	45.7	2.1	65	46.8	7.3	1,915	48.2	1.3
Thyroid	37,245	97.8	0.1	3,184	96.9	0.3	4,708	96.6	0.3	199	96.8	1.3	7,232	97.8	0.2
Hodgkin Lymphoma	5,571	88.0	0.5	841	87.2	1.3	316	88.2	2.0	-	-	-	1,066	88.2	1.1
Non-Hodgkin Lymphoma	35,535	71.9	0.3	3,403	69.4	0.9	2,615	69.6	1.0	152	68.6	4.2	4,913	71.1	0.7
Myeloma	9,129	46.8	0.7	2,812	52.5	1.3	709	52.1	2.5	57	59.7	7.4	1,448	46.8	1.8
Leukemia	21,286	60.0	0.4	2,142	55.0	1.3	1,457	54.6	1.5	103	60.8	5.2	3,516	61.3	0.9
Mesothelioma	931	14.8	1.5	70	25.6	6.6	35	12.2	6.5	-	-	-	125	19.4	4.5

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2004-2010 with follow-up through 2011.

^b Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^c Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

^d Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

- Statistic could not be calculated due to less than 25 cases in the time interval.

Table 33.4

Cancer of the Female Breast

5-Year Cause-Specific Survival^a and Case Distribution by Race/Ethnicity
SEER 18 Registries, 2004-2010

Race/Ethnicity	All Ages				Ages <50				Ages 50+			
	Case Distribution		Survival		Case Distribution		Survival		Case Distribution		Survival	
	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE
All Races	319,890	100.0	87.9	0.1	79,282	100.0	89.6	0.1	240,608	100.0	87.3	0.1
White	258,579	80.8	88.6	0.1	59,900	75.6	90.8	0.1	198,679	82.6	88.0	0.1
White Non-Hispanic	228,009	71.3	88.8	0.1	48,720	61.5	91.5	0.2	179,289	74.5	88.1	0.1
White Hispanic	30,570	9.6	87.4	0.2	11,180	14.1	87.9	0.4	19,390	8.1	87.2	0.3
Black	33,822	10.6	79.5	0.3	10,429	13.2	80.6	0.5	23,393	9.7	79.0	0.3
American Indian/Alaska Native ^c	1,264	0.4	85.9	1.2	409	0.5	85.2	2.1	855	0.4	86.3	1.5
Asian	21,873	6.8	91.7	0.2	7,300	9.2	92.9	0.4	14,573	6.1	91.1	0.3
Asian Indian, Pakistani	1,849	0.6	89.9	0.9	724	0.9	92.1	1.2	1,125	0.5	88.3	1.3
Chinese	4,339	1.4	92.3	0.5	1,526	1.9	93.1	0.8	2,813	1.2	91.9	0.6
Filipino	6,008	1.9	90.1	0.5	1,745	2.2	91.0	0.9	4,263	1.8	89.7	0.6
Japanese	3,200	1.0	93.4	0.5	714	0.9	95.9	1.0	2,486	1.0	92.6	0.6
Korean	1,617	0.5	91.9	0.8	669	0.8	93.7	1.1	948	0.4	90.6	1.2
Vietnamese	1,426	0.4	91.1	1.0	604	0.8	92.7	1.3	822	0.3	89.8	1.4
Other Asian ^d	3,434	1.1	93.2	0.5	1,318	1.7	93.7	0.8	2,116	0.9	92.9	0.7
Pacific Islander	1,751	0.5	87.2	1.0	521	0.7	84.5	2.0	1,230	0.5	88.4	1.2
Hawaiian	1,125	0.4	89.9	1.1	315	0.4	88.7	2.2	810	0.3	90.4	1.3
Other Pacific Islander ^e	626	0.2	82.0	2.0	206	0.3	77.7	3.8	420	0.2	84.2	2.4
Hispanic ^f	31,680	9.9	87.4	0.2	11,562	14.6	87.8	0.4	20,118	8.4	87.1	0.3

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2004-2010 with follow-up through 2011.

^b Percent of cases do not sum to 100 because All Races includes unknown race and other race which are not shown separately.

^c Survival estimates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^d Other Asians include Laotian, Hmong, Kampuchean, Thai, and Asians, Not Otherwise Specified (NOS).

^e Other Pacific Islanders include Micronesian, Chamorro, Guamanian, Polynesian, Tahitian, Samoan, Tongan, Melanesian, Fiji Islander, New Guinean, and Pacific Islander, NOS.

^f Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic not shown due to fewer than 25 cases during the time period.

Table 33.5

Cancer of the Colon and Rectum

5-Year Cause-Specific Survival^a and Case Distribution by Race/Ethnicity
SEER 18 Registries, 2004-2010

Race/Ethnicity	Both Sexes				Males				Females			
	Case Distribution		Survival		Case Distribution		Survival		Case Distribution		Survival	
	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE
All Races	210,577	100.0	65.3	0.1	107,829	100.0	65.6	0.2	102,748	100.0	64.9	0.2
White	166,444	79.0	65.6	0.1	86,058	79.8	66.2	0.2	80,386	78.2	65.0	0.2
White Non-Hispanic	146,882	69.8	65.7	0.1	75,424	69.9	66.3	0.2	71,458	69.5	65.0	0.2
White Hispanic	19,562	9.3	65.0	0.4	10,634	9.9	64.8	0.6	8,928	8.7	65.4	0.6
Black	24,921	11.8	59.2	0.4	11,820	11.0	57.6	0.6	13,101	12.8	60.6	0.5
American Indian/Alaska Native ^c	1,065	0.5	61.1	1.8	535	0.5	61.9	2.6	530	0.5	60.3	2.5
Asian	15,198	7.2	69.8	0.4	7,745	7.2	70.2	0.6	7,453	7.3	69.4	0.6
Asian Indian, Pakistani	752	0.4	72.7	2.0	439	0.4	74.7	2.5	313	0.3	70.3	3.2
Chinese	3,486	1.7	70.7	0.9	1,733	1.6	71.8	1.3	1,753	1.7	69.6	1.3
Filipino	3,289	1.6	68.5	1.0	1,689	1.6	68.7	1.4	1,600	1.6	68.2	1.4
Japanese	2,896	1.4	69.1	1.0	1,466	1.4	68.0	1.4	1,430	1.4	70.3	1.4
Korean	1,490	0.7	68.8	1.4	719	0.7	69.5	2.0	771	0.8	68.2	2.0
Vietnamese	1,284	0.6	66.9	1.6	700	0.6	66.7	2.2	584	0.6	67.0	2.3
Other Asian ^d	2,001	1.0	72.7	1.2	999	0.9	74.1	1.7	1,002	1.0	71.3	1.7
Pacific Islander	947	0.4	59.8	2.0	547	0.5	58.5	2.7	400	0.4	61.6	3.0
Hawaiian	656	0.3	60.0	2.4	387	0.4	58.4	3.2	269	0.3	62.3	3.7
Other Pacific Islander ^e	291	0.1	59.4	3.5	160	0.1	58.7	4.8	131	0.1	59.6	5.2
Hispanic ^f	20,284	9.6	65.1	0.4	10,996	10.2	64.9	0.6	9,288	9.0	65.4	0.6

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2004-2010 with follow-up through 2011.

^b Percent of cases do not sum to 100 because All Races includes unknown race and other race which are not shown separately.

^c Survival estimates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^d Other Asians include Laotian, Hmong, Kampuchean, Thai, and Asians, Not Otherwise Specified (NOS).

^e Other Pacific Islanders include Micronesian, Chamorro, Guamanian, Polynesian, Tahitian, Samoan, Tongan, Melanesian, Fiji Islander, New Guinean, and Pacific Islander, NOS.

^f Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic not shown due to fewer than 25 cases during the time period.

Table 33.6

Cancer of the Liver and Intrahepatic Bile Duct

5-Year Cause-Specific Survival^a and Case Distribution by Race/Ethnicity
SEER 18 Registries, 2004-2010

Race/Ethnicity	Both Sexes				Males				Females			
	Case Distribution		Survival		Case Distribution		Survival		Case Distribution		Survival	
	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE
All Races	37,287	100.0	20.2	0.3	27,043	100.0	20.3	0.3	10,244	100.0	20.0	0.5
White	25,769	69.1	19.7	0.3	18,804	69.5	19.9	0.4	6,965	68.0	19.0	0.6
White Non-Hispanic	19,108	51.2	19.4	0.4	13,965	51.6	19.8	0.4	5,143	50.2	18.4	0.7
White Hispanic	6,661	17.9	20.4	0.6	4,839	17.9	20.3	0.8	1,822	17.8	20.8	1.2
Black	4,648	12.5	15.2	0.7	3,434	12.7	13.7	0.9	1,214	11.9	19.0	1.5
American Indian/Alaska Native ^c	360	1.0	17.4	2.5	259	1.0	20.5	3.0	101	1.0	8.2	4.3
Asian	5,927	15.9	26.3	0.7	4,103	15.2	26.9	0.9	1,824	17.8	24.8	1.3
Asian Indian, Pakistani	204	0.5	22.3	5.3	142	0.5	27.8	5.1	62	0.6	15.9	7.8
Chinese	1,500	4.0	30.0	1.5	1,067	3.9	30.9	1.8	433	4.2	27.3	2.8
Filipino	918	2.5	22.7	1.7	641	2.4	21.4	2.0	277	2.7	26.3	3.3
Japanese	542	1.5	23.5	2.4	263	1.0	28.0	3.5	279	2.7	20.4	3.1
Korean	698	1.9	27.9	2.0	467	1.7	27.3	2.4	231	2.3	29.0	3.4
Vietnamese	1,250	3.4	28.1	1.6	959	3.5	28.2	1.8	291	2.8	27.8	3.3
Other Asian ^d	815	2.2	21.1	1.9	564	2.1	22.2	2.2	251	2.5	18.4	3.5
Pacific Islander	321	0.9	16.2	2.9	250	0.9	15.7	3.3	71	0.7	17.6	6.1
Hawaiian	167	0.4	11.9	3.5	123	0.5	12.8	4.2	44	0.4	10.7	5.8
Other Pacific Islander ^e	154	0.4	23.3	4.7	127	0.5	20.0	5.0	27	0.3	41.6	10.5
Hispanic ^f	6,851	18.4	20.6	0.6	4,971	18.4	20.2	0.8	1,880	18.4	21.5	1.2

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2004-2010 with follow-up through 2011.

^b Percent of cases do not sum to 100 because All Races includes unknown race and other race which are not shown separately.

^c Survival estimates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^d Other Asians include Laotian, Hmong, Kampuchean, Thai, and Asians, Not Otherwise Specified (NOS).

^e Other Pacific Islanders include Micronesian, Chamorro, Guamanian, Polynesian, Tahitian, Samoan, Tongan, Melanesian, Fiji Islander, New Guinean, and Pacific Islander, NOS.

^f Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic not shown due to fewer than 25 cases during the time period.

Table 33.7

Cancer of the Lung and Bronchus

5-Year Cause-Specific Survival^a and Case Distribution by Race/Ethnicity
SEER 18 Registries, 2004-2010

Race/Ethnicity	Both Sexes				Males				Females			
	Case Distribution		Survival		Case Distribution		Survival		Case Distribution		Survival	
	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE
All Races	266,050	100.0	19.0	0.1	141,577	100.0	16.5	0.1	124,473	100.0	21.8	0.1
White	219,621	82.5	19.3	0.1	115,317	81.5	16.8	0.1	104,304	83.8	22.0	0.2
White Non-Hispanic	205,606	77.3	19.3	0.1	107,649	76.0	16.9	0.1	97,957	78.7	22.0	0.2
White Hispanic	14,015	5.3	18.6	0.4	7,668	5.4	14.8	0.5	6,347	5.1	23.2	0.7
Black	29,675	11.2	16.1	0.3	16,649	11.8	13.7	0.3	13,026	10.5	19.0	0.4
American Indian/Alaska Native ^c	971	0.4	12.8	1.4	500	0.4	12.3	1.9	471	0.4	13.5	2.0
Asian	13,554	5.1	20.8	0.4	7,841	5.5	18.3	0.6	5,713	4.6	24.1	0.7
Asian Indian, Pakistani	573	0.2	26.4	2.3	387	0.3	23.3	2.7	186	0.1	32.8	4.5
Chinese	3,422	1.3	20.8	0.9	1,967	1.4	19.2	1.1	1,455	1.2	23.1	1.5
Filipino	3,483	1.3	21.8	0.9	2,066	1.5	18.3	1.1	1,417	1.1	27.0	1.5
Japanese	2,109	0.8	18.4	1.0	1,075	0.8	16.7	1.4	1,034	0.8	20.1	1.6
Korean	1,060	0.4	18.2	1.5	578	0.4	15.4	1.9	482	0.4	21.5	2.3
Vietnamese	1,553	0.6	21.4	1.3	1,031	0.7	18.7	1.6	522	0.4	26.6	2.4
Other Asian ^d	1,354	0.5	21.1	1.4	737	0.5	18.2	1.8	617	0.5	24.6	2.3
Pacific Islander	1,246	0.5	17.6	1.4	733	0.5	14.9	1.7	513	0.4	21.2	2.3
Hawaiian	832	0.3	17.8	1.7	466	0.3	15.6	2.2	366	0.3	20.6	2.5
Other Pacific Islander ^e	414	0.2	16.8	2.5	267	0.2	13.8	2.8	147	0.1	22.3	4.6
Hispanic ^f	14,528	5.5	18.6	0.4	7,946	5.6	14.9	0.5	6,582	5.3	23.1	0.6

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2004-2010 with follow-up through 2011.

^b Percent of cases do not sum to 100 because All Races includes unknown race and other race which are not shown separately.

^c Survival estimates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^d Other Asians include Laotian, Hmong, Kampuchean, Thai, and Asians, Not Otherwise Specified (NOS).

^e Other Pacific Islanders include Micronesian, Chamorro, Guamanian, Polynesian, Tahitian, Samoan, Tongan, Melanesian, Fiji Islander, New Guinean, and Pacific Islander, NOS.

^f Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic not shown due to fewer than 25 cases during the time period.

Table 33.8

Cancer of the Prostate

5-Year Cause-Specific Survival^a and Case Distribution by Race/Ethnicity
SEER 18 Registries, 2004-2010

Race/Ethnicity	All Ages				Ages <65				Ages 65+			
	Case Distribution		Survival		Case Distribution		Survival		Case Distribution		Survival	
	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE
All Races	362,979	100.0	93.8	0.0	157,422	100.0	96.7	0.1	205,557	100.0	91.6	0.1
White	281,715	77.6	94.0	0.1	119,451	75.9	97.0	0.1	162,264	78.9	91.7	0.1
White Non-Hispanic	251,842	69.4	94.1	0.1	106,922	67.9	97.2	0.1	144,920	70.5	91.8	0.1
White Hispanic	29,873	8.2	93.0	0.2	12,529	8.0	95.6	0.2	17,344	8.4	91.1	0.3
Black	51,820	14.3	92.4	0.1	27,736	17.6	95.1	0.2	24,084	11.7	89.1	0.2
American Indian/Alaska Native ^c	922	0.3	89.5	1.2	389	0.2	92.7	1.6	533	0.3	87.0	1.8
Asian	15,297	4.2	94.5	0.2	5,059	3.2	96.9	0.3	10,238	5.0	93.3	0.3
Asian Indian, Pakistani	1,447	0.4	95.4	0.7	620	0.4	97.3	0.8	827	0.4	94.0	1.0
Chinese	3,228	0.9	94.8	0.5	921	0.6	99.0	0.4	2,307	1.1	93.2	0.6
Filipino	4,353	1.2	93.1	0.5	1,534	1.0	95.1	0.7	2,819	1.4	92.0	0.6
Japanese	2,574	0.7	94.9	0.5	680	0.4	97.4	0.7	1,894	0.9	94.0	0.6
Korean	760	0.2	94.7	1.0	175	0.1	94.8	2.1	585	0.3	94.6	1.1
Vietnamese	849	0.2	95.0	0.9	284	0.2	96.0	1.3	565	0.3	94.5	1.1
Other Asian ^d	2,086	0.6	95.6	0.5	845	0.5	98.0	0.6	1,241	0.6	94.0	0.8
Pacific Islander	1,175	0.3	90.1	1.0	434	0.3	93.8	1.4	741	0.4	88.1	1.4
Hawaiian	691	0.2	90.1	1.3	260	0.2	93.7	1.7	431	0.2	88.0	1.8
Other Pacific Islander ^e	484	0.1	90.2	1.6	174	0.1	93.9	2.2	310	0.2	88.2	2.2
Hispanic ^f	31,662	8.7	93.0	0.2	13,177	8.4	95.5	0.2	18,485	9.0	91.2	0.3

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2004-2010 with follow-up through 2011.

^b Percent of cases do not sum to 100 because All Races includes unknown race and other race which are not shown separately.

^c Survival estimates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^d Other Asians include Laotian, Hmong, Kampuchean, Thai, and Asians, Not Otherwise Specified (NOS).

^e Other Pacific Islanders include Micronesian, Chamorroan, Guamanian, Polynesian, Tahitian, Samoan, Tongan, Melanesian, Fiji Islander, New Guinean, and Pacific Islander, NOS.

^f Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic not shown due to fewer than 25 cases during the time period.