


Table 2.18
All Cancer Sites (Invasive)

Age-Adjusted Cancer Death^a Rates by State, All Races, 2008-2012

Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	171.22	0.10		
High Five States				
Kentucky	204.37 ^b	0.94	01(01,01)	19.36 ^c
Mississippi	199.87 ^b	1.14	02(02,04)	16.74 ^c
West Virginia	197.63 ^b	1.31	03(02,05)	15.43 ^c
Louisiana	196.85 ^b	0.93	04(02,05)	14.97 ^c
Arkansas	194.40 ^b	1.09	05(03,07)	13.54 ^c
Low Five States				
New Mexico	151.03 ^b	1.18	47(46,49)	-11.79 ^c
Arizona	150.85 ^b	0.66	48(47,48)	-11.90 ^c
Colorado	147.79 ^b	0.81	49(48,49)	-13.68 ^c
Hawaii	139.29 ^b	1.34	50(50,50)	-18.65 ^c
Utah	127.63 ^b	1.11	51(51,51)	-25.46 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	191.17 ^b	0.85	07(05,09)	11.66 ^c	Montana	160.42 ^b	1.66	41(36,45)	-6.30 ^c
Alaska	178.21	2.89	19(11,32)	4.08	Nebraska	165.96 ^b	1.29	35(32,40)	-3.07
Arizona	150.85 ^b	0.66	48(47,48)	-11.90 ^c	Nevada	172.38	1.17	26(20,33)	0.68
Arkansas	194.40 ^b	1.09	05(03,07)	13.54 ^c	New Hampshire	172.75	1.54	23(19,34)	0.90
California	155.15 ^b	0.30	46(44,46)	-9.38 ^c	New Jersey	167.60 ^b	0.59	34(32,36)	-2.11
Colorado	147.79 ^b	0.81	49(48,49)	-13.68 ^c	New Mexico	151.03 ^b	1.18	47(46,49)	-11.79 ^c
Connecticut	159.92 ^b	0.88	43(40,45)	-6.60 ^c	New York	162.28 ^b	0.39	40(38,43)	-5.22 ^c
Delaware	181.95 ^b	1.89	15(10,20)	6.27 ^c	North Carolina	176.18 ^b	0.60	20(19,22)	2.90
D.C.	186.97 ^b	2.57	10(05,17)	9.20 ^c	North Dakota	158.57 ^b	2.01	45(37,46)	-7.38 ^c
Florida	163.18 ^b	0.36	39(37,41)	-4.69	Ohio	186.92 ^b	0.53	11(09,12)	9.17 ^c
Georgia	173.17	0.64	22(20,30)	1.14	Oklahoma	190.64 ^b	0.97	08(05,10)	11.34 ^c
Hawaii	139.29 ^b	1.34	50(50,50)	-18.65 ^c	Oregon	172.43	0.90	25(20,32)	0.71
Idaho	158.72 ^b	1.43	44(39,46)	-7.30 ^c	Pennsylvania	179.66 ^b	0.48	17(15,19)	4.93
Illinois	178.86 ^b	0.52	18(15,20)	4.46	Rhode Island	172.05	1.68	28(19,34)	0.49
Indiana	188.26 ^b	0.74	09(08,11)	9.95 ^c	South Carolina	183.28 ^b	0.86	14(11,16)	7.05 ^c
Iowa	171.32	0.97	30(21,33)	0.06	South Dakota	165.35	1.88	36(30,43)	-3.43
Kansas	170.48	1.06	31(22,34)	-0.43	Tennessee	193.18 ^b	0.75	06(05,08)	12.83 ^c
Kentucky	204.37 ^b	0.94	01(01,01)	19.36 ^c	Texas	164.57 ^b	0.39	37(35,39)	-3.88
Louisiana	196.85 ^b	0.93	04(02,05)	14.97 ^c	Utah	127.63 ^b	1.11	51(51,51)	-25.46 ^c
Maine	183.33 ^b	1.48	13(10,18)	7.08 ^c	Vermont	173.72	2.18	21(17,34)	1.46
Maryland	171.75	0.77	29(21,33)	0.31	Virginia	172.55	0.66	24(20,31)	0.78
Massachusetts	170.30	0.68	32(26,34)	-0.53	Washington	169.64	0.71	33(27,35)	-0.92
Michigan	179.95 ^b	0.57	16(14,19)	5.10 ^c	West Virginia	197.63 ^b	1.31	03(02,05)	15.43 ^c
Minnesota	164.03 ^b	0.76	38(35,41)	-4.20	Wisconsin	172.32	0.74	27(21,32)	0.64
Mississippi	199.87 ^b	1.14	02(02,04)	16.74 ^c	Wyoming	160.36 ^b	2.38	42(35,46)	-6.34 ^c
Missouri	184.13 ^b	0.74	12(11,15)	7.54 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 5% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 2.19
All Cancer Sites (Invasive)

Age-Adjusted Cancer Death^a Rates by State, All Races, 2008-2012

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	207.86	0.17		
High Five States				
Mississippi	260.36 ^b	2.05	01(01,02)	25.26 ^c
Kentucky	253.64 ^b	1.62	02(01,03)	22.03 ^c
Louisiana	247.35 ^b	1.62	03(02,07)	19.00 ^c
Alabama	246.84 ^b	1.52	04(03,07)	18.76 ^c
Arkansas	246.48 ^b	1.88	05(02,07)	18.58 ^c
Low Five States				
New Mexico	181.02 ^b	1.96	47(44,48)	-12.91 ^c
Arizona	180.04 ^b	1.08	48(46,48)	-13.38 ^c
Colorado	173.84 ^b	1.35	49(48,50)	-16.37 ^c
Hawaii	171.73 ^b	2.22	50(49,50)	-17.38 ^c
Utah	153.02 ^b	1.84	51(51,51)	-26.38 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	246.84 ^b	1.52	04(03,07)	18.76 ^c	Montana	188.46 ^b	2.68	44(39,47)	-9.33 ^c
Alaska	211.37	4.81	23(12,38)	1.69	Nebraska	200.86	2.15	35(27,40)	-3.36
Arizona	180.04 ^b	1.08	48(46,48)	-13.38 ^c	Nevada	201.92	1.91	32(26,39)	-2.86
Arkansas	246.48 ^b	1.88	05(02,07)	18.58 ^c	New Hampshire	208.30	2.59	26(20,35)	0.22
California	183.36 ^b	0.49	46(44,47)	-11.78 ^c	New Jersey	198.99 ^b	1.00	36(33,40)	-4.26
Colorado	173.84 ^b	1.35	49(48,50)	-16.37 ^c	New Mexico	181.02 ^b	1.96	47(44,48)	-12.91 ^c
Connecticut	192.16 ^b	1.49	42(39,45)	-7.55 ^c	New York	193.77 ^b	0.66	41(39,43)	-6.78 ^c
Delaware	218.13	3.18	18(12,25)	4.94	North Carolina	222.23 ^b	1.05	15(12,17)	6.91 ^c
D.C.	227.13 ^b	4.48	12(08,21)	9.27 ^c	North Dakota	198.06	3.39	37(26,44)	-4.72
Florida	197.14 ^b	0.59	38(35,40)	-5.15 ^c	Ohio	228.19 ^b	0.91	11(10,13)	9.78 ^c
Georgia	218.19 ^b	1.15	17(14,22)	4.97	Oklahoma	233.58 ^b	1.65	08(08,11)	12.37 ^c
Hawaii	171.73 ^b	2.22	50(49,50)	-17.38 ^c	Oregon	203.58	1.47	31(26,36)	-2.06
Idaho	189.34 ^b	2.34	43(39,46)	-8.91 ^c	Pennsylvania	218.51 ^b	0.81	16(15,20)	5.13 ^c
Illinois	215.90 ^b	0.89	20(16,22)	3.87	Rhode Island	216.12	2.91	19(13,27)	3.98
Indiana	232.43 ^b	1.27	10(08,11)	11.82 ^c	South Carolina	232.66 ^b	1.51	09(08,12)	11.93 ^c
Iowa	209.05	1.63	24(21,31)	0.58	South Dakota	196.91 ^b	3.08	40(28,45)	-5.27 ^c
Kansas	206.13	1.76	29(23,35)	-0.83	Tennessee	245.91 ^b	1.33	06(03,07)	18.31 ^c
Kentucky	253.64 ^b	1.62	02(01,03)	22.03 ^c	Texas	201.78 ^b	0.67	34(30,37)	-2.92
Louisiana	247.35 ^b	1.62	03(02,07)	19.00 ^c	Utah	153.02 ^b	1.84	51(51,51)	-26.38 ^c
Maine	223.77 ^b	2.49	13(10,19)	7.65 ^c	Vermont	206.21	3.63	28(19,40)	-0.79
Maryland	207.37	1.31	27(22,31)	-0.23	Virginia	211.54	1.14	22(20,27)	1.77
Massachusetts	205.54	1.16	30(24,33)	-1.12	Washington	201.83 ^b	1.17	33(29,38)	-2.90
Michigan	215.86 ^b	0.96	21(16,22)	3.85	West Virginia	242.52 ^b	2.21	07(03,08)	16.68 ^c
Minnesota	196.96 ^b	1.27	39(34,42)	-5.24 ^c	Wisconsin	208.94	1.24	25(22,30)	0.52
Mississippi	260.36 ^b	2.05	01(01,02)	25.26 ^c	Wyoming	187.01 ^b	3.85	45(37,48)	-10.03 ^c
Missouri	222.74 ^b	1.25	14(12,17)	7.16 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 5% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 2.20
All Cancer Sites (Invasive)

Age-Adjusted Cancer Death^a Rates by State, All Races, 2008-2012

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	145.43	0.13		
High Five States				
Kentucky	169.98 ^b	1.14	01(01,02)	16.88 ^c
West Virginia	165.19 ^b	1.62	02(01,04)	13.59 ^c
D.C.	161.62 ^b	3.15	03(01,17)	11.13 ^c
Louisiana	161.00 ^b	1.12	04(02,09)	10.71 ^c
Oklahoma	159.62 ^b	1.20	05(03,12)	9.76 ^c
Low Five States				
Colorado	129.42 ^b	1.01	47(45,49)	-11.01 ^c
New Mexico	128.08 ^b	1.47	48(46,49)	-11.93 ^c
Arizona	128.07 ^b	0.83	49(46,49)	-11.93 ^c
Hawaii	114.84 ^b	1.66	50(50,51)	-21.03 ^c
Utah	108.61 ^b	1.39	51(50,51)	-25.32 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	152.43 ^b	1.02	17(12,21)	4.81	Montana	138.64	2.11	40(31,45)	-4.67
Alaska	151.50	3.59	18(03,37)	4.18	Nebraska	140.80	1.61	38(30,44)	-3.18
Arizona	128.07 ^b	0.83	49(46,49)	-11.93 ^c	Nevada	149.01	1.49	21(16,32)	2.46
Arkansas	156.57 ^b	1.32	11(04,17)	7.66 ^c	New Hampshire	147.58	1.91	24(16,36)	1.48
California	135.26 ^b	0.37	44(42,46)	-6.99 ^c	New Jersey	147.43	0.73	25(19,31)	1.38
Colorado	129.42 ^b	1.01	47(45,49)	-11.01 ^c	New Mexico	128.08 ^b	1.47	48(46,49)	-11.93 ^c
Connecticut	138.45 ^b	1.10	41(36,44)	-4.80	New York	141.74 ^b	0.48	36(33,40)	-2.54
Delaware	156.26 ^b	2.35	12(03,20)	7.45 ^c	North Carolina	144.95	0.72	31(25,36)	-0.33
D.C.	161.62 ^b	3.15	03(01,17)	11.13 ^c	North Dakota	130.16 ^b	2.49	46(42,49)	-10.50 ^c
Florida	136.94 ^b	0.45	43(39,44)	-5.84 ^c	Ohio	158.47 ^b	0.66	06(03,12)	8.97 ^c
Georgia	143.46	0.76	33(28,38)	-1.35	Oklahoma	159.62 ^b	1.20	05(03,12)	9.76 ^c
Hawaii	114.84 ^b	1.66	50(50,51)	-21.03 ^c	Oregon	149.77 ^b	1.14	20(16,27)	2.99
Idaho	134.88 ^b	1.80	45(39,47)	-7.25 ^c	Pennsylvania	153.34 ^b	0.60	16(12,19)	5.44 ^c
Illinois	154.37 ^b	0.64	15(10,18)	6.15 ^c	Rhode Island	143.30	2.04	34(20,41)	-1.46
Indiana	157.84 ^b	0.91	08(03,14)	8.53 ^c	South Carolina	148.79	1.03	22(17,30)	2.31
Iowa	145.30	1.22	30(21,36)	-0.08	South Dakota	142.77	2.38	35(20,44)	-1.83
Kansas	144.68	1.31	32(21,38)	-0.52	Tennessee	156.69 ^b	0.90	10(05,15)	7.74 ^c
Kentucky	169.98 ^b	1.14	01(01,02)	16.88 ^c	Texas	137.70 ^b	0.48	42(38,44)	-5.31 ^c
Louisiana	161.00 ^b	1.12	04(02,09)	10.71 ^c	Utah	108.61 ^b	1.39	51(50,51)	-25.32 ^c
Maine	154.60 ^b	1.83	14(04,21)	6.31 ^c	Vermont	149.89	2.73	19(08,36)	3.07
Maryland	147.97	0.94	23(18,31)	1.75	Virginia	145.79	0.80	29(22,34)	0.25
Massachusetts	147.10	0.84	26(19,33)	1.15	Washington	146.40	0.89	28(20,34)	0.67
Michigan	155.02 ^b	0.71	13(09,18)	6.59 ^c	West Virginia	165.19 ^b	1.62	02(01,04)	13.59 ^c
Minnesota	141.13 ^b	0.96	37(32,41)	-2.96	Wisconsin	146.83	0.92	27(19,33)	0.97
Mississippi	158.45 ^b	1.35	07(03,15)	8.96 ^c	Wyoming	140.27	3.05	39(20,46)	-3.55
Missouri	156.81 ^b	0.92	09(05,16)	7.82 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 5% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 3.18
Cancer of the Brain and Other Nervous System (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	4.31	0.02		
High Five States				
South Dakota	5.52	0.35	01(01,25)	28.29 ^c
Washington	5.16 ^b	0.12	02(01,15)	19.98 ^c
Nebraska	5.10	0.23	03(01,26)	18.48 ^c
Vermont	5.10	0.38	04(01,43)	18.41 ^c
Kansas	5.08 ^b	0.18	05(01,23)	17.92 ^c
Low Five States				
New Mexico	3.79	0.19	47(29,50)	-12.05
New York	3.75 ^b	0.06	48(43,50)	-12.82
New Jersey	3.74 ^b	0.09	49(41,50)	-13.20
D.C.	3.60	0.36	50(12,51)	-16.33 ^c
Hawaii	2.42 ^b	0.18	51(50,51)	-43.88 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	4.92 ^b	0.14	11(01,24)	14.27	Montana	4.56	0.28	22(01,48)	6.00
Alaska	4.73	0.43	18(01,50)	9.83	Nebraska	5.10	0.23	03(01,26)	18.48 ^c
Arizona	4.17	0.11	36(22,47)	-3.05	Nevada	4.14	0.18	39(16,50)	-3.95
Arkansas	4.92	0.18	10(01,27)	14.32	New Hampshire	4.77	0.26	15(01,43)	10.82
California	4.28	0.05	30(24,39)	-0.53	New Jersey	3.74 ^b	0.09	49(41,50)	-13.20
Colorado	4.42	0.14	26(12,42)	2.78	New Mexico	3.79	0.19	47(29,50)	-12.05
Connecticut	4.24	0.15	33(17,47)	-1.48	New York	3.75 ^b	0.06	48(43,50)	-12.82
Delaware	4.24	0.30	32(04,50)	-1.47	North Carolina	4.31	0.09	28(19,42)	0.22
D.C.	3.60	0.36	50(12,51)	-16.33 ^c	North Dakota	3.87	0.33	46(12,50)	-10.12
Florida	4.07 ^b	0.06	41(31,47)	-5.41	Ohio	4.48	0.08	25(15,34)	4.15
Georgia	4.09	0.10	40(26,48)	-5.05	Oklahoma	4.72	0.15	19(03,32)	9.68
Hawaii	2.42 ^b	0.18	51(50,51)	-43.88 ^c	Oregon	4.98 ^b	0.16	08(01,24)	15.70 ^c
Idaho	4.79	0.25	14(01,42)	11.37	Pennsylvania	4.17	0.08	37(25,45)	-3.10
Illinois	4.06	0.08	42(30,47)	-5.59	Rhode Island	4.16	0.27	38(07,50)	-3.39
Indiana	4.59	0.12	21(09,33)	6.65	South Carolina	4.36	0.13	27(14,45)	1.19
Iowa	5.07 ^b	0.17	06(01,22)	17.88 ^c	South Dakota	5.52	0.35	01(01,25)	28.29 ^c
Kansas	5.08 ^b	0.18	05(01,23)	17.92 ^c	Tennessee	4.76 ^b	0.12	16(04,27)	10.49
Kentucky	4.74	0.14	17(04,30)	10.08	Texas	4.03 ^b	0.06	43(32,47)	-6.44
Louisiana	4.24	0.14	35(18,47)	-1.60	Utah	4.31	0.20	29(11,48)	0.15
Maine	5.04	0.25	07(01,31)	17.09 ^c	Vermont	5.10	0.38	04(01,43)	18.41 ^c
Maryland	3.94	0.12	45(30,50)	-8.39	Virginia	3.97	0.10	44(30,50)	-7.76
Massachusetts	4.25	0.11	31(20,45)	-1.18	Washington	5.16 ^b	0.12	02(01,15)	19.98 ^c
Michigan	4.85 ^b	0.10	12(04,23)	12.55	West Virginia	4.24	0.19	34(12,49)	-1.59
Minnesota	4.52	0.13	23(10,37)	4.88	Wisconsin	4.97 ^b	0.13	09(01,21)	15.50 ^c
Mississippi	4.63	0.17	20(04,39)	7.58	Wyoming	4.83	0.41	13(01,49)	12.15
Missouri	4.51	0.12	24(11,37)	4.66					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.


- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 3.19
Cancer of the Brain and Other Nervous System (Invasive)
Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	5.28	0.03		
High Five States				
Maine	6.60	0.42	01(01,28)	25.07 ^c
Idaho	6.60	0.42	02(01,28)	25.02 ^c
Kansas	6.53 ^b	0.31	03(01,20)	23.84 ^c
Iowa	6.39 ^b	0.29	04(01,23)	21.18 ^c
Washington	6.38 ^b	0.20	05(01,16)	20.88 ^c
Low Five States				
New York	4.59 ^b	0.10	47(40,50)	-13.08
New Mexico	4.53	0.30	48(23,50)	-14.22
D.C.	4.39	0.61	49(04,51)	-16.74 ^c
North Dakota	4.15	0.49	50(16,51)	-21.41 ^c
Hawaii	3.03 ^b	0.29	51(49,51)	-42.53 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	6.01	0.23	13(02,28)	13.91	Montana	5.32	0.45	29(02,50)	0.89
Alaska	6.12	0.76	10(01,50)	15.95 ^c	Nebraska	6.07	0.37	12(01,39)	15.00 ^c
Arizona	4.98	0.18	41(21,49)	-5.54	Nevada	5.35	0.29	28(07,48)	1.47
Arkansas	6.00	0.29	14(01,35)	13.68	New Hampshire	6.19	0.43	09(01,42)	17.34 ^c
California	5.23	0.08	33(21,41)	-0.82	New Jersey	4.75	0.15	46(30,50)	-9.93
Colorado	5.46	0.23	23(08,45)	3.57	New Mexico	4.53	0.30	48(23,50)	-14.22
Connecticut	5.27	0.24	31(11,48)	-0.17	New York	4.59 ^b	0.10	47(40,50)	-13.08
Delaware	5.05	0.48	39(04,50)	-4.23	North Carolina	5.48	0.16	22(11,40)	3.85
D.C.	4.39	0.61	49(04,51)	-16.74 ^c	North Dakota	4.15	0.49	50(16,51)	-21.41 ^c
Florida	5.09	0.10	37(24,45)	-3.45	Ohio	5.42	0.14	25(14,39)	2.78
Georgia	5.11	0.16	36(18,47)	-3.09	Oklahoma	5.59	0.25	19(05,42)	6.03
Hawaii	3.03 ^b	0.29	51(49,51)	-42.53 ^c	Oregon	6.21 ^b	0.25	07(01,24)	17.75 ^c
Idaho	6.60	0.42	02(01,28)	25.02 ^c	Pennsylvania	5.07	0.12	38(23,47)	-3.97
Illinois	5.03	0.13	40(23,47)	-4.68	Rhode Island	5.25	0.45	32(03,50)	-0.44
Indiana	5.60	0.19	18(07,37)	6.19	South Carolina	5.56	0.23	20(07,42)	5.34
Iowa	6.39 ^b	0.29	04(01,23)	21.18 ^c	South Dakota	6.20	0.54	08(01,46)	17.53 ^c
Kansas	6.53 ^b	0.31	03(01,20)	23.84 ^c	Tennessee	5.66	0.19	17(07,36)	7.26
Kentucky	5.45	0.23	24(08,45)	3.24	Texas	4.84 ^b	0.10	44(32,49)	-8.19
Louisiana	5.14	0.23	35(14,49)	-2.61	Utah	5.28	0.32	30(06,49)	0.15
Maine	6.60	0.42	01(01,28)	25.07 ^c	Vermont	6.32	0.61	06(01,47)	19.81 ^c
Maryland	4.92	0.19	43(21,49)	-6.64	Virginia	4.78	0.16	45(29,50)	-9.40
Massachusetts	5.14	0.18	34(16,47)	-2.53	Washington	6.38 ^b	0.20	05(01,16)	20.88 ^c
Michigan	5.92 ^b	0.16	15(04,25)	12.25	West Virginia	4.94	0.31	42(13,50)	-6.40
Minnesota	5.51	0.21	21(08,42)	4.50	Wisconsin	6.09 ^b	0.21	11(02,24)	15.36 ^c
Mississippi	5.39	0.29	26(07,48)	2.22	Wyoming	5.66	0.64	16(01,50)	7.36
Missouri	5.37	0.19	27(11,44)	1.73					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 3.20
Cancer of the Brain and Other Nervous System (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	3.48	0.02		
High Five States				
South Dakota	4.89	0.47	01(01,26)	40.48 ^c
Nebraska	4.20	0.29	02(01,35)	20.49 ^c
Kentucky	4.13 ^b	0.18	03(01,25)	18.52 ^c
Washington	4.09 ^b	0.15	04(01,23)	17.31 ^c
Mississippi	4.05	0.22	05(01,35)	16.15 ^c
Low Five States				
New York	3.09 ^b	0.07	47(35,50)	-11.19
D.C.	3.03	0.44	48(02,51)	-12.95
Nevada	3.01	0.21	49(22,50)	-13.56
New Jersey	2.91 ^b	0.11	50(41,50)	-16.49 ^c
Hawaii	1.87 ^b	0.22	51(50,51)	-46.17 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	4.00	0.17	09(01,28)	14.93	Montana	3.87	0.36	16(01,50)	11.05
Alaska	3.58	0.50	26(01,50)	2.93	Nebraska	4.20	0.29	02(01,35)	20.49 ^c
Arizona	3.47	0.14	31(13,47)	-0.45	Nevada	3.01	0.21	49(22,50)	-13.56
Arkansas	4.02	0.22	07(01,35)	15.46 ^c	New Hampshire	3.58	0.30	27(02,50)	2.84
California	3.47	0.06	32(21,40)	-0.48	New Jersey	2.91 ^b	0.11	50(41,50)	-16.49 ^c
Colorado	3.54	0.17	28(06,47)	1.78	New Mexico	3.14	0.24	45(12,50)	-9.94
Connecticut	3.37	0.18	36(12,49)	-3.16	New York	3.09 ^b	0.07	47(35,50)	-11.19
Delaware	3.63	0.37	23(01,50)	4.15	North Carolina	3.39	0.11	35(18,47)	-2.63
D.C.	3.03	0.44	48(02,51)	-12.95	North Dakota	3.60	0.44	25(01,50)	3.31
Florida	3.18 ^b	0.07	43(32,49)	-8.71	Ohio	3.66	0.10	21(10,36)	5.08
Georgia	3.27	0.12	41(22,49)	-5.98	Oklahoma	3.95	0.19	11(01,35)	13.35
Hawaii	1.87 ^b	0.22	51(50,51)	-46.17 ^c	Oregon	3.87	0.19	15(02,38)	11.14
Idaho	3.12	0.28	46(09,50)	-10.47	Pennsylvania	3.42	0.09	34(19,45)	-1.83
Illinois	3.28	0.10	40(25,48)	-5.85	Rhode Island	3.23	0.32	42(05,50)	-7.34
Indiana	3.74	0.14	19(04,38)	7.40	South Carolina	3.34	0.16	37(16,49)	-4.00
Iowa	3.95	0.21	12(01,36)	13.32	South Dakota	4.89	0.47	01(01,26)	40.48 ^c
Kansas	3.84	0.22	17(02,43)	10.27	Tennessee	3.99	0.15	10(02,26)	14.57
Kentucky	4.13 ^b	0.18	03(01,25)	18.52 ^c	Texas	3.34	0.07	38(24,46)	-4.09
Louisiana	3.49	0.17	30(10,47)	0.25	Utah	3.45	0.24	33(04,50)	-0.85
Maine	3.69	0.30	20(01,49)	5.84	Vermont	3.91	0.46	14(01,50)	12.30
Maryland	3.15	0.14	44(24,50)	-9.64	Virginia	3.29	0.12	39(22,49)	-5.55
Massachusetts	3.51	0.13	29(12,46)	0.86	Washington	4.09 ^b	0.15	04(01,23)	17.31 ^c
Michigan	3.93 ^b	0.12	13(03,26)	12.91	West Virginia	3.62	0.25	24(02,49)	3.88
Minnesota	3.64	0.16	22(05,43)	4.44	Wisconsin	4.03	0.16	06(01,28)	15.59 ^c
Mississippi	4.05	0.22	05(01,35)	16.15 ^c	Wyoming	4.02	0.52	08(01,50)	15.32 ^c
Missouri	3.75	0.15	18(04,38)	7.72					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 4.23
Cancer of the Female Breast (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	21.90	0.05		
High Five States				
D.C.	29.01 ^b	1.34	01(01,02)	32.51 ^c
Louisiana	25.01 ^b	0.44	02(01,07)	14.24 ^c
Mississippi	24.45 ^b	0.54	03(02,15)	11.68 ^c
New Jersey	23.92 ^b	0.30	04(02,13)	9.23
Ohio	23.76 ^b	0.26	05(03,14)	8.50
Low Five States				
Colorado	19.67 ^b	0.39	47(32,50)	-10.15 ^c
Wyoming	19.47	1.12	48(11,50)	-11.10 ^c
Maine	19.40 ^b	0.66	49(26,50)	-11.40 ^c
Vermont	18.68	0.97	50(25,50)	-14.68 ^c
Hawaii	15.06 ^b	0.62	51(51,51)	-31.24 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	22.59	0.40	17(05,27)	3.17	Montana	20.30	0.82	39(15,50)	-7.29
Alaska	20.98	1.30	28(02,50)	-4.16	Nebraska	19.77	0.61	45(24,50)	-9.70
Arizona	19.68 ^b	0.33	46(32,50)	-10.14 ^c	Nevada	23.27	0.58	08(02,25)	6.26
Arkansas	22.28	0.50	21(05,34)	1.77	New Hampshire	20.36	0.71	37(18,50)	-7.03
California	21.24 ^b	0.15	26(23,35)	-3.02	New Jersey	23.92 ^b	0.30	04(02,13)	9.23
Colorado	19.67 ^b	0.39	47(32,50)	-10.15 ^c	New Mexico	20.39	0.59	36(21,50)	-6.86
Connecticut	20.31 ^b	0.42	38(24,49)	-7.26	New York	21.52	0.19	24(21,33)	-1.72
Delaware	22.10	0.89	23(02,47)	0.95	North Carolina	22.19	0.28	22(11,27)	1.36
D.C.	29.01 ^b	1.34	01(01,02)	32.51 ^c	North Dakota	19.83	0.98	43(13,50)	-9.44
Florida	20.95 ^b	0.18	30(24,39)	-4.31	Ohio	23.76 ^b	0.26	05(03,14)	8.50
Georgia	22.87	0.30	14(05,23)	4.45	Oklahoma	23.21	0.46	09(02,24)	6.00
Hawaii	15.06 ^b	0.62	51(51,51)	-31.24 ^c	Oregon	20.87	0.43	31(21,46)	-4.67
Idaho	20.69	0.71	34(15,50)	-5.52	Pennsylvania	23.21 ^b	0.24	10(04,19)	5.99
Illinois	23.05 ^b	0.25	13(05,22)	5.25	Rhode Island	19.80	0.77	44(21,50)	-9.59
Indiana	22.74	0.35	16(05,25)	3.85	South Carolina	23.20	0.41	11(03,23)	5.97
Iowa	20.67	0.47	35(22,48)	-5.60	South Dakota	20.73	0.92	33(07,50)	-5.34
Kansas	21.30	0.51	25(15,44)	-2.74	Tennessee	22.57	0.34	19(06,26)	3.06
Kentucky	22.57	0.42	18(05,28)	3.06	Texas	21.04 ^b	0.18	27(24,38)	-3.89
Louisiana	25.01 ^b	0.44	02(01,07)	14.24 ^c	Utah	20.79	0.61	32(17,49)	-5.03
Maine	19.40 ^b	0.66	49(26,50)	-11.40 ^c	Vermont	18.68	0.97	50(25,50)	-14.68 ^c
Maryland	23.71 ^b	0.38	06(02,19)	8.29	Virginia	22.78	0.32	15(05,24)	4.05
Massachusetts	20.26 ^b	0.31	41(27,48)	-7.46	Washington	20.27 ^b	0.33	40(27,48)	-7.42
Michigan	23.10 ^b	0.28	12(04,22)	5.52	West Virginia	22.55	0.61	20(03,34)	2.98
Minnesota	20.05 ^b	0.36	42(28,49)	-8.45	Wisconsin	20.98	0.35	29(21,44)	-4.17
Mississippi	24.45 ^b	0.54	03(02,15)	11.68 ^c	Wyoming	19.47	1.12	48(11,50)	-11.10 ^c
Missouri	23.40 ^b	0.36	07(03,21)	6.87					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 10% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 5.14
Cancer of the Cervix Uteri (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	2.31	0.02		
High Five States				
Arkansas	3.39 ^b	0.21	01(01,10)	47.14 ^c
Mississippi	3.37 ^b	0.21	02(01,11)	46.10 ^c
West Virginia	3.29 ^b	0.26	03(01,17)	42.78 ^c
Oklahoma	3.16 ^b	0.18	04(01,14)	37.07 ^c
Alabama	3.03 ^b	0.15	05(01,17)	31.16 ^c
Low Five States				
Massachusetts	1.40 ^b	0.09	47(40,51)	-39.29 ^c
Rhode Island	1.34 ^b	0.21	48(30,51)	-41.89 ^c
Utah	1.29 ^b	0.15	49(38,51)	-44.02 ^c
Vermont	1.20 ^b	0.26	50(29,51)	-47.85 ^c
North Dakota	1.16 ^b	0.25	51(28,51)	-49.52 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	3.03 ^b	0.15	05(01,17)	31.16 ^c	Montana	1.51	0.24	45(21,51)	-34.56 ^c
Alaska	2.03	0.36	30(03,51)	-11.88	Nebraska	1.95	0.21	33(14,48)	-15.66 ^c
Arizona	1.98	0.11	32(21,42)	-14.11	Nevada	2.20	0.18	23(10,40)	-4.51
Arkansas	3.39 ^b	0.21	01(01,10)	47.14 ^c	New Hampshire	1.64	0.21	41(22,51)	-28.72 ^c
California	2.23	0.05	22(18,30)	-3.37	New Jersey	2.38	0.10	20(12,29)	3.15
Colorado	1.53 ^b	0.11	44(35,50)	-33.77 ^c	New Mexico	2.14	0.20	25(10,44)	-7.02
Connecticut	1.66 ^b	0.13	40(28,49)	-28.11 ^c	New York	2.29	0.06	21(16,30)	-0.75
Delaware	2.56	0.33	17(01,41)	10.80	North Carolina	2.05	0.09	29(21,39)	-11.32
D.C.	2.84	0.43	09(01,40)	23.01 ^c	North Dakota	1.16 ^b	0.25	51(28,51)	-49.52 ^c
Florida	2.71 ^b	0.07	13(06,18)	17.48 ^c	Ohio	2.57	0.09	15(08,22)	11.30
Georgia	2.71 ^b	0.10	12(05,20)	17.60 ^c	Oklahoma	3.16 ^b	0.18	04(01,14)	37.07 ^c
Hawaii	1.83	0.22	38(16,50)	-20.68 ^c	Oregon	1.88	0.13	37(21,45)	-18.42 ^c
Idaho	1.90	0.22	35(14,49)	-17.41 ^c	Pennsylvania	2.13	0.08	26(19,35)	-7.69
Illinois	2.53	0.08	18(09,23)	9.90	Rhode Island	1.34 ^b	0.21	48(30,51)	-41.89 ^c
Indiana	2.41	0.12	19(10,29)	4.38	South Carolina	2.73	0.15	11(03,22)	18.20 ^c
Iowa	1.94	0.16	34(19,45)	-15.91 ^c	South Dakota	2.06	0.31	28(05,50)	-10.77
Kansas	1.99	0.17	31(17,45)	-13.53	Tennessee	2.70	0.12	14(05,21)	17.01 ^c
Kentucky	2.95 ^b	0.16	07(02,18)	27.72 ^c	Texas	2.77 ^b	0.07	10(05,16)	20.06 ^c
Louisiana	3.00 ^b	0.16	06(01,17)	30.21 ^c	Utah	1.29 ^b	0.15	49(38,51)	-44.02 ^c
Maine	1.59 ^b	0.20	42(24,51)	-31.21 ^c	Vermont	1.20 ^b	0.26	50(29,51)	-47.85 ^c
Maryland	2.16	0.12	24(16,38)	-6.28	Virginia	1.90 ^b	0.09	36(23,42)	-17.54 ^c
Massachusetts	1.40 ^b	0.09	47(40,51)	-39.29 ^c	Washington	1.81 ^b	0.10	39(27,45)	-21.70 ^c
Michigan	2.08	0.09	27(19,38)	-9.62	West Virginia	3.29 ^b	0.26	03(01,17)	42.78 ^c
Minnesota	1.46 ^b	0.10	46(37,51)	-36.82 ^c	Wisconsin	1.53 ^b	0.10	43(35,50)	-33.61 ^c
Mississippi	3.37 ^b	0.21	02(01,11)	46.10 ^c	Wyoming	2.86	0.45	08(01,41)	23.92 ^c
Missouri	2.56	0.13	16(07,25)	10.84					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 6.25
Cancer of the Colon and Rectum (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	15.50	0.03		
High Five States				
Mississippi	19.69 ^b	0.36	01(01,03)	27.01 ^c
Arkansas	18.54 ^b	0.34	02(01,08)	19.65 ^c
Louisiana	18.46 ^b	0.28	03(01,07)	19.12 ^c
West Virginia	18.43 ^b	0.40	04(01,10)	18.89 ^c
Kentucky	18.10 ^b	0.28	05(02,10)	16.76 ^c
Low Five States				
Arizona	13.44 ^b	0.20	47(43,50)	-13.31 ^c
Hawaii	13.40 ^b	0.41	48(36,50)	-13.54 ^c
Colorado	13.37 ^b	0.24	49(42,50)	-13.76 ^c
Connecticut	12.68 ^b	0.25	50(47,51)	-18.16 ^c
Utah	11.62 ^b	0.34	51(50,51)	-25.01 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	17.19 ^b	0.26	09(05,18)	10.94 ^c	Montana	14.22	0.49	42(23,50)	-8.22
Alaska	15.84	0.86	21(03,48)	2.19	Nebraska	16.96 ^b	0.41	12(04,23)	9.41
Arizona	13.44 ^b	0.20	47(43,50)	-13.31 ^c	Nevada	17.15 ^b	0.37	10(04,21)	10.67 ^c
Arkansas	18.54 ^b	0.34	02(01,08)	19.65 ^c	New Hampshire	14.43	0.44	39(22,49)	-6.90
California	14.22 ^b	0.09	43(36,46)	-8.27	New Jersey	16.26 ^b	0.18	18(13,24)	4.92
Colorado	13.37 ^b	0.24	49(42,50)	-13.76 ^c	New Mexico	14.82	0.37	34(21,46)	-4.37
Connecticut	12.68 ^b	0.25	50(47,51)	-18.16 ^c	New York	15.08 ^b	0.12	31(24,37)	-2.71
Delaware	14.60	0.54	37(19,49)	-5.81	North Carolina	14.75 ^b	0.17	35(27,42)	-4.85
D.C.	17.47	0.78	08(01,29)	12.72 ^c	North Dakota	15.97	0.63	20(05,43)	3.07
Florida	14.40 ^b	0.11	40(33,45)	-7.07	Ohio	17.00 ^b	0.16	11(07,18)	9.70
Georgia	15.76	0.19	22(18,29)	1.66	Oklahoma	17.60 ^b	0.30	06(03,16)	13.58 ^c
Hawaii	13.40 ^b	0.41	48(36,50)	-13.54 ^c	Oregon	14.86	0.26	33(23,44)	-4.13
Idaho	13.75 ^b	0.42	46(30,50)	-11.31 ^c	Pennsylvania	16.73 ^b	0.15	14(09,20)	7.94
Illinois	16.68 ^b	0.16	17(09,21)	7.64	Rhode Island	15.08	0.49	30(16,47)	-2.70
Indiana	16.73 ^b	0.22	15(07,21)	7.92	South Carolina	16.18	0.26	19(11,27)	4.42
Iowa	16.71 ^b	0.30	16(06,23)	7.82	South Dakota	15.69	0.57	23(08,43)	1.26
Kansas	15.52	0.32	25(17,37)	0.16	Tennessee	17.56 ^b	0.23	07(04,13)	13.27 ^c
Kentucky	18.10 ^b	0.28	05(02,10)	16.76 ^c	Texas	15.38	0.12	26(21,32)	-0.77
Louisiana	18.46 ^b	0.28	03(01,07)	19.12 ^c	Utah	11.62 ^b	0.34	51(50,51)	-25.01 ^c
Maine	15.19	0.43	28(17,45)	-2.02	Vermont	14.65	0.63	36(16,49)	-5.46
Maryland	15.36	0.23	27(20,36)	-0.90	Virginia	15.12	0.20	29(22,38)	-2.44
Massachusetts	14.27 ^b	0.20	41(32,47)	-7.90	Washington	14.09 ^b	0.20	44(34,48)	-9.10
Michigan	15.57	0.17	24(19,32)	0.46	West Virginia	18.43 ^b	0.40	04(01,10)	18.89 ^c
Minnesota	13.94 ^b	0.22	45(36,49)	-10.03 ^c	Wisconsin	14.52 ^b	0.21	38(28,45)	-6.30
Mississippi	19.69 ^b	0.36	01(01,03)	27.01 ^c	Wyoming	14.98	0.72	32(09,49)	-3.33
Missouri	16.77 ^b	0.22	13(07,21)	8.20					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 10% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 6.26
Cancer of the Colon and Rectum (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	18.59	0.05		
High Five States				
Mississippi	23.88 ^b	0.62	01(01,05)	28.43 ^c
West Virginia	22.88 ^b	0.69	02(01,10)	23.06 ^c
Arkansas	22.72 ^b	0.57	03(01,09)	22.18 ^c
Louisiana	22.69 ^b	0.49	04(01,08)	22.04 ^c
Kentucky	21.77 ^b	0.48	05(02,13)	17.10 ^c
Low Five States				
Idaho	15.97 ^b	0.68	47(29,50)	-14.08 ^c
Arizona	15.88 ^b	0.32	48(39,50)	-14.61 ^c
Colorado	15.30 ^b	0.40	49(42,50)	-17.73 ^c
Connecticut	14.79 ^b	0.41	50(45,51)	-20.44 ^c
Utah	13.19 ^b	0.53	51(49,51)	-29.09 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	21.24 ^b	0.45	08(03,16)	14.26 ^c	Montana	16.12	0.78	46(25,50)	-13.32 ^c
Alaska	18.01	1.38	29(04,50)	-3.14	Nebraska	19.84	0.67	15(05,32)	6.73
Arizona	15.88 ^b	0.32	48(39,50)	-14.61 ^c	Nevada	20.96 ^b	0.62	09(03,22)	12.71 ^c
Arkansas	22.72 ^b	0.57	03(01,09)	22.18 ^c	New Hampshire	16.22	0.73	45(26,50)	-12.76 ^c
California	16.77 ^b	0.15	40(35,46)	-9.78	New Jersey	19.58	0.31	18(11,26)	5.29
Colorado	15.30 ^b	0.40	49(42,50)	-17.73 ^c	New Mexico	18.01	0.62	30(15,45)	-3.14
Connecticut	14.79 ^b	0.41	50(45,51)	-20.44 ^c	New York	17.88 ^b	0.20	33(25,37)	-3.85
Delaware	17.45	0.90	35(13,50)	-6.12	North Carolina	17.99	0.30	31(22,38)	-3.26
D.C.	18.72	1.27	23(03,49)	0.67	North Dakota	19.59	1.06	17(03,42)	5.37
Florida	17.19 ^b	0.18	38(31,43)	-7.53	Ohio	20.88 ^b	0.28	10(05,16)	12.29 ^c
Georgia	19.49	0.34	20(11,27)	4.82	Oklahoma	21.70 ^b	0.50	06(02,14)	16.70 ^c
Hawaii	16.77	0.69	41(22,50)	-9.80	Oregon	17.29	0.43	37(25,46)	-7.00
Idaho	15.97 ^b	0.68	47(29,50)	-14.08 ^c	Pennsylvania	20.16 ^b	0.25	14(09,21)	8.41
Illinois	20.28 ^b	0.27	13(08,20)	9.07	Rhode Island	17.63	0.83	34(13,49)	-5.18
Indiana	20.32 ^b	0.38	12(06,22)	9.27	South Carolina	19.57	0.44	19(10,29)	5.25
Iowa	19.78	0.50	16(07,29)	6.39	South Dakota	18.46	0.94	26(08,47)	-0.73
Kansas	18.68	0.53	24(12,39)	0.46	Tennessee	21.33 ^b	0.39	07(03,15)	14.74 ^c
Kentucky	21.77 ^b	0.48	05(02,13)	17.10 ^c	Texas	18.95	0.20	21(17,29)	1.89
Louisiana	22.69 ^b	0.49	04(01,08)	22.04 ^c	Utah	13.19 ^b	0.53	51(49,51)	-29.09 ^c
Maine	18.36	0.72	27(11,45)	-1.25	Vermont	16.24	1.03	44(19,51)	-12.68 ^c
Maryland	18.86	0.39	22(15,34)	1.42	Virginia	17.90	0.33	32(22,40)	-3.72
Massachusetts	17.15 ^b	0.33	39(27,46)	-7.77	Washington	16.37 ^b	0.33	43(35,49)	-11.98 ^c
Michigan	18.59	0.28	25(18,33)	0.00	West Virginia	22.88 ^b	0.69	02(01,10)	23.06 ^c
Minnesota	16.54 ^b	0.37	42(32,48)	-11.03 ^c	Wisconsin	17.37	0.36	36(25,45)	-6.56
Mississippi	23.88 ^b	0.62	01(01,05)	28.43 ^c	Wyoming	18.27	1.19	28(06,50)	-1.76
Missouri	20.51 ^b	0.38	11(05,21)	10.33 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 10% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 6.27
Cancer of the Colon and Rectum (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	13.07	0.04		
High Five States				
D.C.	16.62 ^b	1.00	01(01,19)	27.21 ^c
Mississippi	16.52 ^b	0.43	02(01,05)	26.38 ^c
Arkansas	15.28 ^b	0.41	03(01,14)	16.96 ^c
Kentucky	15.17 ^b	0.34	04(02,13)	16.09 ^c
Louisiana	15.14 ^b	0.34	05(02,13)	15.82 ^c
Low Five States				
Idaho	11.82	0.53	47(21,51)	-9.57
Arizona	11.33 ^b	0.25	48(41,50)	-13.29 ^c
Connecticut	11.05 ^b	0.30	49(42,51)	-15.45 ^c
Hawaii	10.68 ^b	0.50	50(39,51)	-18.29 ^c
Utah	10.22 ^b	0.43	51(46,51)	-21.79 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	14.13	0.31	11(05,24)	8.17	Montana	12.46	0.63	35(10,50)	-4.67
Alaska	13.74	1.09	19(01,50)	5.16	Nebraska	14.59	0.51	08(02,26)	11.64 ^c
Arizona	11.33 ^b	0.25	48(41,50)	-13.29 ^c	Nevada	13.83	0.46	17(04,35)	5.86
Arkansas	15.28 ^b	0.41	03(01,14)	16.96 ^c	New Hampshire	13.03	0.56	27(06,48)	-0.25
California	12.19 ^b	0.11	42(32,46)	-6.72	New Jersey	13.79	0.22	18(08,26)	5.56
Colorado	11.83 ^b	0.30	45(30,50)	-9.48	New Mexico	12.16	0.45	43(20,50)	-6.97
Connecticut	11.05 ^b	0.30	49(42,51)	-15.45 ^c	New York	13.01	0.14	28(20,35)	-0.45
Delaware	12.20	0.65	39(11,51)	-6.64	North Carolina	12.30 ^b	0.21	37(28,47)	-5.85
D.C.	16.62 ^b	1.00	01(01,19)	27.21 ^c	North Dakota	13.08	0.77	26(03,50)	0.07
Florida	12.06 ^b	0.13	44(33,47)	-7.69	Ohio	14.06 ^b	0.19	13(07,23)	7.56
Georgia	13.10	0.23	25(17,36)	0.24	Oklahoma	14.43 ^b	0.36	09(03,23)	10.41 ^c
Hawaii	10.68 ^b	0.50	50(39,51)	-18.29 ^c	Oregon	12.83	0.33	31(17,45)	-1.79
Idaho	11.82	0.53	47(21,51)	-9.57	Pennsylvania	14.11 ^b	0.18	12(07,22)	8.00
Illinois	14.00 ^b	0.19	14(07,23)	7.15	Rhode Island	13.11	0.60	24(05,48)	0.31
Indiana	13.95	0.27	15(06,26)	6.76	South Carolina	13.39	0.31	20(10,35)	2.50
Iowa	14.35	0.37	10(03,24)	9.79	South Dakota	13.18	0.70	23(03,49)	0.87
Kansas	12.84	0.39	30(13,46)	-1.75	Tennessee	14.76 ^b	0.28	07(03,17)	12.93 ^c
Kentucky	15.17 ^b	0.34	04(02,13)	16.09 ^c	Texas	12.53 ^b	0.14	34(26,42)	-4.11
Louisiana	15.14 ^b	0.34	05(02,13)	15.82 ^c	Utah	10.22 ^b	0.43	51(46,51)	-21.79 ^c
Maine	12.78	0.52	32(10,48)	-2.17	Vermont	13.30	0.80	21(03,49)	1.77
Maryland	12.76	0.27	33(19,44)	-2.38	Virginia	12.93	0.24	29(18,40)	-1.06
Massachusetts	12.19 ^b	0.24	41(27,48)	-6.68	Washington	12.20	0.25	40(26,48)	-6.66
Michigan	13.24	0.21	22(16,34)	1.36	West Virginia	14.99 ^b	0.48	06(01,21)	14.73 ^c
Minnesota	11.82 ^b	0.27	46(32,49)	-9.52	Wisconsin	12.31	0.26	36(24,47)	-5.79
Mississippi	16.52 ^b	0.43	02(01,05)	26.38 ^c	Wyoming	12.22	0.90	38(06,51)	-6.51
Missouri	13.90	0.27	16(07,27)	6.35					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 10% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 7.14
Cancer of the Corpus and Uterus, NOS (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	4.37	0.02		
High Five States				
D.C.	7.36 ^b	0.67	01(01,04)	68.33 ^c
Vermont	5.46	0.52	02(01,36)	24.99 ^c
New York	5.35 ^b	0.09	03(02,09)	22.45 ^c
New Jersey	5.33 ^b	0.14	04(02,11)	21.83 ^c
Maryland	5.29 ^b	0.18	05(02,13)	21.06 ^c
Low Five States				
New Mexico	3.26 ^b	0.23	47(34,51)	-25.45 ^c
Wyoming	3.26	0.46	48(13,51)	-25.45 ^c
Arkansas	3.22 ^b	0.19	49(37,51)	-26.44 ^c
Nevada	3.21 ^b	0.21	50(36,51)	-26.67 ^c
Alaska	2.94	0.50	51(16,51)	-32.80 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	3.46 ^b	0.15	44(35,51)	-20.78 ^c	Montana	3.81	0.35	37(10,51)	-12.95
Alaska	2.94	0.50	51(16,51)	-32.80 ^c	Nebraska	4.32	0.28	26(06,45)	-1.19
Arizona	3.42 ^b	0.13	45(36,51)	-21.73 ^c	Nevada	3.21 ^b	0.21	50(36,51)	-26.67 ^c
Arkansas	3.22 ^b	0.19	49(37,51)	-26.44 ^c	New Hampshire	4.84	0.34	12(02,38)	10.79
California	4.33	0.07	24(15,31)	-0.93	New Jersey	5.33 ^b	0.14	04(02,11)	21.83 ^c
Colorado	3.68 ^b	0.17	40(27,49)	-15.79 ^c	New Mexico	3.26 ^b	0.23	47(34,51)	-25.45 ^c
Connecticut	4.35	0.20	23(09,39)	-0.52	New York	5.35 ^b	0.09	03(02,09)	22.45 ^c
Delaware	5.21	0.43	06(01,35)	19.06 ^c	North Carolina	4.15	0.12	31(17,39)	-5.12
D.C.	7.36 ^b	0.67	01(01,04)	68.33 ^c	North Dakota	3.42	0.41	46(14,51)	-21.75 ^c
Florida	3.99 ^b	0.08	34(25,40)	-8.73	Ohio	4.90 ^b	0.12	11(04,18)	12.12
Georgia	4.03	0.13	33(19,42)	-7.83	Oklahoma	3.79	0.18	38(22,48)	-13.37
Hawaii	4.42	0.33	18(03,46)	1.16	Oregon	4.08	0.19	32(14,44)	-6.60
Idaho	4.37	0.32	21(03,45)	0.03	Pennsylvania	4.97 ^b	0.11	09(04,16)	13.65
Illinois	4.99 ^b	0.12	08(03,16)	14.18	Rhode Island	4.33	0.35	25(03,47)	-1.04
Indiana	4.50	0.15	16(09,32)	2.92	South Carolina	4.59	0.18	14(06,33)	5.02
Iowa	4.96	0.23	10(02,24)	13.53	South Dakota	4.82	0.45	13(02,44)	10.19
Kansas	4.16	0.22	30(11,45)	-4.75	Tennessee	3.74 ^b	0.14	39(27,48)	-14.46
Kentucky	3.60 ^b	0.17	43(30,50)	-17.58 ^c	Texas	3.66 ^b	0.08	42(34,47)	-16.27 ^c
Louisiana	3.67 ^b	0.17	41(26,50)	-16.07 ^c	Utah	3.87	0.26	36(13,50)	-11.52
Maine	4.36	0.30	22(04,45)	-0.29	Vermont	5.46	0.52	02(01,36)	24.99 ^c
Maryland	5.29 ^b	0.18	05(02,13)	21.06 ^c	Virginia	4.42	0.14	19(11,33)	1.05
Massachusetts	4.39	0.15	20(11,35)	0.43	Washington	4.18	0.15	29(14,40)	-4.30
Michigan	5.06 ^b	0.13	07(02,16)	15.82 ^c	West Virginia	4.27	0.26	27(07,44)	-2.29
Minnesota	4.22	0.17	28(13,40)	-3.38	Wisconsin	4.52	0.16	15(08,32)	3.42
Mississippi	3.90	0.21	35(16,48)	-10.89	Wyoming	3.26	0.46	48(13,51)	-25.45 ^c
Missouri	4.43	0.15	17(10,34)	1.44					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 8.18
Cancer of the Esophagus (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	4.18	0.02		
High Five States				
New Hampshire	5.55 ^b	0.27	01(01,13)	32.76 ^c
Maine	5.47 ^b	0.25	02(01,14)	30.88 ^c
Ohio	5.03 ^b	0.09	03(02,14)	20.35 ^c
West Virginia	4.97 ^b	0.21	04(01,26)	18.95 ^c
Indiana	4.93 ^b	0.12	05(02,19)	17.99 ^c
Low Five States				
California	3.47 ^b	0.04	47(46,50)	-16.94 ^c
Texas	3.38 ^b	0.06	48(46,50)	-19.10 ^c
New Mexico	3.37 ^b	0.18	49(43,51)	-19.52 ^c
Hawaii	3.08 ^b	0.20	50(46,51)	-26.41 ^c
Utah	2.91 ^b	0.17	51(48,51)	-30.47 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	4.06	0.12	39(22,46)	-2.99	Montana	4.29	0.27	30(03,47)	2.50
Alaska	4.46	0.44	22(01,50)	6.68	Nebraska	4.62	0.22	18(02,40)	10.61
Arizona	4.07	0.11	37(23,46)	-2.67	Nevada	4.78	0.19	11(02,31)	14.43
Arkansas	3.89	0.15	44(26,48)	-7.07	New Hampshire	5.55 ^b	0.27	01(01,13)	32.76 ^c
California	3.47 ^b	0.04	47(46,50)	-16.94 ^c	New Jersey	3.98	0.09	42(28,46)	-4.74
Colorado	3.98	0.13	43(25,46)	-4.91	New Mexico	3.37 ^b	0.18	49(43,51)	-19.52 ^c
Connecticut	4.11	0.14	36(19,46)	-1.80	New York	4.01	0.06	41(30,45)	-4.17
Delaware	4.32	0.29	27(03,47)	3.22	North Carolina	3.82 ^b	0.09	45(35,47)	-8.76
D.C.	4.73	0.41	16(01,46)	13.09	North Dakota	4.35	0.34	25(02,48)	4.07
Florida	4.04	0.06	40(29,44)	-3.49	Ohio	5.03 ^b	0.09	03(02,14)	20.35 ^c
Georgia	3.76 ^b	0.09	46(36,47)	-10.04	Oklahoma	4.49	0.15	20(07,37)	7.45
Hawaii	3.08 ^b	0.20	50(46,51)	-26.41 ^c	Oregon	4.39	0.14	24(10,41)	4.91
Idaho	4.16	0.23	34(09,47)	-0.49	Pennsylvania	4.81 ^b	0.08	10(04,20)	14.94
Illinois	4.46	0.08	21(13,31)	6.75	Rhode Island	4.78	0.28	12(01,41)	14.40
Indiana	4.93 ^b	0.12	05(02,19)	17.99 ^c	South Carolina	4.30	0.13	29(15,42)	2.94
Iowa	4.76 ^b	0.16	15(02,28)	13.81	South Dakota	4.82	0.32	09(01,43)	15.27 ^c
Kansas	4.23	0.17	31(14,46)	1.06	Tennessee	4.16	0.11	33(20,44)	-0.43
Kentucky	4.43	0.14	23(09,39)	5.97	Texas	3.38 ^b	0.06	48(46,50)	-19.10 ^c
Louisiana	4.34	0.14	26(12,42)	3.76	Utah	2.91 ^b	0.17	51(48,51)	-30.47 ^c
Maine	5.47 ^b	0.25	02(01,14)	30.88 ^c	Vermont	4.77	0.36	14(01,45)	14.03
Maryland	4.07	0.12	38(23,46)	-2.75	Virginia	4.18	0.10	32(21,43)	-0.03
Massachusetts	4.93 ^b	0.12	06(02,19)	17.91 ^c	Washington	4.71 ^b	0.12	17(04,27)	12.56
Michigan	4.82 ^b	0.09	08(03,20)	15.36 ^c	West Virginia	4.97 ^b	0.21	04(01,26)	18.95 ^c
Minnesota	4.58	0.13	19(06,32)	9.42	Wisconsin	4.77 ^b	0.12	13(03,24)	14.19
Mississippi	4.14	0.16	35(16,46)	-1.08	Wyoming	4.90	0.41	07(01,45)	17.26 ^c
Missouri	4.31	0.11	28(15,41)	2.97					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 8.19
Cancer of the Esophagus (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	7.45	0.03		
High Five States				
New Hampshire	10.13 ^b	0.56	01(01,13)	35.94 ^c
Maine	9.85 ^b	0.51	02(01,14)	32.15 ^c
Ohio	9.14 ^b	0.18	03(01,13)	22.62 ^c
West Virginia	9.09 ^b	0.42	04(01,23)	21.93 ^c
Indiana	8.97 ^b	0.24	05(02,16)	20.28 ^c
Low Five States				
Texas	6.10 ^b	0.11	47(45,50)	-18.12 ^c
California	6.05 ^b	0.09	48(45,50)	-18.85 ^c
New Mexico	5.80 ^b	0.34	49(43,51)	-22.26 ^c
Hawaii	5.58 ^b	0.40	50(43,51)	-25.20 ^c
Utah	5.15 ^b	0.33	51(47,51)	-30.95 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	7.30	0.25	37(17,46)	-2.13	Montana	7.40	0.52	33(06,49)	-0.75
Alaska	7.10	0.82	40(02,51)	-4.71	Nebraska	8.35	0.43	13(02,40)	11.95
Arizona	7.10	0.21	41(23,46)	-4.79	Nevada	8.08	0.37	18(04,41)	8.33
Arkansas	6.79	0.30	45(23,49)	-8.87	New Hampshire	10.13 ^b	0.56	01(01,13)	35.94 ^c
California	6.05 ^b	0.09	48(45,50)	-18.85 ^c	New Jersey	7.03	0.18	43(26,46)	-5.72
Colorado	7.08	0.26	42(20,46)	-5.04	New Mexico	5.80 ^b	0.34	49(43,51)	-22.26 ^c
Connecticut	7.37	0.29	34(15,46)	-1.07	New York	7.13	0.12	39(27,45)	-4.37
Delaware	7.43	0.57	32(03,49)	-0.33	North Carolina	7.00	0.18	44(28,46)	-6.09
D.C.	7.46	0.81	30(01,51)	0.06	North Dakota	8.26	0.70	15(01,46)	10.85
Florida	7.15	0.11	38(28,45)	-4.13	Ohio	9.14 ^b	0.18	03(01,13)	22.62 ^c
Georgia	6.64 ^b	0.19	46(35,48)	-10.89	Oklahoma	8.07	0.30	19(06,36)	8.19
Hawaii	5.58 ^b	0.40	50(43,51)	-25.20 ^c	Oregon	7.65	0.28	23(12,44)	2.65
Idaho	7.33	0.45	35(09,48)	-1.61	Pennsylvania	8.77 ^b	0.16	09(03,17)	17.62 ^c
Illinois	8.01	0.17	21(11,30)	7.41	Rhode Island	8.72	0.59	10(01,41)	16.95 ^c
Indiana	8.97 ^b	0.24	05(02,16)	20.28 ^c	South Carolina	7.65	0.26	24(12,44)	2.62
Iowa	8.78 ^b	0.33	08(02,24)	17.72 ^c	South Dakota	8.96	0.65	06(01,40)	20.22 ^c
Kansas	7.79	0.34	22(08,44)	4.51	Tennessee	7.52	0.22	29(16,44)	0.87
Kentucky	8.02	0.28	20(07,36)	7.56	Texas	6.10 ^b	0.11	47(45,50)	-18.12 ^c
Louisiana	7.62	0.27	27(12,44)	2.26	Utah	5.15 ^b	0.33	51(47,51)	-30.95 ^c
Maine	9.85 ^b	0.51	02(01,14)	32.15 ^c	Vermont	8.22	0.71	16(01,47)	10.30
Maryland	7.32	0.24	36(18,46)	-1.82	Virginia	7.62	0.21	26(14,41)	2.26
Massachusetts	8.94 ^b	0.24	07(02,17)	19.91 ^c	Washington	8.16	0.23	17(07,31)	9.48
Michigan	8.65 ^b	0.19	11(03,19)	16.04 ^c	West Virginia	9.09 ^b	0.42	04(01,23)	21.93 ^c
Minnesota	7.63	0.25	25(12,43)	2.42	Wisconsin	8.38 ^b	0.24	12(04,26)	12.44
Mississippi	7.44	0.33	31(12,46)	-0.18	Wyoming	8.32	0.79	14(01,47)	11.62
Missouri	7.53	0.22	28(15,44)	1.04					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 8.20
Cancer of the Esophagus (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	1.55	0.01		
High Five States				
D.C.	2.77	0.41	01(01,20)	78.92 ^c
Minnesota	2.05 ^b	0.12	02(01,16)	32.74 ^c
Vermont	1.95	0.31	03(01,49)	25.77 ^c
Alaska	1.88	0.40	04(01,51)	21.72 ^c
Wyoming	1.85	0.34	05(01,50)	19.22 ^c
Low Five States				
Kansas	1.28	0.12	47(20,51)	-17.23 ^c
Texas	1.17 ^b	0.04	48(43,51)	-24.16 ^c
North Dakota	1.03	0.22	49(10,51)	-33.24 ^c
Utah	0.97 ^b	0.13	50(41,51)	-37.16 ^c
Hawaii	0.97 ^b	0.15	51(35,51)	-37.62 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	1.47	0.10	34(11,48)	-5.06	Montana	1.49	0.22	32(02,51)	-3.92
Alaska	1.88	0.40	04(01,51)	21.72 ^c	Nebraska	1.61	0.17	25(02,49)	4.15
Arizona	1.48	0.09	33(12,48)	-4.40	Nevada	1.82	0.16	08(01,42)	17.66 ^c
Arkansas	1.45	0.13	40(07,49)	-6.38	New Hampshire	1.76	0.20	14(01,48)	13.72
California	1.41 ^b	0.04	43(29,46)	-9.07	New Jersey	1.63	0.08	23(06,41)	5.37
Colorado	1.42	0.11	42(11,49)	-8.37	New Mexico	1.30	0.15	46(11,51)	-15.71 ^c
Connecticut	1.59	0.12	27(04,47)	2.39	New York	1.61	0.05	26(11,37)	3.81
Delaware	1.77	0.24	12(01,49)	14.16	North Carolina	1.34	0.07	45(28,49)	-13.42
D.C.	2.77	0.41	01(01,20)	78.92 ^c	North Dakota	1.03	0.22	49(10,51)	-33.24 ^c
Florida	1.45	0.05	39(23,45)	-6.05	Ohio	1.81 ^b	0.07	09(03,26)	16.75 ^c
Georgia	1.46	0.08	36(16,47)	-5.39	Oklahoma	1.55	0.12	28(05,48)	-0.11
Hawaii	0.97 ^b	0.15	51(35,51)	-37.62 ^c	Oregon	1.66	0.12	21(03,44)	7.10
Idaho	1.37	0.18	44(05,51)	-11.33	Pennsylvania	1.67	0.06	20(06,33)	8.09
Illinois	1.71	0.07	16(04,32)	10.15	Rhode Island	1.80	0.22	10(01,47)	16.19 ^c
Indiana	1.70	0.09	17(03,38)	10.04	South Carolina	1.64	0.11	22(04,44)	5.85
Iowa	1.46	0.12	38(08,49)	-5.92	South Dakota	1.51	0.24	30(02,51)	-2.36
Kansas	1.28	0.12	47(20,51)	-17.23 ^c	Tennessee	1.46	0.09	37(13,48)	-5.79
Kentucky	1.47	0.11	35(10,48)	-5.17	Texas	1.17 ^b	0.04	48(43,51)	-24.16 ^c
Louisiana	1.70	0.11	19(03,42)	9.74	Utah	0.97 ^b	0.13	50(41,51)	-37.16 ^c
Maine	1.78	0.20	11(01,47)	15.17 ^c	Vermont	1.95	0.31	03(01,49)	25.77 ^c
Maryland	1.53	0.10	29(08,47)	-0.88	Virginia	1.44	0.08	41(17,48)	-7.25
Massachusetts	1.83	0.09	06(02,28)	18.19 ^c	Washington	1.83	0.10	07(02,31)	17.95 ^c
Michigan	1.72	0.07	15(04,34)	10.96	West Virginia	1.61	0.16	24(02,48)	4.16
Minnesota	2.05 ^b	0.12	02(01,16)	32.74 ^c	Wisconsin	1.77	0.10	13(02,33)	14.11
Mississippi	1.51	0.13	31(05,49)	-2.58	Wyoming	1.85	0.34	05(01,50)	19.22 ^c
Missouri	1.70	0.10	18(03,39)	9.88					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 9.18
Hodgkin Lymphoma

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	0.37	0.00		
High Five States				
Vermont	0.62	0.14	01(01,44)	66.76 ^c
Louisiana	0.50	0.05	02(01,25)	36.71 ^c
New Hampshire	0.48	0.09	03(01,44)	30.55 ^c
West Virginia	0.46	0.07	04(01,43)	25.65 ^c
Nebraska	0.45	0.07	05(01,44)	22.42 ^c
Low Five States				
Colorado	0.31	0.04	42(10,46)	-15.26 ^c
Kentucky	0.31	0.04	43(08,46)	-15.27 ^c
Arizona	0.27	0.03	44(27,46)	-26.46 ^c
Utah	0.25	0.05	45(12,46)	-32.46 ^c
Hawaii	0.19	0.05	46(26,46)	-47.20 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	0.41	0.04	13(02,42)	11.54	Montana	0.33	0.08	40(01,46)	-11.53
Alaska	-	-	-(-,-)	-	Nebraska	0.45	0.07	05(01,44)	22.42 ^c
Arizona	0.27	0.03	44(27,46)	-26.46 ^c	Nevada	0.36	0.05	28(02,46)	-1.86
Arkansas	0.38	0.05	22(02,45)	2.63	New Hampshire	0.48	0.09	03(01,44)	30.55 ^c
California	0.39	0.01	17(08,34)	4.42	New Jersey	0.34	0.03	34(10,44)	-7.17
Colorado	0.31	0.04	42(10,46)	-15.26 ^c	New Mexico	0.37	0.06	25(01,46)	0.23
Connecticut	0.37	0.04	27(03,45)	-0.38	New York	0.39	0.02	15(05,35)	6.97
Delaware	0.38	0.09	21(01,46)	3.01	North Carolina	0.35	0.03	31(07,44)	-3.87
D.C.	-	-	-(-,-)	-	North Dakota	-	-	-(-,-)	-
Florida	0.37	0.02	26(08,39)	0.19	Ohio	0.39	0.03	16(05,39)	4.65
Georgia	0.35	0.03	33(07,44)	-4.99	Oklahoma	0.44	0.05	08(01,40)	17.99 ^c
Hawaii	0.19	0.05	46(26,46)	-47.20 ^c	Oregon	0.42	0.04	10(01,42)	13.92
Idaho	0.44	0.08	07(01,45)	19.29 ^c	Pennsylvania	0.38	0.02	20(06,39)	3.54
Illinois	0.33	0.02	39(14,44)	-10.61	Rhode Island	0.43	0.09	09(01,46)	15.98 ^c
Indiana	0.36	0.03	29(05,44)	-3.25	South Carolina	0.36	0.04	30(04,45)	-3.49
Iowa	0.33	0.04	38(04,46)	-10.31	South Dakota	-	-	-(-,-)	-
Kansas	0.38	0.05	19(02,45)	4.05	Tennessee	0.41	0.04	12(02,40)	11.66
Kentucky	0.31	0.04	43(08,46)	-15.27 ^c	Texas	0.37	0.02	24(08,38)	1.03
Louisiana	0.50	0.05	02(01,25)	36.71 ^c	Utah	0.25	0.05	45(12,46)	-32.46 ^c
Maine	0.45	0.08	06(01,45)	21.04 ^c	Vermont	0.62	0.14	01(01,44)	66.76 ^c
Maryland	0.34	0.03	36(07,45)	-8.67	Virginia	0.38	0.03	23(05,43)	1.74
Massachusetts	0.35	0.03	32(06,44)	-4.35	Washington	0.32	0.03	41(12,45)	-13.68
Michigan	0.41	0.03	14(03,37)	11.24	West Virginia	0.46	0.07	04(01,43)	25.65 ^c
Minnesota	0.34	0.04	35(06,45)	-7.25	Wisconsin	0.42	0.04	11(02,39)	13.56
Mississippi	0.38	0.05	18(02,45)	4.34	Wyoming	-	-	-(-,-)	-
Missouri	0.33	0.03	37(08,45)	-9.57					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 9.19
Hodgkin Lymphoma

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	0.46	0.01		
High Five States				
Rhode Island	0.72	0.17	01(01,41)	56.20 ^c
New Hampshire	0.68	0.16	02(01,41)	48.19 ^c
Nebraska	0.67	0.12	03(01,39)	45.54 ^c
Louisiana	0.62	0.08	04(01,32)	35.19 ^c
Oregon	0.57	0.08	05(01,37)	24.20 ^c
Low Five States				
Maryland	0.38	0.05	38(07,42)	-17.55 ^c
Colorado	0.34	0.06	39(11,42)	-27.02 ^c
Kentucky	0.34	0.06	40(13,42)	-27.31 ^c
Arizona	0.31 ^b	0.04	41(26,42)	-33.70 ^c
Utah	0.30	0.08	42(08,42)	-34.27 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	0.45	0.06	26(03,42)	-1.60	Montana	-	-	-(-,-)	-
Alaska	-	-	-(-,-)	-	Nebraska	0.67	0.12	03(01,39)	45.54 ^c
Arizona	0.31 ^b	0.04	41(26,42)	-33.70 ^c	Nevada	0.52	0.10	07(01,42)	12.30
Arkansas	0.42	0.08	32(03,42)	-9.21	New Hampshire	0.68	0.16	02(01,41)	48.19 ^c
California	0.49	0.02	16(05,31)	6.55	New Jersey	0.42	0.05	31(06,41)	-9.07
Colorado	0.34	0.06	39(11,42)	-27.02 ^c	New Mexico	0.52	0.10	08(01,42)	11.84
Connecticut	0.38	0.07	37(05,42)	-16.51 ^c	New York	0.50	0.03	14(04,33)	8.67
Delaware	-	-	-(-,-)	-	North Carolina	0.45	0.05	27(05,40)	-2.67
D.C.	-	-	-(-,-)	-	North Dakota	-	-	-(-,-)	-
Florida	0.46	0.03	24(07,37)	0.00	Ohio	0.48	0.04	20(04,38)	3.90
Georgia	0.48	0.05	21(03,39)	3.08	Oklahoma	0.49	0.08	17(01,41)	6.54
Hawaii	-	-	-(-,-)	-	Oregon	0.57	0.08	05(01,37)	24.20 ^c
Idaho	0.49	0.11	18(01,42)	6.31	Pennsylvania	0.45	0.04	28(06,39)	-3.08
Illinois	0.42	0.04	33(09,41)	-9.39	Rhode Island	0.72	0.17	01(01,41)	56.20 ^c
Indiana	0.47	0.06	22(03,40)	2.56	South Carolina	0.46	0.07	25(02,41)	-0.08
Iowa	0.41	0.07	34(03,42)	-10.24	South Dakota	-	-	-(-,-)	-
Kansas	0.44	0.08	29(02,42)	-3.63	Tennessee	0.51	0.06	10(02,38)	11.21
Kentucky	0.34	0.06	40(13,42)	-27.31 ^c	Texas	0.51	0.03	12(03,31)	10.02
Louisiana	0.62	0.08	04(01,32)	35.19 ^c	Utah	0.30	0.08	42(08,42)	-34.27 ^c
Maine	0.52	0.13	06(01,42)	12.57	Vermont	-	-	-(-,-)	-
Maryland	0.38	0.05	38(07,42)	-17.55 ^c	Virginia	0.52	0.05	09(02,37)	11.81
Massachusetts	0.46	0.05	23(03,40)	0.53	Washington	0.40	0.05	35(06,42)	-12.31
Michigan	0.51	0.05	13(02,37)	9.72	West Virginia	0.49	0.10	19(01,42)	5.37
Minnesota	0.43	0.06	30(04,42)	-7.40	Wisconsin	0.49	0.06	15(02,39)	7.30
Mississippi	0.51	0.09	11(01,42)	10.80	Wyoming	-	-	-(-,-)	-
Missouri	0.39	0.05	36(07,42)	-16.35 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 9.20
Hodgkin Lymphoma

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	0.29	0.01		
High Five States				
West Virginia	0.43	0.09	01(01,35)	48.30 ^c
Louisiana	0.41	0.06	02(01,29)	40.53 ^c
Oklahoma	0.41	0.06	03(01,32)	39.24 ^c
Alabama	0.37	0.05	04(01,33)	27.53 ^c
Maine	0.37	0.10	05(01,36)	25.00 ^c
Low Five States				
Texas	0.26	0.02	32(11,36)	-10.52
Illinois	0.26	0.03	33(09,36)	-11.89
Washington	0.25	0.04	34(07,36)	-15.89 ^c
Arizona	0.24	0.04	35(07,36)	-18.33 ^c
Nevada	0.23	0.06	36(03,36)	-20.90 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	0.37	0.05	04(01,33)	27.53 ^c	Montana	-	-	-(-,-)	-
Alaska	-	-	-(-,-)	-	Nebraska	-	-	-(-,-)	-
Arizona	0.24	0.04	35(07,36)	-18.33 ^c	Nevada	0.23	0.06	36(03,36)	-20.90 ^c
Arkansas	0.35	0.06	06(01,36)	20.02 ^c	New Hampshire	-	-	-(-,-)	-
California	0.30	0.02	16(07,31)	2.27	New Jersey	0.28	0.03	23(05,36)	-5.03
Colorado	0.29	0.05	19(02,36)	-0.28	New Mexico	-	-	-(-,-)	-
Connecticut	0.34	0.06	08(01,36)	17.49 ^c	New York	0.31	0.02	14(04,32)	5.58
Delaware	-	-	-(-,-)	-	North Carolina	0.28	0.03	24(05,36)	-5.45
D.C.	-	-	-(-,-)	-	North Dakota	-	-	-(-,-)	-
Florida	0.29	0.02	22(07,34)	-1.89	Ohio	0.31	0.03	13(03,34)	5.59
Georgia	0.27	0.03	27(05,36)	-6.33	Oklahoma	0.41	0.06	03(01,32)	39.24 ^c
Hawaii	-	-	-(-,-)	-	Oregon	0.29	0.05	21(02,36)	-1.86
Idaho	-	-	-(-,-)	-	Pennsylvania	0.33	0.03	10(02,29)	13.91
Illinois	0.26	0.03	33(09,36)	-11.89	Rhode Island	-	-	-(-,-)	-
Indiana	0.28	0.04	25(04,36)	-5.68	South Carolina	0.27	0.05	29(03,36)	-8.40
Iowa	0.27	0.05	28(02,36)	-7.60	South Dakota	-	-	-(-,-)	-
Kansas	0.34	0.07	09(01,36)	15.74 ^c	Tennessee	0.32	0.04	12(01,35)	9.92
Kentucky	0.29	0.05	20(02,36)	-1.64	Texas	0.26	0.02	32(11,36)	-10.52
Louisiana	0.41	0.06	02(01,29)	40.53 ^c	Utah	-	-	-(-,-)	-
Maine	0.37	0.10	05(01,36)	25.00 ^c	Vermont	-	-	-(-,-)	-
Maryland	0.29	0.04	18(02,36)	0.22	Virginia	0.27	0.03	31(05,36)	-9.57
Massachusetts	0.27	0.04	30(05,36)	-8.54	Washington	0.25	0.04	34(07,36)	-15.89 ^c
Michigan	0.33	0.03	11(02,32)	12.43	West Virginia	0.43	0.09	01(01,35)	48.30 ^c
Minnesota	0.28	0.04	26(03,36)	-6.06	Wisconsin	0.35	0.05	07(01,34)	17.74 ^c
Mississippi	0.30	0.06	15(01,36)	2.52	Wyoming	-	-	-(-,-)	-
Missouri	0.30	0.04	17(02,36)	0.99					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 11.18
Cancer of the Kidney and Renal Pelvis (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	3.92	0.02		
High Five States				
Oklahoma	5.30 ^b	0.16	01(01,04)	35.23 ^c
Louisiana	5.14 ^b	0.15	02(01,07)	31.05 ^c
North Dakota	4.96	0.36	03(01,29)	26.46 ^c
Arkansas	4.65 ^b	0.17	04(02,22)	18.54 ^c
Kentucky	4.62 ^b	0.14	05(02,20)	17.77 ^c
Low Five States				
Wyoming	3.30	0.33	47(15,51)	-15.95 ^c
Massachusetts	3.29 ^b	0.09	48(37,50)	-16.05 ^c
Connecticut	3.27 ^b	0.13	49(35,50)	-16.63 ^c
D.C.	2.79 ^b	0.31	50(32,51)	-28.74 ^c
Hawaii	2.68 ^b	0.18	51(46,51)	-31.71 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	3.96	0.12	28(15,37)	0.94	Montana	3.43	0.24	43(20,51)	-12.54
Alaska	4.44	0.45	14(01,49)	13.29	Nebraska	4.43	0.21	15(02,32)	12.86
Arizona	3.75	0.10	33(22,43)	-4.48	Nevada	3.76	0.17	32(16,48)	-4.09
Arkansas	4.65 ^b	0.17	04(02,22)	18.54 ^c	New Hampshire	3.66	0.22	35(14,50)	-6.67
California	3.52 ^b	0.04	39(34,45)	-10.16	New Jersey	3.38 ^b	0.08	45(35,49)	-13.81
Colorado	3.47 ^b	0.12	42(30,49)	-11.65	New Mexico	4.48	0.20	13(02,30)	14.18
Connecticut	3.27 ^b	0.13	49(35,50)	-16.63 ^c	New York	3.30 ^b	0.06	46(39,50)	-15.76 ^c
Delaware	3.50	0.26	41(15,51)	-10.80	North Carolina	3.97	0.09	27(18,36)	1.19
D.C.	2.79 ^b	0.31	50(32,51)	-28.74 ^c	North Dakota	4.96	0.36	03(01,29)	26.46 ^c
Florida	3.42 ^b	0.05	44(36,48)	-12.79	Ohio	4.27 ^b	0.08	19(09,26)	8.83
Georgia	3.51 ^b	0.09	40(32,48)	-10.43	Oklahoma	5.30 ^b	0.16	01(01,04)	35.23 ^c
Hawaii	2.68 ^b	0.18	51(46,51)	-31.71 ^c	Oregon	3.94	0.13	29(15,40)	0.47
Idaho	4.07	0.23	24(05,44)	3.66	Pennsylvania	3.98	0.07	26(19,34)	1.51
Illinois	4.13	0.08	22(14,30)	5.20	Rhode Island	3.55	0.24	38(17,50)	-9.61
Indiana	4.56 ^b	0.12	09(03,20)	16.22 ^c	South Carolina	4.07	0.13	23(12,35)	3.73
Iowa	4.59 ^b	0.16	07(02,22)	17.06 ^c	South Dakota	4.41	0.31	16(01,40)	12.40
Kansas	4.57 ^b	0.17	08(02,24)	16.53 ^c	Tennessee	4.19	0.11	20(09,31)	6.79
Kentucky	4.62 ^b	0.14	05(02,20)	17.77 ^c	Texas	4.54 ^b	0.06	11(04,17)	15.78 ^c
Louisiana	5.14 ^b	0.15	02(01,07)	31.05 ^c	Utah	3.56	0.19	37(21,50)	-9.20
Maine	4.01	0.22	25(05,45)	2.32	Vermont	3.56	0.31	36(10,51)	-9.13
Maryland	3.70	0.11	34(24,45)	-5.64	Virginia	3.81	0.10	31(21,41)	-2.98
Massachusetts	3.29 ^b	0.09	48(37,50)	-16.05 ^c	Washington	3.92	0.11	30(18,38)	-0.13
Michigan	4.13	0.09	21(13,31)	5.37	West Virginia	4.62	0.20	06(01,26)	17.71 ^c
Minnesota	4.27	0.12	18(06,29)	8.87	Wisconsin	4.34 ^b	0.12	17(04,26)	10.59
Mississippi	4.55 ^b	0.17	10(02,25)	16.05 ^c	Wyoming	3.30	0.33	47(15,51)	-15.95 ^c
Missouri	4.51 ^b	0.12	12(03,21)	14.92					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 11.19
Cancer of the Kidney and Renal Pelvis (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	5.73	0.03		
High Five States				
Oklahoma	7.82 ^b	0.30	01(01,05)	36.52 ^c
North Dakota	7.29	0.66	02(01,35)	27.37 ^c
Louisiana	7.28 ^b	0.27	03(01,13)	27.08 ^c
Iowa	6.93 ^b	0.30	04(01,20)	20.97 ^c
Mississippi	6.76	0.32	05(01,26)	18.08 ^c
Low Five States				
Wyoming	4.81	0.59	47(08,51)	-15.93 ^c
Delaware	4.72	0.47	48(20,51)	-17.64 ^c
Massachusetts	4.69 ^b	0.17	49(37,51)	-18.09 ^c
D.C.	4.65	0.63	50(09,51)	-18.80 ^c
Hawaii	4.20 ^b	0.34	51(37,51)	-26.72 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	5.82	0.23	25(11,41)	1.62	Montana	4.99	0.43	43(14,51)	-12.85
Alaska	5.99	0.79	21(01,51)	4.65	Nebraska	6.46	0.38	13(02,37)	12.80
Arizona	5.44	0.19	35(21,46)	-5.03	Nevada	5.28	0.30	37(17,50)	-7.80
Arkansas	6.67	0.30	06(01,27)	16.57 ^c	New Hampshire	5.72	0.42	30(04,49)	-0.09
California	5.09 ^b	0.08	39(34,48)	-11.16	New Jersey	5.14 ^b	0.16	38(30,49)	-10.23
Colorado	5.07	0.22	41(26,50)	-11.38	New Mexico	6.02	0.35	20(03,44)	5.20
Connecticut	4.94	0.24	45(28,51)	-13.64	New York	5.02 ^b	0.11	42(35,49)	-12.40
Delaware	4.72	0.47	48(20,51)	-17.64 ^c	North Carolina	6.13	0.17	18(07,32)	7.02
D.C.	4.65	0.63	50(09,51)	-18.80 ^c	North Dakota	7.29	0.66	02(01,35)	27.37 ^c
Florida	4.98 ^b	0.09	44(36,49)	-12.97	Ohio	5.94	0.14	23(13,35)	3.72
Georgia	5.39	0.18	36(23,46)	-5.86	Oklahoma	7.82 ^b	0.30	01(01,05)	36.52 ^c
Hawaii	4.20 ^b	0.34	51(37,51)	-26.72 ^c	Oregon	5.69	0.24	32(13,44)	-0.61
Idaho	5.81	0.40	27(04,49)	1.50	Pennsylvania	5.67	0.13	33(19,39)	-1.03
Illinois	5.95	0.15	22(13,34)	3.95	Rhode Island	5.82	0.48	26(02,50)	1.60
Indiana	6.57 ^b	0.21	09(03,24)	14.71	South Carolina	6.33	0.25	17(03,31)	10.65
Iowa	6.93 ^b	0.30	04(01,20)	20.97 ^c	South Dakota	5.82	0.53	24(02,50)	1.70
Kansas	6.65	0.31	07(01,29)	16.12 ^c	Tennessee	6.37	0.21	15(03,28)	11.33
Kentucky	6.52	0.26	11(02,29)	13.91	Texas	6.53 ^b	0.12	10(04,20)	14.11
Louisiana	7.28 ^b	0.27	03(01,13)	27.08 ^c	Utah	5.08	0.33	40(19,51)	-11.24
Maine	6.11	0.41	19(02,46)	6.71	Vermont	4.82	0.55	46(12,51)	-15.88 ^c
Maryland	5.61	0.21	34(16,44)	-2.03	Virginia	5.74	0.18	29(15,40)	0.27
Massachusetts	4.69 ^b	0.17	49(37,51)	-18.09 ^c	Washington	5.80	0.20	28(13,39)	1.24
Michigan	5.70	0.15	31(17,39)	-0.51	West Virginia	6.52	0.36	12(01,35)	13.83
Minnesota	6.40	0.23	14(03,30)	11.86	Wisconsin	6.35	0.21	16(04,29)	10.86
Mississippi	6.76	0.32	05(01,26)	18.08 ^c	Wyoming	4.81	0.59	47(08,51)	-15.93 ^c
Missouri	6.57 ^b	0.21	08(02,24)	14.71					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 11.20
Cancer of the Kidney And Renal Pelvis (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	2.51	0.02		
High Five States				
Louisiana	3.42 ^b	0.16	01(01,12)	36.38 ^c
Oklahoma	3.31 ^b	0.17	02(01,17)	31.93 ^c
South Dakota	3.21	0.35	03(01,39)	28.11 ^c
New Mexico	3.16	0.23	04(01,27)	26.17 ^c
West Virginia	3.15	0.22	05(01,27)	25.59 ^c
Low Five States				
Rhode Island	1.99	0.24	47(19,51)	-20.69 ^c
Wyoming	1.98	0.36	48(05,51)	-20.91 ^c
New Hampshire	1.94	0.22	49(21,51)	-22.60 ^c
Hawaii	1.45 ^b	0.19	50(45,51)	-42.32 ^c
D.C.	1.44 ^b	0.29	51(29,51)	-42.40 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	2.55	0.13	24(11,41)	1.72	Montana	2.03	0.25	45(14,51)	-19.12 ^c
Alaska	3.07	0.52	07(01,50)	22.60 ^c	Nebraska	2.70	0.22	20(02,45)	7.56
Arizona	2.31	0.11	35(21,47)	-7.95	Nevada	2.41	0.19	31(09,49)	-3.91
Arkansas	3.02	0.18	09(01,28)	20.55 ^c	New Hampshire	1.94	0.22	49(21,51)	-22.60 ^c
California	2.27 ^b	0.05	38(28,44)	-9.56	New Jersey	2.06 ^b	0.09	43(33,49)	-17.95 ^c
Colorado	2.14	0.13	41(24,50)	-14.59	New Mexico	3.16	0.23	04(01,27)	26.17 ^c
Connecticut	2.00 ^b	0.13	46(28,50)	-20.29 ^c	New York	2.05 ^b	0.06	44(36,49)	-18.20 ^c
Delaware	2.58	0.30	22(01,49)	2.87	North Carolina	2.41	0.09	30(19,43)	-3.88
D.C.	1.44 ^b	0.29	51(29,51)	-42.40 ^c	North Dakota	3.07	0.38	08(01,46)	22.46 ^c
Florida	2.15 ^b	0.06	40(32,48)	-14.11	Ohio	2.96 ^b	0.09	11(03,21)	18.27 ^c
Georgia	2.13 ^b	0.09	42(28,49)	-14.80	Oklahoma	3.31 ^b	0.17	02(01,17)	31.93 ^c
Hawaii	1.45 ^b	0.19	50(45,51)	-42.32 ^c	Oregon	2.50	0.15	27(10,44)	-0.20
Idaho	2.52	0.25	25(03,49)	0.55	Pennsylvania	2.67	0.08	21(11,30)	6.55
Illinois	2.75	0.09	18(08,28)	9.64	Rhode Island	1.99	0.24	47(19,51)	-20.69 ^c
Indiana	2.97 ^b	0.12	10(02,24)	18.69 ^c	South Carolina	2.37	0.13	32(17,47)	-5.40
Iowa	2.79	0.17	17(03,36)	11.33	South Dakota	3.21	0.35	03(01,39)	28.11 ^c
Kansas	2.91	0.19	14(01,32)	16.32 ^c	Tennessee	2.50	0.11	28(14,41)	-0.24
Kentucky	3.15 ^b	0.15	06(01,21)	25.54 ^c	Texas	2.94 ^b	0.07	12(04,20)	17.29 ^c
Louisiana	3.42 ^b	0.16	01(01,12)	36.38 ^c	Utah	2.29	0.20	37(12,49)	-8.70
Maine	2.44	0.23	29(05,49)	-2.69	Vermont	2.51	0.35	26(01,50)	0.10
Maryland	2.30	0.12	36(20,47)	-8.17	Virginia	2.33	0.10	34(21,46)	-6.99
Massachusetts	2.26	0.10	39(23,48)	-9.95	Washington	2.37	0.11	33(18,45)	-5.46
Michigan	2.89 ^b	0.10	15(04,24)	15.42 ^c	West Virginia	3.15	0.22	05(01,27)	25.59 ^c
Minnesota	2.58	0.13	23(10,40)	2.80	Wisconsin	2.70	0.12	19(07,34)	7.62
Mississippi	2.81	0.18	16(03,36)	12.26	Wyoming	1.98	0.36	48(05,51)	-20.91 ^c
Missouri	2.92	0.12	13(02,26)	16.73 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 12.18
Cancer of the Larynx (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	1.08	0.01		
High Five States				
D.C.	1.85	0.26	01(01,22)	71.22 ^c
West Virginia	1.63 ^b	0.12	02(01,10)	50.81 ^c
Kentucky	1.46 ^b	0.08	03(01,14)	35.10 ^c
Alabama	1.40 ^b	0.07	04(01,17)	29.45 ^c
Tennessee	1.35 ^b	0.06	05(02,18)	25.39 ^c
Low Five States				
Oregon	0.75 ^b	0.06	47(35,50)	-30.66 ^c
Colorado	0.70 ^b	0.06	48(38,50)	-35.14 ^c
Idaho	0.62 ^b	0.09	49(37,51)	-42.58 ^c
Wyoming	0.58 ^b	0.14	50(30,51)	-46.50 ^c
Utah	0.38 ^b	0.06	51(49,51)	-65.24 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	1.40 ^b	0.07	04(01,17)	29.45 ^c	Montana	1.16	0.14	19(02,44)	7.80
Alaska	0.98	0.21	36(02,50)	-9.32	Nebraska	0.79	0.09	45(28,50)	-26.97 ^c
Arizona	0.79 ^b	0.05	43(35,49)	-26.70 ^c	Nevada	1.09	0.09	26(06,41)	0.91
Arkansas	1.09	0.08	27(08,40)	0.77	New Hampshire	1.00	0.12	34(06,48)	-7.62
California	0.80 ^b	0.02	42(37,48)	-25.84 ^c	New Jersey	1.14	0.05	22(10,35)	5.21
Colorado	0.70 ^b	0.06	48(38,50)	-35.14 ^c	New Mexico	0.99	0.09	35(11,47)	-8.70
Connecticut	0.92	0.07	37(23,47)	-15.21 ^c	New York	1.13	0.03	24(12,32)	4.38
Delaware	1.28	0.15	10(01,40)	18.27 ^c	North Carolina	1.17	0.05	18(08,32)	7.98
D.C.	1.85	0.26	01(01,22)	71.22 ^c	North Dakota	1.03	0.16	30(02,50)	-4.38
Florida	1.20 ^b	0.03	15(09,26)	11.60	Ohio	1.35 ^b	0.05	07(03,16)	24.74 ^c
Georgia	1.08	0.05	28(12,37)	0.09	Oklahoma	1.02	0.07	32(12,42)	-5.85
Hawaii	0.80	0.10	40(24,50)	-25.42 ^c	Oregon	0.75 ^b	0.06	47(35,50)	-30.66 ^c
Idaho	0.62 ^b	0.09	49(37,51)	-42.58 ^c	Pennsylvania	1.16	0.04	20(09,31)	7.77
Illinois	1.16	0.04	21(10,32)	7.30	Rhode Island	1.24	0.14	12(01,42)	14.67
Indiana	1.20	0.06	16(06,32)	11.45	South Carolina	1.35 ^b	0.07	06(02,22)	24.90 ^c
Iowa	1.00	0.07	33(13,44)	-7.29	South Dakota	0.78	0.13	46(15,50)	-27.34 ^c
Kansas	0.79 ^b	0.07	44(30,50)	-26.76 ^c	Tennessee	1.35 ^b	0.06	05(02,18)	25.39 ^c
Kentucky	1.46 ^b	0.08	03(01,14)	35.10 ^c	Texas	1.13	0.03	23(12,32)	4.56
Louisiana	1.33	0.07	08(02,25)	23.32 ^c	Utah	0.38 ^b	0.06	51(49,51)	-65.24 ^c
Maine	1.28	0.12	11(01,36)	18.21 ^c	Vermont	1.21	0.18	14(01,47)	11.81
Maryland	1.22	0.06	13(04,31)	12.72	Virginia	1.03	0.05	29(16,39)	-4.09
Massachusetts	1.03	0.05	31(16,39)	-4.86	Washington	0.91	0.05	38(27,46)	-15.83 ^c
Michigan	1.19	0.05	17(07,30)	10.49	West Virginia	1.63 ^b	0.12	02(01,10)	50.81 ^c
Minnesota	0.80 ^b	0.05	41(33,49)	-25.54 ^c	Wisconsin	0.86 ^b	0.05	39(30,48)	-20.48 ^c
Mississippi	1.32	0.09	09(02,29)	21.89 ^c	Wyoming	0.58 ^b	0.14	50(30,51)	-46.50 ^c
Missouri	1.11	0.06	25(09,36)	2.98					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 12.19
Cancer of the Larynx (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	1.93	0.02		
High Five States				
D.C.	3.57	0.55	01(01,23)	85.14 ^c
West Virginia	2.74 ^b	0.23	02(01,17)	42.00 ^c
Alabama	2.54 ^b	0.15	03(01,17)	31.89 ^c
South Carolina	2.50 ^b	0.15	04(01,20)	29.99 ^c
Kentucky	2.47 ^b	0.15	05(01,22)	28.21 ^c
Low Five States				
Kansas	1.34 ^b	0.14	46(32,49)	-30.64 ^c
Oregon	1.26 ^b	0.11	47(35,49)	-34.44 ^c
Colorado	1.14 ^b	0.11	48(40,49)	-40.85 ^c
Idaho	1.05 ^b	0.17	49(38,50)	-45.51 ^c
Utah	0.65 ^b	0.12	50(49,50)	-66.22 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	2.54 ^b	0.15	03(01,17)	31.89 ^c	Montana	2.13	0.28	15(02,45)	10.34
Alaska	1.72	0.43	36(01,50)	-10.46	Nebraska	1.40	0.18	44(23,49)	-27.29 ^c
Arizona	1.38 ^b	0.09	45(34,49)	-28.25 ^c	Nevada	1.76	0.17	33(10,46)	-8.68
Arkansas	1.95	0.16	26(06,41)	1.34	New Hampshire	1.78	0.24	32(05,49)	-7.52
California	1.48 ^b	0.04	42(35,46)	-23.30 ^c	New Jersey	1.94	0.10	28(12,37)	0.79
Colorado	1.14 ^b	0.11	48(40,49)	-40.85 ^c	New Mexico	1.75	0.19	34(09,47)	-9.06
Connecticut	1.67	0.14	37(17,46)	-13.37	New York	2.02	0.07	21(11,32)	4.90
Delaware	1.94	0.28	27(02,47)	0.83	North Carolina	2.15	0.10	13(05,31)	11.76
D.C.	3.57	0.55	01(01,23)	85.14 ^c	North Dakota	1.78	0.32	31(02,49)	-7.44
Florida	2.14 ^b	0.06	14(08,27)	10.93	Ohio	2.40 ^b	0.09	07(02,17)	24.64 ^c
Georgia	2.00	0.10	23(09,35)	3.84	Oklahoma	1.83	0.14	30(11,44)	-5.00
Hawaii	1.54	0.21	40(12,49)	-19.98 ^c	Oregon	1.26 ^b	0.11	47(35,49)	-34.44 ^c
Idaho	1.05 ^b	0.17	49(38,50)	-45.51 ^c	Pennsylvania	2.08	0.08	18(09,31)	8.22
Illinois	2.09	0.09	17(08,31)	8.68	Rhode Island	2.19	0.29	12(01,43)	13.87
Indiana	2.01	0.11	22(07,36)	4.34	South Carolina	2.50 ^b	0.15	04(01,20)	29.99 ^c
Iowa	1.75	0.15	35(13,45)	-9.09	South Dakota	1.52	0.28	41(08,50)	-20.87 ^c
Kansas	1.34 ^b	0.14	46(32,49)	-30.64 ^c	Tennessee	2.42 ^b	0.13	06(01,22)	25.56 ^c
Kentucky	2.47 ^b	0.15	05(01,22)	28.21 ^c	Texas	2.08	0.07	19(09,30)	8.15
Louisiana	2.31	0.15	10(02,30)	20.14 ^c	Utah	0.65 ^b	0.12	50(49,50)	-66.22 ^c
Maine	2.36	0.25	09(01,35)	22.57 ^c	Vermont	2.06	0.35	20(01,48)	6.72
Maryland	2.24	0.13	11(03,30)	16.28 ^c	Virginia	1.96	0.11	25(09,37)	1.67
Massachusetts	1.85	0.11	29(13,40)	-4.05	Washington	1.55 ^b	0.10	39(28,47)	-19.62 ^c
Michigan	2.11	0.09	16(07,31)	9.67	West Virginia	2.74 ^b	0.23	02(01,17)	42.00 ^c
Minnesota	1.44 ^b	0.11	43(31,48)	-25.51 ^c	Wisconsin	1.57	0.11	38(27,46)	-18.30 ^c
Mississippi	2.40	0.19	08(01,30)	24.39 ^c	Wyoming	-	-	-(-,-)	-
Missouri	1.96	0.12	24(09,37)	1.99					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 12.20
Cancer of the Larynx (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	0.41	0.01		
High Five States				
Delaware	0.73	0.16	01(01,35)	78.44 ^c
West Virginia	0.71	0.10	02(01,20)	73.50 ^c
Kentucky	0.67 ^b	0.07	03(01,13)	64.13 ^c
Louisiana	0.57	0.07	04(01,25)	37.65 ^c
Rhode Island	0.56	0.13	05(01,39)	36.46 ^c
Low Five States				
Oregon	0.33	0.05	35(11,39)	-19.30 ^c
Wisconsin	0.30	0.04	36(18,39)	-25.74 ^c
Minnesota	0.30	0.04	37(17,39)	-26.32 ^c
Arizona	0.29	0.04	38(21,39)	-28.39 ^c
California	0.27 ^b	0.02	39(31,39)	-33.75 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	0.49	0.06	11(02,33)	18.49 ^c	Montana	-	-	-(-,-)	-
Alaska	-	-	-(-,-)	-	Nebraska	-	-	-(-,-)	-
Arizona	0.29	0.04	38(21,39)	-28.39 ^c	Nevada	0.51	0.09	10(01,37)	23.64 ^c
Arkansas	0.38	0.07	26(05,39)	-7.25	New Hampshire	-	-	-(-,-)	-
California	0.27 ^b	0.02	39(31,39)	-33.75 ^c	New Jersey	0.53	0.04	07(02,23)	29.52 ^c
Colorado	0.36	0.05	30(08,39)	-12.81	New Mexico	0.34	0.08	33(05,39)	-17.89 ^c
Connecticut	0.33	0.05	34(10,39)	-18.78 ^c	New York	0.46	0.03	14(05,27)	13.07
Delaware	0.73	0.16	01(01,35)	78.44 ^c	North Carolina	0.42	0.04	21(07,35)	1.92
D.C.	-	-	-(-,-)	-	North Dakota	-	-	-(-,-)	-
Florida	0.42	0.02	20(10,31)	2.90	Ohio	0.52	0.04	08(02,22)	26.98 ^c
Georgia	0.38	0.04	27(11,38)	-7.67	Oklahoma	0.36	0.06	28(07,39)	-11.22
Hawaii	-	-	-(-,-)	-	Oregon	0.33	0.05	35(11,39)	-19.30 ^c
Idaho	-	-	-(-,-)	-	Pennsylvania	0.45	0.03	16(05,29)	10.73
Illinois	0.45	0.03	17(06,31)	8.65	Rhode Island	0.56	0.13	05(01,39)	36.46 ^c
Indiana	0.55	0.05	06(01,24)	34.07 ^c	South Carolina	0.44	0.06	18(03,37)	7.55
Iowa	0.42	0.07	22(03,39)	1.17	South Dakota	-	-	-(-,-)	-
Kansas	0.34	0.06	32(07,39)	-17.06 ^c	Tennessee	0.52	0.05	09(02,27)	25.93 ^c
Kentucky	0.67 ^b	0.07	03(01,13)	64.13 ^c	Texas	0.36	0.02	29(17,37)	-11.23
Louisiana	0.57	0.07	04(01,25)	37.65 ^c	Utah	-	-	-(-,-)	-
Maine	0.38	0.09	25(02,39)	-6.33	Vermont	-	-	-(-,-)	-
Maryland	0.43	0.05	19(04,37)	3.96	Virginia	0.35	0.04	31(13,39)	-13.75
Massachusetts	0.41	0.04	23(06,37)	-1.30	Washington	0.39	0.05	24(08,38)	-5.08
Michigan	0.47	0.04	12(04,30)	13.84	West Virginia	0.71	0.10	02(01,20)	73.50 ^c
Minnesota	0.30	0.04	37(17,39)	-26.32 ^c	Wisconsin	0.30	0.04	36(18,39)	-25.74 ^c
Mississippi	0.47	0.07	13(02,38)	13.34	Wyoming	-	-	-(-,-)	-
Missouri	0.46	0.05	15(04,34)	10.94					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 13.26
Leukemia

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	6.96	0.02		
High Five States				
Wyoming	8.01	0.54	01(01,44)	15.07 ^c
Wisconsin	7.95 ^b	0.16	02(01,13)	14.17
West Virginia	7.82	0.26	03(01,25)	12.28
Oklahoma	7.77 ^b	0.20	04(01,21)	11.63
Minnesota	7.77 ^b	0.17	05(01,19)	11.55
Low Five States				
Vermont	6.34	0.42	47(09,51)	-8.96
Utah	6.23	0.25	48(24,51)	-10.54
New Mexico	6.17	0.24	49(28,51)	-11.42
D.C.	5.78	0.45	50(21,51)	-17.02 ^c
Hawaii	5.49 ^b	0.27	51(45,51)	-21.14 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	7.66 ^b	0.17	08(01,22)	10.05	Montana	6.75	0.34	34(05,50)	-3.08
Alaska	6.76	0.58	33(01,51)	-2.90	Nebraska	7.08	0.27	21(03,47)	1.62
Arizona	6.46 ^b	0.14	46(27,50)	-7.17	Nevada	6.57	0.23	41(18,50)	-5.61
Arkansas	7.51	0.22	10(01,30)	7.90	New Hampshire	6.88	0.31	29(05,50)	-1.20
California	6.55 ^b	0.06	43(31,48)	-5.86	New Jersey	6.58	0.12	39(26,49)	-5.46
Colorado	6.63	0.17	38(21,49)	-4.83	New Mexico	6.17	0.24	49(28,51)	-11.42
Connecticut	6.95	0.19	26(11,46)	-0.23	New York	6.58 ^b	0.08	40(29,48)	-5.57
Delaware	7.23	0.38	20(01,48)	3.87	North Carolina	6.79	0.12	31(21,45)	-2.49
D.C.	5.78	0.45	50(21,51)	-17.02 ^c	North Dakota	6.86	0.42	30(02,50)	-1.47
Florida	6.57 ^b	0.07	42(29,48)	-5.68	Ohio	7.29	0.11	19(08,27)	4.67
Georgia	6.48 ^b	0.13	45(28,49)	-6.93	Oklahoma	7.77 ^b	0.20	04(01,21)	11.63
Hawaii	5.49 ^b	0.27	51(45,51)	-21.14 ^c	Oregon	7.00	0.18	24(09,45)	0.55
Idaho	6.89	0.30	27(04,49)	-1.10	Pennsylvania	7.45 ^b	0.10	12(05,23)	7.00
Illinois	7.42 ^b	0.11	15(05,24)	6.58	Rhode Island	6.76	0.33	32(05,50)	-2.90
Indiana	7.76 ^b	0.15	06(01,18)	11.51	South Carolina	6.68	0.17	36(20,49)	-4.13
Iowa	7.04	0.20	23(08,45)	1.17	South Dakota	7.41	0.40	16(01,48)	6.38
Kansas	7.70	0.22	07(01,25)	10.53	Tennessee	7.50 ^b	0.15	11(02,25)	7.66
Kentucky	7.58	0.18	09(01,24)	8.91	Texas	6.88	0.08	28(21,38)	-1.18
Louisiana	7.43	0.18	14(02,28)	6.75	Utah	6.23	0.25	48(24,51)	-10.54
Maine	7.06	0.29	22(02,48)	1.38	Vermont	6.34	0.42	47(09,51)	-8.96
Maryland	6.53	0.15	44(24,49)	-6.23	Virginia	6.65	0.13	37(23,48)	-4.55
Massachusetts	6.72	0.14	35(21,47)	-3.48	Washington	6.98	0.15	25(13,41)	0.25
Michigan	7.35	0.12	17(05,26)	5.58	West Virginia	7.82	0.26	03(01,25)	12.28
Minnesota	7.77 ^b	0.17	05(01,19)	11.55	Wisconsin	7.95 ^b	0.16	02(01,13)	14.17
Mississippi	7.44	0.22	13(01,31)	6.88	Wyoming	8.01	0.54	01(01,44)	15.07 ^c
Missouri	7.31	0.15	18(04,29)	4.97					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 13.27
Leukemia

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	9.36	0.04		
High Five States				
Wisconsin	11.12 ^b	0.29	01(01,11)	18.87 ^c
Alabama	10.87 ^b	0.33	02(01,19)	16.13 ^c
Minnesota	10.72 ^b	0.30	03(01,19)	14.52
Wyoming	10.66	0.93	04(01,49)	13.92
Kentucky	10.66 ^b	0.34	05(01,23)	13.91
Low Five States				
California	8.51 ^b	0.11	47(37,50)	-9.04
Colorado	8.51	0.30	48(26,50)	-9.05
D.C.	8.26	0.87	49(03,51)	-11.78
New Mexico	7.89 ^b	0.41	50(31,51)	-15.70 ^c
Hawaii	7.00 ^b	0.45	51(46,51)	-25.15 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	10.87 ^b	0.33	02(01,19)	16.13 ^c	Montana	8.71	0.58	43(06,51)	-6.92
Alaska	9.76	1.05	19(01,51)	4.33	Nebraska	9.52	0.47	24(02,49)	1.75
Arizona	8.52 ^b	0.24	46(29,50)	-8.97	Nevada	8.84	0.41	41(13,50)	-5.53
Arkansas	10.33	0.39	08(01,33)	10.36	New Hampshire	8.96	0.55	37(05,50)	-4.25
California	8.51 ^b	0.11	47(37,50)	-9.04	New Jersey	8.95	0.21	38(21,48)	-4.34
Colorado	8.51	0.30	48(26,50)	-9.05	New Mexico	7.89 ^b	0.41	50(31,51)	-15.70 ^c
Connecticut	9.55	0.34	23(05,45)	2.01	New York	8.98	0.14	35(25,46)	-4.08
Delaware	10.01	0.69	14(01,49)	6.95	North Carolina	9.33	0.22	29(14,43)	-0.28
D.C.	8.26	0.87	49(03,51)	-11.78	North Dakota	9.46	0.74	26(01,50)	1.09
Florida	8.78 ^b	0.13	42(30,48)	-6.14	Ohio	9.93	0.19	18(06,29)	6.09
Georgia	8.94	0.24	39(21,49)	-4.47	Oklahoma	10.38	0.35	07(01,29)	10.92
Hawaii	7.00 ^b	0.45	51(46,51)	-25.15 ^c	Oregon	9.43	0.32	27(08,46)	0.77
Idaho	9.25	0.52	31(03,50)	-1.21	Pennsylvania	10.02 ^b	0.18	13(05,26)	7.12
Illinois	10.00	0.19	15(05,27)	6.81	Rhode Island	9.42	0.61	28(01,50)	0.61
Indiana	10.43 ^b	0.27	06(01,23)	11.49	South Carolina	9.18	0.31	33(13,48)	-1.95
Iowa	9.58	0.35	22(05,45)	2.40	South Dakota	9.74	0.70	20(01,50)	4.09
Kansas	10.16	0.40	12(01,36)	8.57	Tennessee	10.23	0.28	09(02,28)	9.36
Kentucky	10.66 ^b	0.34	05(01,23)	13.91	Texas	9.12	0.14	34(22,44)	-2.56
Louisiana	10.19	0.34	10(01,31)	8.90	Utah	8.55	0.44	45(16,51)	-8.60
Maine	9.51	0.52	25(02,50)	1.64	Vermont	8.96	0.77	36(01,51)	-4.23
Maryland	8.62	0.27	44(24,50)	-7.86	Virginia	9.19	0.24	32(16,46)	-1.76
Massachusetts	8.91	0.24	40(21,49)	-4.75	Washington	9.33	0.26	30(12,45)	-0.34
Michigan	9.65	0.21	21(09,36)	3.11	West Virginia	10.19	0.47	11(01,40)	8.87
Minnesota	10.72 ^b	0.30	03(01,19)	14.52	Wisconsin	11.12 ^b	0.29	01(01,11)	18.87 ^c
Mississippi	9.94	0.41	17(01,41)	6.21	Wyoming	10.66	0.93	04(01,49)	13.92
Missouri	9.97	0.27	16(03,32)	6.59					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 13.28
Leukemia

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	5.22	0.02		
High Five States				
West Virginia	6.09	0.31	01(01,29)	16.72 ^c
Wyoming	5.96	0.63	02(01,49)	14.30
Indiana	5.94 ^b	0.18	03(01,19)	13.78
Oklahoma	5.88	0.23	04(01,25)	12.62
Kansas	5.86	0.26	05(01,31)	12.22
Low Five States				
Utah	4.36	0.28	47(26,51)	-16.51 ^c
Vermont	4.34	0.47	48(06,51)	-16.83 ^c
Hawaii	4.28	0.33	49(20,51)	-17.91 ^c
D.C.	4.22	0.50	50(05,51)	-19.19 ^c
Alaska	4.12	0.61	51(02,51)	-21.05 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	5.48	0.19	17(02,38)	4.99	Montana	5.22	0.42	25(01,51)	-0.02
Alaska	4.12	0.61	51(02,51)	-21.05 ^c	Nebraska	5.27	0.31	22(01,48)	1.00
Arizona	4.85	0.16	41(19,49)	-7.03	Nevada	4.71	0.27	46(14,51)	-9.81
Arkansas	5.51	0.25	14(01,42)	5.54	New Hampshire	5.40	0.37	19(01,49)	3.42
California	5.08	0.07	32(21,41)	-2.73	New Jersey	4.95	0.13	36(19,48)	-5.17
Colorado	5.15	0.20	29(07,47)	-1.36	New Mexico	4.76	0.28	45(10,51)	-8.78
Connecticut	5.09	0.21	30(06,48)	-2.45	New York	4.89 ^b	0.09	39(27,47)	-6.37
Delaware	5.09	0.42	31(01,51)	-2.50	North Carolina	5.07	0.14	33(14,45)	-2.76
D.C.	4.22	0.50	50(05,51)	-19.19 ^c	North Dakota	4.86	0.47	40(01,51)	-6.90
Florida	4.83 ^b	0.09	42(29,48)	-7.36	Ohio	5.42	0.12	18(05,33)	3.83
Georgia	4.79	0.14	43(24,49)	-8.21	Oklahoma	5.88	0.23	04(01,25)	12.62
Hawaii	4.28	0.33	49(20,51)	-17.91 ^c	Oregon	5.19	0.21	27(05,46)	-0.45
Idaho	4.98	0.34	35(02,51)	-4.58	Pennsylvania	5.62 ^b	0.11	09(02,24)	7.68
Illinois	5.66 ^b	0.12	08(02,24)	8.48	Rhode Island	4.92	0.38	37(02,51)	-5.80
Indiana	5.94 ^b	0.18	03(01,19)	13.78	South Carolina	4.91	0.19	38(15,49)	-5.94
Iowa	5.23	0.23	24(04,47)	0.22	South Dakota	5.52	0.46	13(01,49)	5.72
Kansas	5.86	0.26	05(01,31)	12.22	Tennessee	5.53	0.17	12(02,34)	5.92
Kentucky	5.48	0.21	16(02,40)	5.09	Texas	5.24	0.09	23(13,37)	0.44
Louisiana	5.49	0.21	15(01,40)	5.21	Utah	4.36	0.28	47(26,51)	-16.51 ^c
Maine	5.16	0.34	28(01,50)	-1.07	Vermont	4.34	0.47	48(06,51)	-16.83 ^c
Maryland	5.05	0.18	34(11,48)	-3.18	Virginia	4.77	0.15	44(24,49)	-8.60
Massachusetts	5.28	0.16	21(06,41)	1.12	Washington	5.21	0.17	26(07,44)	-0.14
Michigan	5.67	0.14	07(02,25)	8.58	West Virginia	6.09	0.31	01(01,29)	16.72 ^c
Minnesota	5.58	0.19	10(01,34)	7.00	Wisconsin	5.58	0.18	11(01,33)	6.85
Mississippi	5.67	0.26	06(01,37)	8.64	Wyoming	5.96	0.63	02(01,49)	14.30
Missouri	5.36	0.17	20(04,40)	2.66					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 14.18
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	5.96	0.02		
High Five States				
D.C.	9.17 ^b	0.56	01(01,04)	53.83 ^c
Texas	7.71 ^b	0.08	02(01,04)	29.35 ^c
Louisiana	7.69 ^b	0.18	03(01,06)	29.00 ^c
Hawaii	7.58 ^b	0.31	04(01,08)	27.15 ^c
California	7.24 ^b	0.06	05(03,07)	21.58 ^c
Low Five States				
Nebraska	4.55 ^b	0.21	47(36,51)	-23.58 ^c
Idaho	4.33 ^b	0.23	48(38,51)	-27.32 ^c
Utah	4.19 ^b	0.20	49(42,51)	-29.63 ^c
Montana	4.17 ^b	0.26	50(40,51)	-30.04 ^c
North Dakota	3.78 ^b	0.31	51(44,51)	-36.61 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	6.14	0.15	14(07,22)	3.08	Montana	4.17 ^b	0.26	50(40,51)	-30.04 ^c
Alaska	5.69	0.47	22(06,48)	-4.57	Nebraska	4.55 ^b	0.21	47(36,51)	-23.58 ^c
Arizona	5.72	0.13	21(13,34)	-4.00	Nevada	5.48	0.20	29(14,42)	-8.04
Arkansas	6.34	0.19	09(06,21)	6.45	New Hampshire	4.63 ^b	0.25	45(32,51)	-22.29 ^c
California	7.24 ^b	0.06	05(03,07)	21.58 ^c	New Jersey	5.50 ^b	0.11	27(19,37)	-7.66
Colorado	5.16 ^b	0.15	37(24,44)	-13.37	New Mexico	6.68	0.25	07(04,17)	12.14
Connecticut	5.45	0.16	33(16,41)	-8.53	New York	5.93	0.07	17(12,23)	-0.50
Delaware	6.28	0.35	10(04,34)	5.47	North Carolina	5.47 ^b	0.10	32(19,38)	-8.27
D.C.	9.17 ^b	0.56	01(01,04)	53.83 ^c	North Dakota	3.78 ^b	0.31	51(44,51)	-36.61 ^c
Florida	5.66 ^b	0.07	23(17,31)	-5.04	Ohio	5.27 ^b	0.09	35(26,41)	-11.50
Georgia	5.47 ^b	0.11	31(19,38)	-8.19	Oklahoma	5.78	0.17	20(10,34)	-2.95
Hawaii	7.58 ^b	0.31	04(01,08)	27.15 ^c	Oregon	6.05	0.16	16(08,27)	1.48
Idaho	4.33 ^b	0.23	48(38,51)	-27.32 ^c	Pennsylvania	5.53 ^b	0.08	25(19,35)	-7.26
Illinois	5.65	0.09	24(16,33)	-5.16	Rhode Island	6.18	0.32	13(06,35)	3.67
Indiana	5.10 ^b	0.12	38(29,44)	-14.39	South Carolina	5.81	0.15	19(11,33)	-2.52
Iowa	4.71 ^b	0.16	44(36,49)	-21.03 ^c	South Dakota	4.62 ^b	0.31	46(25,51)	-22.49 ^c
Kansas	4.81 ^b	0.17	43(33,49)	-19.34 ^c	Tennessee	6.19	0.13	12(07,20)	3.97
Kentucky	5.52	0.15	26(16,39)	-7.38	Texas	7.71 ^b	0.08	02(01,04)	29.35 ^c
Louisiana	7.69 ^b	0.18	03(01,06)	29.00 ^c	Utah	4.19 ^b	0.20	49(42,51)	-29.63 ^c
Maine	4.86 ^b	0.24	41(26,50)	-18.49 ^c	Vermont	5.32	0.38	34(08,49)	-10.68
Maryland	6.06	0.14	15(08,25)	1.69	Virginia	5.49 ^b	0.12	28(19,38)	-7.84
Massachusetts	6.24	0.13	11(07,19)	4.65	Washington	6.40	0.14	08(06,16)	7.40
Michigan	5.47 ^b	0.10	30(20,37)	-8.19	West Virginia	4.84 ^b	0.20	42(29,49)	-18.80 ^c
Minnesota	5.02 ^b	0.13	39(31,46)	-15.79 ^c	Wisconsin	5.17 ^b	0.13	36(26,44)	-13.22
Mississippi	7.04 ^b	0.21	06(03,09)	18.22 ^c	Wyoming	4.94	0.41	40(12,51)	-17.05 ^c
Missouri	5.82	0.13	18(11,31)	-2.26					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 14.19
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	8.83	0.03		
High Five States				
D.C.	14.71 ^b	1.08	01(01,02)	66.48 ^c
Louisiana	11.71 ^b	0.34	02(01,05)	32.54 ^c
Texas	11.29 ^b	0.15	03(02,05)	27.87 ^c
Hawaii	11.06 ^b	0.55	04(02,12)	25.26 ^c
California	10.55 ^b	0.11	05(04,08)	19.45 ^c
Low Five States				
Idaho	6.37 ^b	0.41	47(36,51)	-27.92 ^c
South Dakota	5.85 ^b	0.52	48(36,51)	-33.75 ^c
Montana	5.62 ^b	0.44	49(41,51)	-36.35 ^c
Utah	5.53 ^b	0.33	50(45,51)	-37.40 ^c
North Dakota	5.40 ^b	0.55	51(40,51)	-38.82 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	9.06	0.28	15(07,27)	2.61	Montana	5.62 ^b	0.44	49(41,51)	-36.35 ^c
Alaska	8.15	0.84	31(04,49)	-7.70	Nebraska	6.62 ^b	0.38	46(34,50)	-25.05 ^c
Arizona	8.12	0.22	32(18,38)	-8.09	Nevada	7.83	0.36	35(16,44)	-11.36
Arkansas	9.48	0.36	11(05,24)	7.35	New Hampshire	6.96 ^b	0.45	44(22,49)	-21.15 ^c
California	10.55 ^b	0.11	05(04,08)	19.45 ^c	New Jersey	8.27	0.20	26(17,36)	-6.33
Colorado	7.61 ^b	0.27	36(23,44)	-13.82	New Mexico	9.67	0.43	10(04,27)	9.45
Connecticut	8.52	0.31	21(11,37)	-3.57	New York	8.90	0.14	17(11,23)	0.80
Delaware	9.82	0.65	08(02,34)	11.13	North Carolina	8.28	0.19	24(17,36)	-6.28
D.C.	14.71 ^b	1.08	01(01,02)	66.48 ^c	North Dakota	5.40 ^b	0.55	51(40,51)	-38.82 ^c
Florida	8.54	0.12	20(16,30)	-3.36	Ohio	7.98 ^b	0.16	34(22,38)	-9.64
Georgia	8.24	0.21	27(17,37)	-6.70	Oklahoma	8.22	0.30	29(14,39)	-6.97
Hawaii	11.06 ^b	0.55	04(02,12)	25.26 ^c	Oregon	9.00	0.30	16(08,30)	1.85
Idaho	6.37 ^b	0.41	47(36,51)	-27.92 ^c	Pennsylvania	8.32	0.15	23(18,35)	-5.78
Illinois	8.28	0.17	25(17,36)	-6.28	Rhode Island	9.72	0.60	09(02,32)	10.08
Indiana	7.32 ^b	0.22	38(32,45)	-17.12 ^c	South Carolina	8.84	0.28	18(09,32)	0.06
Iowa	6.72 ^b	0.29	45(36,49)	-23.94 ^c	South Dakota	5.85 ^b	0.52	48(36,51)	-33.75 ^c
Kansas	6.98 ^b	0.31	42(32,48)	-21.03 ^c	Tennessee	9.46	0.25	12(06,20)	7.14
Kentucky	8.21	0.28	30(15,38)	-7.06	Texas	11.29 ^b	0.15	03(02,05)	27.87 ^c
Louisiana	11.71 ^b	0.34	02(01,05)	32.54 ^c	Utah	5.53 ^b	0.33	50(45,51)	-37.40 ^c
Maine	7.16 ^b	0.44	40(21,49)	-18.99 ^c	Vermont	8.22	0.70	28(06,47)	-6.90
Maryland	9.29	0.27	14(07,23)	5.20	Virginia	8.36	0.22	22(15,36)	-5.37
Massachusetts	9.85 ^b	0.25	07(05,15)	11.53	Washington	9.34	0.24	13(07,21)	5.71
Michigan	7.99 ^b	0.18	33(21,38)	-9.51	West Virginia	6.99 ^b	0.37	41(29,49)	-20.92 ^c
Minnesota	7.19 ^b	0.24	39(32,47)	-18.62 ^c	Wisconsin	7.45 ^b	0.23	37(28,45)	-15.67 ^c
Mississippi	10.43 ^b	0.40	06(02,14)	18.09 ^c	Wyoming	6.97	0.71	43(13,51)	-21.10 ^c
Missouri	8.69	0.24	19(11,32)	-1.61					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 14.20
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	3.53	0.02		
High Five States				
D.C.	4.81	0.54	01(01,29)	36.01 ^c
Texas	4.60 ^b	0.09	02(01,06)	30.31 ^c
Hawaii	4.41	0.32	03(01,17)	24.80 ^c
California	4.40 ^b	0.07	04(01,07)	24.68 ^c
Louisiana	4.40 ^b	0.19	05(01,09)	24.60 ^c
Low Five States				
Vermont	2.80	0.38	47(07,51)	-20.79 ^c
Montana	2.78	0.30	48(10,51)	-21.22 ^c
New Hampshire	2.55 ^b	0.25	49(24,51)	-27.77 ^c
Idaho	2.52 ^b	0.24	50(27,51)	-28.65 ^c
North Dakota	2.32 ^b	0.32	51(22,51)	-34.42 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	3.77	0.16	09(05,27)	6.75	Montana	2.78	0.30	48(10,51)	-21.22 ^c
Alaska	3.34	0.52	25(01,51)	-5.59	Nebraska	2.84	0.23	46(15,51)	-19.68 ^c
Arizona	3.58	0.14	12(07,33)	1.30	Nevada	3.35	0.22	23(07,48)	-5.18
Arkansas	3.67	0.20	11(05,38)	3.96	New Hampshire	2.55 ^b	0.25	49(24,51)	-27.77 ^c
California	4.40 ^b	0.07	04(01,07)	24.68 ^c	New Jersey	3.30	0.11	27(11,43)	-6.56
Colorado	3.01	0.15	40(16,50)	-14.69	New Mexico	4.02	0.26	07(01,28)	13.90
Connecticut	2.96 ^b	0.16	43(18,50)	-16.18 ^c	New York	3.51	0.08	14(09,28)	-0.62
Delaware	3.35	0.34	22(03,51)	-5.16	North Carolina	3.15 ^b	0.11	35(17,46)	-10.85
D.C.	4.81	0.54	01(01,29)	36.01 ^c	North Dakota	2.32 ^b	0.32	51(22,51)	-34.42 ^c
Florida	3.17 ^b	0.07	33(21,43)	-10.18	Ohio	3.06 ^b	0.09	38(22,47)	-13.31
Georgia	3.25	0.11	29(13,45)	-8.06	Oklahoma	3.68	0.18	10(06,34)	4.24
Hawaii	4.41	0.32	03(01,17)	24.80 ^c	Oregon	3.41	0.17	19(07,44)	-3.53
Idaho	2.52 ^b	0.24	50(27,51)	-28.65 ^c	Pennsylvania	3.20 ^b	0.09	32(18,44)	-9.51
Illinois	3.52	0.10	13(08,31)	-0.34	Rhode Island	3.43	0.32	18(04,49)	-2.87
Indiana	3.27	0.13	28(11,45)	-7.36	South Carolina	3.31	0.15	26(09,45)	-6.45
Iowa	2.95	0.17	44(17,50)	-16.46 ^c	South Dakota	3.46	0.37	16(02,50)	-2.13
Kansas	2.91	0.19	45(16,51)	-17.55 ^c	Tennessee	3.51	0.13	15(08,37)	-0.64
Kentucky	3.24	0.16	30(10,47)	-8.39	Texas	4.60 ^b	0.09	02(01,06)	30.31 ^c
Louisiana	4.40 ^b	0.19	05(01,09)	24.60 ^c	Utah	2.97	0.23	42(11,51)	-15.93 ^c
Maine	3.01	0.26	41(09,51)	-14.78	Vermont	2.80	0.38	47(07,51)	-20.79 ^c
Maryland	3.45	0.14	17(08,40)	-2.30	Virginia	3.14	0.12	36(16,47)	-11.14
Massachusetts	3.35	0.13	24(10,43)	-5.24	Washington	3.79	0.14	08(05,23)	7.33
Michigan	3.36	0.10	21(11,40)	-5.02	West Virginia	3.06	0.22	39(10,50)	-13.43
Minnesota	3.13	0.14	37(14,48)	-11.42	Wisconsin	3.23	0.14	31(12,46)	-8.55
Mississippi	4.30	0.22	06(01,13)	21.66 ^c	Wyoming	3.16	0.46	34(02,51)	-10.65
Missouri	3.38	0.13	20(09,42)	-4.25					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 15.24
Cancer of the Lung and Bronchus (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	47.22	0.05		
High Five States				
Kentucky	70.92 ^b	0.55	01(01,01)	50.18 ^c
Arkansas	63.26 ^b	0.62	02(02,04)	33.95 ^c
West Virginia	62.70 ^b	0.73	03(02,05)	32.77 ^c
Tennessee	61.39 ^b	0.42	04(03,05)	30.00 ^c
Mississippi	61.32 ^b	0.63	05(02,05)	29.86 ^c
Low Five States				
California	36.15 ^b	0.14	47(47,47)	-23.44 ^c
Colorado	34.45 ^b	0.39	48(48,50)	-27.05 ^c
Hawaii	33.46 ^b	0.66	49(48,50)	-29.14 ^c
New Mexico	33.27 ^b	0.55	50(48,50)	-29.54 ^c
Utah	20.41 ^b	0.45	51(51,51)	-56.79 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	58.14 ^b	0.47	07(06,10)	23.12 ^c	Montana	42.77 ^b	0.86	37(34,44)	-9.42 ^c
Alaska	52.07	1.57	17(11,26)	10.25 ^c	Nebraska	44.18 ^b	0.67	35(32,39)	-6.45 ^c
Arizona	38.47 ^b	0.33	46(44,46)	-18.53 ^c	Nevada	49.37	0.63	22(17,27)	4.54
Arkansas	63.26 ^b	0.62	02(02,04)	33.95 ^c	New Hampshire	48.65	0.82	24(17,31)	3.02
California	36.15 ^b	0.14	47(47,47)	-23.44 ^c	New Jersey	42.14 ^b	0.30	39(37,43)	-10.77 ^c
Colorado	34.45 ^b	0.39	48(48,50)	-27.05 ^c	New Mexico	33.27 ^b	0.55	50(48,50)	-29.54 ^c
Connecticut	41.58 ^b	0.45	41(37,44)	-11.95 ^c	New York	42.03 ^b	0.20	40(37,43)	-11.00 ^c
Delaware	54.20 ^b	1.03	12(10,17)	14.78 ^c	North Carolina	53.03 ^b	0.33	15(12,17)	12.30 ^c
D.C.	42.51 ^b	1.23	38(32,45)	-9.98 ^c	North Dakota	40.84 ^b	1.03	43(36,46)	-13.52 ^c
Florida	46.58	0.19	30(28,33)	-1.36	Ohio	55.32 ^b	0.29	11(11,13)	17.14 ^c
Georgia	50.06 ^b	0.34	18(17,23)	6.01 ^c	Oklahoma	58.42 ^b	0.54	06(06,10)	23.71 ^c
Hawaii	33.46 ^b	0.66	49(48,50)	-29.14 ^c	Oregon	47.23	0.47	28(24,32)	0.02
Idaho	39.15 ^b	0.71	45(42,46)	-17.10 ^c	Pennsylvania	48.63 ^b	0.25	25(21,27)	2.98
Illinois	49.74 ^b	0.28	20(17,24)	5.32 ^c	Rhode Island	49.99	0.91	19(16,27)	5.86 ^c
Indiana	57.75 ^b	0.41	08(06,10)	22.30 ^c	South Carolina	53.32 ^b	0.46	14(12,17)	12.91 ^c
Iowa	47.75	0.52	27(22,31)	1.11	South Dakota	44.97	0.98	34(27,39)	-4.77
Kansas	48.92	0.57	23(18,28)	3.59	Tennessee	61.39 ^b	0.42	04(03,05)	30.00 ^c
Kentucky	70.92 ^b	0.55	01(01,01)	50.18 ^c	Texas	43.53 ^b	0.20	36(34,38)	-7.83 ^c
Louisiana	57.64 ^b	0.50	09(06,10)	22.06 ^c	Utah	20.41 ^b	0.45	51(51,51)	-56.79 ^c
Maine	53.77 ^b	0.80	13(11,17)	13.87 ^c	Vermont	49.53	1.16	21(15,31)	4.88
Maryland	46.41	0.40	31(27,34)	-1.72	Virginia	48.55 ^b	0.35	26(20,28)	2.81
Massachusetts	46.80	0.36	29(26,33)	-0.90	Washington	45.77 ^b	0.37	33(29,35)	-3.08
Michigan	52.17 ^b	0.31	16(14,18)	10.47 ^c	West Virginia	62.70 ^b	0.73	03(02,05)	32.77 ^c
Minnesota	41.56 ^b	0.39	42(37,44)	-11.98 ^c	Wisconsin	45.83 ^b	0.38	32(29,35)	-2.95
Mississippi	61.32 ^b	0.63	05(02,05)	29.86 ^c	Wyoming	39.62 ^b	1.19	44(37,46)	-16.10 ^c
Missouri	56.95 ^b	0.41	10(07,11)	20.60 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 5% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 15.25
Cancer of the Lung and Bronchus (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	59.79	0.09		
High Five States				
Kentucky	92.19 ^b	0.96	01(01,02)	54.19 ^c
Mississippi	88.45 ^b	1.18	02(01,03)	47.94 ^c
Arkansas	88.00 ^b	1.10	03(01,03)	47.18 ^c
Tennessee	83.57 ^b	0.76	04(04,06)	39.77 ^c
Alabama	82.56 ^b	0.86	05(04,06)	38.09 ^c
Low Five States				
Hawaii	44.37 ^b	1.14	47(44,49)	-25.78 ^c
California	43.67 ^b	0.24	48(46,48)	-26.96 ^c
New Mexico	40.90 ^b	0.93	49(48,50)	-31.59 ^c
Colorado	40.75 ^b	0.65	50(48,50)	-31.84 ^c
Utah	26.35 ^b	0.76	51(51,51)	-55.93 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	82.56 ^b	0.86	05(04,06)	38.09 ^c	Montana	49.66 ^b	1.37	42(37,46)	-16.94 ^c
Alaska	61.25	2.59	24(14,37)	2.44	Nebraska	55.46 ^b	1.13	34(25,38)	-7.24 ^c
Arizona	46.63 ^b	0.55	44(43,47)	-22.00 ^c	Nevada	55.94 ^b	0.99	33(25,38)	-6.44 ^c
Arkansas	88.00 ^b	1.10	03(01,03)	47.18 ^c	New Hampshire	57.71	1.35	27(23,37)	-3.47
California	43.67 ^b	0.24	48(46,48)	-26.96 ^c	New Jersey	52.05 ^b	0.51	39(37,41)	-12.95 ^c
Colorado	40.75 ^b	0.65	50(48,50)	-31.84 ^c	New Mexico	40.90 ^b	0.93	49(48,50)	-31.59 ^c
Connecticut	49.73 ^b	0.76	41(39,44)	-16.83 ^c	New York	51.94 ^b	0.34	40(37,41)	-13.13 ^c
Delaware	66.37 ^b	1.73	16(13,23)	11.01 ^c	North Carolina	72.03 ^b	0.59	12(10,13)	20.48 ^c
D.C.	54.60	2.19	36(23,43)	-8.68 ^c	North Dakota	53.31 ^b	1.76	38(25,43)	-10.84 ^c
Florida	58.11 ^b	0.32	26(24,30)	-2.82	Ohio	70.92 ^b	0.50	13(11,14)	18.62 ^c
Georgia	68.00 ^b	0.63	14(14,16)	13.73 ^c	Oklahoma	75.70 ^b	0.93	08(07,10)	26.61 ^c
Hawaii	44.37 ^b	1.14	47(44,49)	-25.78 ^c	Oregon	55.15 ^b	0.76	35(27,38)	-7.75 ^c
Idaho	45.86 ^b	1.14	46(43,48)	-23.30 ^c	Pennsylvania	62.44 ^b	0.43	20(17,24)	4.44
Illinois	62.40 ^b	0.47	21(17,24)	4.37	Rhode Island	62.52	1.56	18(15,26)	4.56
Indiana	75.33 ^b	0.72	09(07,10)	25.99 ^c	South Carolina	72.72 ^b	0.83	11(09,13)	21.62 ^c
Iowa	61.40	0.88	23(18,26)	2.69	South Dakota	58.16	1.67	25(18,37)	-2.72
Kansas	61.68	0.96	22(17,26)	3.16	Tennessee	83.57 ^b	0.76	04(04,06)	39.77 ^c
Kentucky	92.19 ^b	0.96	01(01,02)	54.19 ^c	Texas	56.44 ^b	0.35	30(26,35)	-5.60 ^c
Louisiana	77.40 ^b	0.90	07(06,09)	29.45 ^c	Utah	26.35 ^b	0.76	51(51,51)	-55.93 ^c
Maine	66.50 ^b	1.34	15(14,19)	11.23 ^c	Vermont	57.23	1.89	28(19,39)	-4.27
Maryland	57.07 ^b	0.69	29(24,35)	-4.54	Virginia	62.48 ^b	0.61	19(17,24)	4.51
Massachusetts	56.18 ^b	0.60	32(26,37)	-6.03 ^c	Washington	53.85 ^b	0.61	37(33,39)	-9.93 ^c
Michigan	64.69 ^b	0.52	17(15,19)	8.20 ^c	West Virginia	80.46 ^b	1.26	06(04,07)	34.57 ^c
Minnesota	49.66 ^b	0.64	43(40,44)	-16.94 ^c	Wisconsin	56.30 ^b	0.64	31(26,36)	-5.83 ^c
Mississippi	88.45 ^b	1.18	02(01,03)	47.94 ^c	Wyoming	46.47 ^b	1.91	45(39,49)	-22.28 ^c
Missouri	72.76 ^b	0.71	10(09,13)	21.70 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 5% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 15.26
Cancer of the Lung And Bronchus (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	37.80	0.06		
High Five States				
Kentucky	55.23 ^b	0.65	01(01,01)	46.11 ^c
West Virginia	49.26 ^b	0.88	02(02,03)	30.32 ^c
Oklahoma	45.31 ^b	0.64	03(03,13)	19.86 ^c
Missouri	45.23 ^b	0.49	04(03,12)	19.65 ^c
Delaware	45.18 ^b	1.26	05(02,18)	19.53 ^c
Low Five States				
California	30.51 ^b	0.18	47(45,48)	-19.29 ^c
Colorado	29.72 ^b	0.49	48(46,49)	-21.38 ^c
New Mexico	27.21 ^b	0.68	49(48,50)	-28.01 ^c
Hawaii	24.98 ^b	0.77	50(49,50)	-33.92 ^c
Utah	15.56 ^b	0.53	51(51,51)	-58.83 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	40.13 ^b	0.52	22(17,28)	6.18	Montana	37.19	1.09	34(21,42)	-1.61
Alaska	44.81	1.98	07(02,25)	18.55 ^c	Nebraska	35.53	0.81	37(30,44)	-5.99
Arizona	31.89 ^b	0.41	45(43,46)	-15.64 ^c	Nevada	44.11 ^b	0.81	10(03,17)	16.69 ^c
Arkansas	44.21 ^b	0.70	09(03,17)	16.96 ^c	New Hampshire	42.14 ^b	1.03	16(05,25)	11.49 ^c
California	30.51 ^b	0.18	47(45,48)	-19.29 ^c	New Jersey	35.26 ^b	0.36	38(34,42)	-6.72
Colorado	29.72 ^b	0.49	48(46,49)	-21.38 ^c	New Mexico	27.21 ^b	0.68	49(48,50)	-28.01 ^c
Connecticut	35.82 ^b	0.56	35(32,42)	-5.25	New York	35.11 ^b	0.24	39(35,42)	-7.12
Delaware	45.18 ^b	1.26	05(02,18)	19.53 ^c	North Carolina	39.15 ^b	0.37	24(21,31)	3.57
D.C.	33.91	1.44	41(29,48)	-10.29 ^c	North Dakota	31.77 ^b	1.24	46(37,48)	-15.94 ^c
Florida	37.26	0.23	32(29,35)	-1.42	Ohio	43.80 ^b	0.35	12(06,15)	15.87 ^c
Georgia	37.23	0.39	33(28,36)	-1.52	Oklahoma	45.31 ^b	0.64	03(03,13)	19.86 ^c
Hawaii	24.98 ^b	0.77	50(49,50)	-33.92 ^c	Oregon	41.18 ^b	0.60	19(14,24)	8.94
Idaho	33.63 ^b	0.90	44(35,46)	-11.04 ^c	Pennsylvania	38.72	0.30	27(23,31)	2.43
Illinois	40.72 ^b	0.33	20(16,23)	7.73	Rhode Island	41.58	1.11	17(06,29)	10.00 ^c
Indiana	44.72 ^b	0.49	08(03,13)	18.29 ^c	South Carolina	38.93	0.53	26(20,32)	2.99
Iowa	37.49	0.62	31(24,37)	-0.82	South Dakota	35.04	1.18	40(28,46)	-7.31
Kansas	39.13	0.68	25(18,33)	3.51	Tennessee	45.11 ^b	0.48	06(03,12)	19.33 ^c
Kentucky	55.23 ^b	0.65	01(01,01)	46.11 ^c	Texas	33.71 ^b	0.24	42(40,45)	-10.81 ^c
Louisiana	42.84 ^b	0.58	14(08,19)	13.34 ^c	Utah	15.56 ^b	0.53	51(51,51)	-58.83 ^c
Maine	44.11 ^b	0.98	11(03,18)	16.68 ^c	Vermont	43.79 ^b	1.48	13(03,24)	15.85 ^c
Maryland	38.63	0.49	28(21,33)	2.20	Virginia	38.19	0.41	29(24,34)	1.03
Massachusetts	40.31 ^b	0.44	21(17,26)	6.63	Washington	39.70 ^b	0.47	23(18,30)	5.02
Michigan	42.84 ^b	0.38	15(10,18)	13.34 ^c	West Virginia	49.26 ^b	0.88	02(02,03)	30.32 ^c
Minnesota	35.60 ^b	0.49	36(33,42)	-5.81	Wisconsin	38.13	0.47	30(23,35)	0.86
Mississippi	41.40 ^b	0.69	18(12,24)	9.52	Wyoming	33.68	1.50	43(30,48)	-10.89 ^c
Missouri	45.23 ^b	0.49	04(03,12)	19.65 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 5% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 16.18
Melanoma of the Skin (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	2.72	0.01		
High Five States				
Utah	3.62 ^b	0.18	01(01,12)	33.27 ^c
Idaho	3.45	0.21	02(01,26)	26.85 ^c
Oregon	3.35 ^b	0.13	03(01,19)	23.11 ^c
West Virginia	3.32	0.17	04(01,28)	22.13 ^c
Colorado	3.27 ^b	0.12	05(01,23)	20.45 ^c
Low Five States				
New York	2.23 ^b	0.05	47(42,49)	-18.01 ^c
Alaska	2.17	0.31	48(07,50)	-20.34 ^c
North Dakota	2.14	0.24	49(24,50)	-21.40 ^c
Hawaii	1.48 ^b	0.14	50(49,51)	-45.55 ^c
D.C.	1.11 ^b	0.20	51(50,51)	-59.06 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	2.87	0.11	27(08,36)	5.76	Montana	3.02	0.23	14(01,43)	11.24
Alaska	2.17	0.31	48(07,50)	-20.34 ^c	Nebraska	3.15	0.18	08(01,34)	15.78 ^c
Arizona	2.94	0.09	21(06,33)	8.27	Nevada	3.05	0.16	13(02,35)	12.24
Arkansas	2.83	0.13	29(06,41)	4.21	New Hampshire	2.84	0.20	28(02,45)	4.53
California	2.57 ^b	0.04	35(30,41)	-5.51	New Jersey	2.48	0.07	36(30,47)	-8.75
Colorado	3.27 ^b	0.12	05(01,23)	20.45 ^c	New Mexico	2.71	0.16	32(08,47)	-0.44
Connecticut	2.45	0.11	38(28,49)	-9.70	New York	2.23 ^b	0.05	47(42,49)	-18.01 ^c
Delaware	2.83	0.24	30(02,48)	4.01	North Carolina	2.91	0.08	24(09,33)	7.07
D.C.	1.11 ^b	0.20	51(50,51)	-59.06 ^c	North Dakota	2.14	0.24	49(24,50)	-21.40 ^c
Florida	2.94 ^b	0.05	20(10,30)	8.31	Ohio	2.95 ^b	0.07	19(08,31)	8.66
Georgia	2.43 ^b	0.08	40(31,48)	-10.43	Oklahoma	3.21 ^b	0.13	07(01,27)	18.13 ^c
Hawaii	1.48 ^b	0.14	50(49,51)	-45.55 ^c	Oregon	3.35 ^b	0.13	03(01,19)	23.11 ^c
Idaho	3.45	0.21	02(01,26)	26.85 ^c	Pennsylvania	2.92	0.06	23(10,31)	7.49
Illinois	2.44 ^b	0.06	39(32,47)	-10.30	Rhode Island	2.43	0.20	42(13,49)	-10.72
Indiana	3.06 ^b	0.09	12(03,30)	12.43	South Carolina	2.66	0.10	33(15,44)	-1.99
Iowa	2.94	0.13	22(04,36)	8.12	South Dakota	2.32	0.23	44(13,49)	-14.78
Kansas	2.99	0.14	16(02,36)	10.18	Tennessee	3.08 ^b	0.10	11(03,29)	13.43
Kentucky	3.22 ^b	0.12	06(01,26)	18.46 ^c	Texas	2.41 ^b	0.05	43(34,47)	-11.16
Louisiana	2.28 ^b	0.10	45(34,49)	-16.21 ^c	Utah	3.62 ^b	0.18	01(01,12)	33.27 ^c
Maine	2.98	0.19	17(02,42)	9.73	Vermont	3.01	0.29	15(01,47)	10.91
Maryland	2.66	0.10	34(19,44)	-2.23	Virginia	2.90	0.09	25(08,34)	6.80
Massachusetts	2.90	0.09	26(07,34)	6.79	Washington	2.96	0.09	18(06,33)	8.74
Michigan	2.43 ^b	0.07	41(32,47)	-10.60	West Virginia	3.32	0.17	04(01,28)	22.13 ^c
Minnesota	2.48	0.09	37(29,48)	-8.83	Wisconsin	2.73	0.09	31(14,40)	0.29
Mississippi	2.27 ^b	0.12	46(33,49)	-16.39 ^c	Wyoming	3.11	0.33	09(01,48)	14.57
Missouri	3.09 ^b	0.10	10(03,29)	13.53					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 16.19
Melanoma of the Skin (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	4.07	0.02		
High Five States				
Utah	5.62 ^b	0.35	01(01,11)	38.17 ^c
West Virginia	5.04	0.32	02(01,29)	23.85 ^c
Tennessee	4.95 ^b	0.19	03(01,20)	21.74 ^c
Idaho	4.89	0.37	04(01,38)	20.24 ^c
Oklahoma	4.87	0.24	05(01,27)	19.72 ^c
Low Five States				
North Dakota	3.25	0.44	46(08,50)	-20.10 ^c
New York	3.22 ^b	0.08	47(42,49)	-20.80 ^c
South Dakota	2.90	0.38	48(29,50)	-28.55 ^c
Alaska	2.76	0.55	49(12,50)	-32.16 ^c
Hawaii	2.33 ^b	0.26	50(47,50)	-42.67 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	4.47	0.20	13(02,35)	9.82	Montana	4.05	0.39	31(02,48)	-0.33
Alaska	2.76	0.55	49(12,50)	-32.16 ^c	Nebraska	4.27	0.31	24(02,45)	5.08
Arizona	4.41	0.17	18(04,34)	8.41	Nevada	4.67	0.29	08(01,36)	14.89
Arkansas	4.48	0.25	12(02,38)	10.21	New Hampshire	4.15	0.37	29(02,47)	1.99
California	3.86	0.07	35(28,41)	-5.13	New Jersey	3.66	0.13	40(29,47)	-9.92
Colorado	4.72	0.22	07(01,31)	16.05 ^c	New Mexico	3.96	0.29	34(06,47)	-2.53
Connecticut	3.40	0.20	45(31,49)	-16.33 ^c	New York	3.22 ^b	0.08	47(42,49)	-20.80 ^c
Delaware	4.02	0.43	32(02,49)	-1.16	North Carolina	4.44	0.15	15(04,32)	9.14
D.C.	-	-	-(-,-)	-	North Dakota	3.25	0.44	46(08,50)	-20.10 ^c
Florida	4.41 ^b	0.09	17(07,30)	8.49	Ohio	4.35	0.13	21(07,33)	6.94
Georgia	3.85	0.15	37(21,45)	-5.42	Oklahoma	4.87	0.24	05(01,27)	19.72 ^c
Hawaii	2.33 ^b	0.26	50(47,50)	-42.67 ^c	Oregon	4.53	0.22	10(02,34)	11.38
Idaho	4.89	0.37	04(01,38)	20.24 ^c	Pennsylvania	4.30	0.11	23(09,33)	5.78
Illinois	3.62 ^b	0.11	42(31,47)	-10.98	Rhode Island	3.85	0.38	36(03,49)	-5.37
Indiana	4.50	0.17	11(03,32)	10.69	South Carolina	4.17	0.20	28(06,42)	2.64
Iowa	4.38	0.24	19(02,39)	7.70	South Dakota	2.90	0.38	48(29,50)	-28.55 ^c
Kansas	4.46	0.26	14(02,39)	9.63	Tennessee	4.95 ^b	0.19	03(01,20)	21.74 ^c
Kentucky	4.60	0.22	09(01,33)	13.11	Texas	3.70 ^b	0.09	39(30,45)	-9.00
Louisiana	3.47	0.19	44(28,49)	-14.56	Utah	5.62 ^b	0.35	01(01,11)	38.17 ^c
Maine	4.44	0.35	16(01,44)	9.10	Vermont	4.27	0.53	25(01,49)	5.05
Maryland	4.21	0.19	26(05,39)	3.66	Virginia	4.35	0.16	20(05,35)	7.10
Massachusetts	4.20	0.16	27(07,39)	3.29	Washington	4.31	0.17	22(05,36)	5.96
Michigan	3.59 ^b	0.12	43(31,47)	-11.58	West Virginia	5.04	0.32	02(01,29)	23.85 ^c
Minnesota	3.72	0.17	38(22,47)	-8.63	Wisconsin	4.11	0.17	30(09,41)	1.11
Mississippi	3.62	0.24	41(17,49)	-10.92	Wyoming	4.01	0.56	33(01,50)	-1.36
Missouri	4.78 ^b	0.18	06(01,26)	17.54 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 16.20
Melanoma of the Skin (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	1.70	0.01		
High Five States				
Oregon	2.42 ^b	0.15	01(01,11)	42.29 ^c
Wyoming	2.33	0.40	02(01,49)	37.21 ^c
Nebraska	2.24	0.21	03(01,33)	32.04 ^c
Idaho	2.20	0.23	04(01,40)	29.87 ^c
Colorado	2.16 ^b	0.13	05(01,25)	27.31 ^c
Low Five States				
Louisiana	1.42	0.11	47(27,50)	-16.40 ^c
Mississippi	1.36	0.13	48(27,50)	-20.14 ^c
North Dakota	1.17	0.23	49(13,51)	-30.84 ^c
D.C.	1.14	0.27	50(07,51)	-32.58 ^c
Hawaii	0.77 ^b	0.13	51(48,51)	-54.43 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	1.71	0.11	32(08,47)	0.70	Montana	2.11	0.28	07(01,47)	24.45 ^c
Alaska	1.74	0.37	29(01,51)	2.42	Nebraska	2.24	0.21	03(01,33)	32.04 ^c
Arizona	1.72	0.10	30(10,45)	1.29	Nevada	1.67	0.16	33(05,49)	-1.76
Arkansas	1.58	0.14	37(10,50)	-6.93	New Hampshire	1.86	0.22	20(01,49)	9.76
California	1.56 ^b	0.04	40(29,46)	-7.88	New Jersey	1.62	0.08	35(18,47)	-4.51
Colorado	2.16 ^b	0.13	05(01,25)	27.31 ^c	New Mexico	1.71	0.17	31(04,49)	0.94
Connecticut	1.80	0.13	25(05,46)	5.94	New York	1.52 ^b	0.05	42(30,48)	-10.66
Delaware	1.98	0.27	10(01,48)	16.50 ^c	North Carolina	1.83	0.08	23(07,37)	8.02
D.C.	1.14	0.27	50(07,51)	-32.58 ^c	North Dakota	1.17	0.23	49(13,51)	-30.84 ^c
Florida	1.78	0.05	27(13,35)	4.85	Ohio	1.92	0.07	14(05,31)	13.18
Georgia	1.43	0.08	45(30,50)	-15.54 ^c	Oklahoma	1.91	0.13	15(02,42)	12.33
Hawaii	0.77 ^b	0.13	51(48,51)	-54.43 ^c	Oregon	2.42 ^b	0.15	01(01,11)	42.29 ^c
Idaho	2.20	0.23	04(01,40)	29.87 ^c	Pennsylvania	1.88	0.07	18(06,32)	11.00
Illinois	1.59	0.07	36(23,47)	-6.62	Rhode Island	1.49	0.21	44(06,50)	-12.05
Indiana	1.98	0.10	11(03,32)	16.36 ^c	South Carolina	1.57	0.11	38(14,49)	-7.37
Iowa	1.84	0.15	22(03,46)	8.13	South Dakota	1.84	0.28	21(01,50)	8.37
Kansas	1.86	0.15	19(03,46)	9.83	Tennessee	1.75	0.10	28(08,45)	2.88
Kentucky	2.16	0.13	06(01,26)	27.09 ^c	Texas	1.42 ^b	0.05	46(36,50)	-16.14 ^c
Louisiana	1.42	0.11	47(27,50)	-16.40 ^c	Utah	1.99	0.19	09(01,44)	17.38 ^c
Maine	1.91	0.21	16(01,48)	12.27	Vermont	2.08	0.32	08(01,49)	22.62 ^c
Maryland	1.57	0.10	39(18,49)	-7.37	Virginia	1.78	0.09	26(07,42)	5.05
Massachusetts	1.95	0.10	12(03,33)	14.95	Washington	1.89	0.10	17(04,37)	11.51
Michigan	1.53	0.07	41(25,49)	-10.17	West Virginia	1.94	0.19	13(01,46)	14.25
Minnesota	1.50	0.10	43(22,50)	-11.53	Wisconsin	1.65	0.10	34(12,48)	-2.56
Mississippi	1.36	0.13	48(27,50)	-20.14 ^c	Wyoming	2.33	0.40	02(01,49)	37.21 ^c
Missouri	1.81	0.10	24(05,42)	6.62					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 18.18
Myeloma

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	3.34	0.01		
High Five States				
D.C.	4.95 ^b	0.42	01(01,09)	48.14 ^c
Alabama	3.88 ^b	0.12	02(01,22)	15.96 ^c
Maryland	3.80 ^b	0.11	03(02,25)	13.62
Louisiana	3.79	0.13	04(01,26)	13.38
South Carolina	3.77	0.12	05(02,27)	12.76
Low Five States				
New Mexico	2.86	0.16	47(29,51)	-14.39
Vermont	2.85	0.27	48(10,51)	-14.71
Rhode Island	2.75	0.21	49(25,51)	-17.74 ^c
Nevada	2.74 ^b	0.15	50(36,51)	-18.06 ^c
Hawaii	2.42 ^b	0.17	51(45,51)	-27.72 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	3.88 ^b	0.12	02(01,22)	15.96 ^c	Montana	3.32	0.24	36(02,50)	-0.79
Alaska	3.46	0.42	22(01,51)	3.54	Nebraska	3.33	0.18	33(03,48)	-0.33
Arizona	2.90 ^b	0.09	46(36,50)	-13.15	Nevada	2.74 ^b	0.15	50(36,51)	-18.06 ^c
Arkansas	3.42	0.14	24(04,44)	2.44	New Hampshire	3.34	0.21	32(02,49)	-0.22
California	3.13 ^b	0.04	40(32,45)	-6.38	New Jersey	3.22	0.08	38(22,45)	-3.70
Colorado	3.01	0.12	43(29,50)	-9.90	New Mexico	2.86	0.16	47(29,51)	-14.39
Connecticut	2.92 ^b	0.12	45(32,51)	-12.71	New York	3.07 ^b	0.05	41(33,47)	-8.05
Delaware	3.68	0.27	10(01,47)	10.13	North Carolina	3.72 ^b	0.09	06(02,25)	11.28
D.C.	4.95 ^b	0.42	01(01,09)	48.14 ^c	North Dakota	3.58	0.31	17(01,49)	6.97
Florida	3.00 ^b	0.05	44(36,48)	-10.14	Ohio	3.56	0.07	18(05,31)	6.43
Georgia	3.64	0.09	13(02,29)	8.81	Oklahoma	3.50	0.13	21(03,40)	4.64
Hawaii	2.42 ^b	0.17	51(45,51)	-27.72 ^c	Oregon	3.41	0.13	26(05,43)	2.10
Idaho	3.39	0.21	29(02,48)	1.44	Pennsylvania	3.33	0.07	35(19,39)	-0.46
Illinois	3.37	0.07	30(15,38)	0.93	Rhode Island	2.75	0.21	49(25,51)	-17.74 ^c
Indiana	3.54	0.10	19(04,35)	5.90	South Carolina	3.77	0.12	05(02,27)	12.76
Iowa	3.40	0.14	28(05,44)	1.75	South Dakota	3.64	0.28	12(01,47)	8.86
Kansas	3.64	0.15	11(02,37)	9.00	Tennessee	3.63	0.10	14(02,32)	8.51
Kentucky	3.42	0.12	25(06,43)	2.26	Texas	3.17	0.05	39(29,44)	-5.07
Louisiana	3.79	0.13	04(01,26)	13.38	Utah	3.44	0.19	23(02,46)	2.83
Maine	3.03	0.19	42(16,51)	-9.43	Vermont	2.85	0.27	48(10,51)	-14.71
Maryland	3.80 ^b	0.11	03(02,25)	13.62	Virginia	3.72 ^b	0.10	07(02,25)	11.26
Massachusetts	3.26	0.09	37(18,44)	-2.60	Washington	3.37	0.10	31(11,41)	0.90
Michigan	3.70 ^b	0.08	09(02,25)	10.68	West Virginia	3.41	0.17	27(03,46)	1.96
Minnesota	3.50	0.11	20(04,38)	4.70	Wisconsin	3.70	0.11	08(02,29)	10.77
Mississippi	3.60	0.15	16(02,39)	7.79	Wyoming	3.33	0.35	34(02,51)	-0.46
Missouri	3.61	0.10	15(02,33)	8.10					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 18.19
Myeloma

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	4.24	0.02		
High Five States				
D.C.	6.18	0.74	01(01,36)	45.76 ^c
Louisiana	4.98	0.23	02(01,29)	17.51 ^c
Alabama	4.98	0.22	03(01,28)	17.40 ^c
Alaska	4.97	0.77	04(01,51)	17.15 ^c
Tennessee	4.87	0.19	05(01,28)	14.79
Low Five States				
New Mexico	3.55	0.27	47(24,51)	-16.26 ^c
Nevada	3.55	0.26	48(27,51)	-16.28 ^c
Vermont	3.40	0.46	49(08,51)	-19.85 ^c
Rhode Island	3.38	0.37	50(19,51)	-20.27 ^c
Hawaii	3.27	0.30	51(30,51)	-22.95 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	4.98	0.22	03(01,28)	17.40 ^c	Montana	4.03	0.39	39(02,51)	-4.90
Alaska	4.97	0.77	04(01,51)	17.15 ^c	Nebraska	4.41	0.32	26(02,49)	4.05
Arizona	3.76	0.16	43(29,50)	-11.29	Nevada	3.55	0.26	48(27,51)	-16.28 ^c
Arkansas	4.44	0.25	25(02,45)	4.77	New Hampshire	4.33	0.37	30(02,50)	2.05
California	4.01	0.07	40(28,45)	-5.38	New Jersey	4.07	0.14	37(20,47)	-3.94
Colorado	3.91	0.21	41(18,50)	-7.72	New Mexico	3.55	0.27	47(24,51)	-16.26 ^c
Connecticut	3.68	0.21	46(27,51)	-13.20	New York	3.79 ^b	0.09	42(34,49)	-10.56
Delaware	4.52	0.45	20(01,50)	6.71	North Carolina	4.64	0.15	12(03,32)	9.44
D.C.	6.18	0.74	01(01,36)	45.76 ^c	North Dakota	4.74	0.53	09(01,51)	11.80
Florida	3.73 ^b	0.08	45(36,50)	-12.08	Ohio	4.45	0.13	24(05,36)	5.03
Georgia	4.54	0.17	18(03,37)	7.19	Oklahoma	4.64	0.23	13(02,40)	9.36
Hawaii	3.27	0.30	51(30,51)	-22.95 ^c	Oregon	4.47	0.22	22(02,42)	5.44
Idaho	4.36	0.36	29(02,50)	2.74	Pennsylvania	4.15	0.11	33(19,43)	-2.02
Illinois	4.36	0.13	28(08,39)	2.77	Rhode Island	3.38	0.37	50(19,51)	-20.27 ^c
Indiana	4.70	0.18	10(02,34)	10.86	South Carolina	4.77	0.22	07(01,34)	12.40
Iowa	4.51	0.24	21(02,42)	6.34	South Dakota	4.20	0.45	32(01,51)	-0.83
Kansas	4.75	0.27	08(01,39)	11.97	Tennessee	4.87	0.19	05(01,28)	14.79
Kentucky	4.55	0.22	17(02,41)	7.22	Texas	4.05	0.10	38(25,45)	-4.57
Louisiana	4.98	0.23	02(01,29)	17.51 ^c	Utah	4.46	0.32	23(02,48)	5.20
Maine	3.73	0.32	44(10,51)	-11.96	Vermont	3.40	0.46	49(08,51)	-19.85 ^c
Maryland	4.52	0.20	19(03,40)	6.72	Virginia	4.69	0.17	11(02,33)	10.58
Massachusetts	4.13	0.16	35(14,46)	-2.52	Washington	4.36	0.17	27(06,43)	2.78
Michigan	4.63	0.14	14(03,32)	9.18	West Virginia	4.14	0.29	34(04,50)	-2.32
Minnesota	4.55	0.20	16(02,38)	7.33	Wisconsin	4.78	0.19	06(02,31)	12.67
Mississippi	4.27	0.27	31(03,48)	0.70	Wyoming	4.12	0.59	36(01,51)	-2.94
Missouri	4.58	0.18	15(02,37)	8.06					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 18.20
Myeloma

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	2.69	0.02		
High Five States				
D.C.	3.97	0.48	01(01,31)	47.76 ^c
Maryland	3.32 ^b	0.14	02(01,17)	23.55 ^c
Mississippi	3.18	0.19	03(01,30)	18.53 ^c
South Dakota	3.15	0.34	04(01,47)	17.29 ^c
Alabama	3.09	0.14	05(01,27)	15.13 ^c
Low Five States				
New Mexico	2.32	0.20	47(12,51)	-13.82
Alaska	2.22	0.45	48(01,51)	-17.39 ^c
Arizona	2.19 ^b	0.11	49(35,51)	-18.33 ^c
Nevada	2.05 ^b	0.18	50(32,51)	-23.76 ^c
Hawaii	1.79 ^b	0.20	51(39,51)	-33.32 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	3.09	0.14	05(01,27)	15.13 ^c	Montana	2.73	0.30	22(01,50)	1.80
Alaska	2.22	0.45	48(01,51)	-17.39 ^c	Nebraska	2.51	0.21	39(04,51)	-6.49
Arizona	2.19 ^b	0.11	49(35,51)	-18.33 ^c	Nevada	2.05 ^b	0.18	50(32,51)	-23.76 ^c
Arkansas	2.66	0.17	26(04,48)	-1.03	New Hampshire	2.59	0.25	32(02,51)	-3.40
California	2.48 ^b	0.05	40(27,46)	-7.78	New Jersey	2.63	0.10	27(13,45)	-2.20
Colorado	2.34	0.14	45(21,50)	-12.81	New Mexico	2.32	0.20	47(12,51)	-13.82
Connecticut	2.37	0.14	44(18,50)	-11.73	New York	2.58	0.06	35(20,44)	-3.90
Delaware	3.06	0.33	08(01,48)	14.09	North Carolina	3.09 ^b	0.10	06(02,21)	15.00 ^c
D.C.	3.97	0.48	01(01,31)	47.76 ^c	North Dakota	2.69	0.35	23(01,51)	0.02
Florida	2.44 ^b	0.06	42(28,48)	-9.29	Ohio	2.94	0.09	13(03,27)	9.47
Georgia	3.01	0.11	10(02,27)	11.98	Oklahoma	2.60	0.15	31(07,48)	-3.32
Hawaii	1.79 ^b	0.20	51(39,51)	-33.32 ^c	Oregon	2.56	0.15	36(09,49)	-4.78
Idaho	2.59	0.25	34(02,50)	-3.53	Pennsylvania	2.76	0.08	19(10,37)	2.59
Illinois	2.68	0.08	24(13,42)	-0.32	Rhode Island	2.34	0.26	46(06,51)	-12.91
Indiana	2.74	0.12	20(05,43)	2.15	South Carolina	3.07	0.15	07(01,29)	14.35
Iowa	2.56	0.16	37(09,49)	-4.85	South Dakota	3.15	0.34	04(01,47)	17.29 ^c
Kansas	2.85	0.18	16(02,45)	6.06	Tennessee	2.78	0.12	18(05,42)	3.44
Kentucky	2.63	0.14	28(08,48)	-2.23	Texas	2.52	0.06	38(22,46)	-6.13
Louisiana	2.97	0.15	12(02,37)	10.69	Utah	2.59	0.22	33(04,50)	-3.48
Maine	2.48	0.23	41(04,51)	-7.79	Vermont	2.39	0.33	43(02,51)	-10.95
Maryland	3.32 ^b	0.14	02(01,17)	23.55 ^c	Virginia	3.05	0.12	09(01,25)	13.37
Massachusetts	2.62	0.11	29(12,46)	-2.56	Washington	2.60	0.12	30(12,47)	-3.08
Michigan	3.00	0.10	11(02,25)	11.72	West Virginia	2.81	0.20	17(02,47)	4.71
Minnesota	2.67	0.13	25(07,46)	-0.66	Wisconsin	2.94	0.13	14(02,34)	9.43
Mississippi	3.18	0.19	03(01,30)	18.53 ^c	Wyoming	2.74	0.43	21(01,51)	1.96
Missouri	2.87	0.12	15(03,37)	6.84					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 19.22
Non-Hodgkin Lymphoma

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	6.15	0.02		
High Five States				
Minnesota	7.15 ^b	0.16	01(01,11)	16.23 ^c
Michigan	7.02 ^b	0.11	02(01,12)	14.18
Kentucky	7.01 ^b	0.18	03(01,17)	13.89
West Virginia	6.93	0.25	04(01,23)	12.69
Oklahoma	6.89 ^b	0.19	05(01,21)	12.05
Low Five States				
Nevada	5.48	0.21	47(24,51)	-10.85
Wyoming	5.42	0.45	48(07,51)	-11.84
North Dakota	5.36	0.37	49(16,51)	-12.92
New Mexico	5.12 ^b	0.22	50(36,51)	-16.78 ^c
D.C.	5.06	0.43	51(17,51)	-17.68 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	6.27	0.16	22(08,36)	1.94	Montana	6.01	0.32	26(03,50)	-2.31
Alaska	6.37	0.56	18(01,51)	3.52	Nebraska	6.29	0.25	21(03,43)	2.29
Arizona	5.79	0.13	36(23,47)	-5.88	Nevada	5.48	0.21	47(24,51)	-10.85
Arkansas	6.64	0.20	12(01,27)	8.00	New Hampshire	5.56	0.28	44(18,51)	-9.64
California	5.84 ^b	0.06	33(25,42)	-5.14	New Jersey	5.79	0.11	35(23,47)	-5.84
Colorado	5.48 ^b	0.16	46(26,51)	-10.85	New Mexico	5.12 ^b	0.22	50(36,51)	-16.78 ^c
Connecticut	5.58	0.16	42(25,51)	-9.29	New York	5.89 ^b	0.07	29(23,42)	-4.33
Delaware	5.75	0.34	38(08,51)	-6.48	North Carolina	5.86	0.11	30(22,45)	-4.67
D.C.	5.06	0.43	51(17,51)	-17.68 ^c	North Dakota	5.36	0.37	49(16,51)	-12.92
Florida	5.84 ^b	0.07	32(25,43)	-5.13	Ohio	6.86 ^b	0.10	06(01,16)	11.50
Georgia	5.49 ^b	0.12	45(32,51)	-10.77	Oklahoma	6.89 ^b	0.19	05(01,21)	12.05
Hawaii	5.75	0.27	39(13,51)	-6.57	Oregon	6.45	0.17	17(04,29)	4.81
Idaho	6.32	0.29	20(02,46)	2.73	Pennsylvania	6.72 ^b	0.09	08(03,19)	9.16
Illinois	6.34	0.10	19(11,27)	3.12	Rhode Island	5.69	0.31	40(12,51)	-7.55
Indiana	6.69 ^b	0.14	11(02,22)	8.81	South Carolina	5.82	0.16	34(20,48)	-5.39
Iowa	6.70	0.19	10(01,24)	8.94	South Dakota	6.09	0.36	25(01,50)	-0.99
Kansas	6.51	0.21	16(02,31)	5.80	Tennessee	6.59	0.14	13(03,24)	7.06
Kentucky	7.01 ^b	0.18	03(01,17)	13.89	Texas	5.95	0.08	28(22,40)	-3.29
Louisiana	6.58	0.17	14(02,26)	6.95	Utah	5.78	0.24	37(16,50)	-6.08
Maine	6.83	0.29	07(01,28)	11.08	Vermont	6.23	0.41	23(01,50)	1.30
Maryland	5.57 ^b	0.14	43(27,50)	-9.43	Virginia	6.01	0.12	27(17,41)	-2.38
Massachusetts	5.68 ^b	0.12	41(25,49)	-7.69	Washington	6.51	0.14	15(04,26)	5.82
Michigan	7.02 ^b	0.11	02(01,12)	14.18	West Virginia	6.93	0.25	04(01,23)	12.69
Minnesota	7.15 ^b	0.16	01(01,11)	16.23 ^c	Wisconsin	6.70 ^b	0.15	09(02,22)	8.95
Mississippi	5.84	0.20	31(17,50)	-5.12	Wyoming	5.42	0.45	48(07,51)	-11.84
Missouri	6.20	0.14	24(13,36)	0.80					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 19.23
Non-Hodgkin Lymphoma

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	7.92	0.03		
High Five States				
Minnesota	9.36 ^b	0.28	01(01,14)	18.13 ^c
Michigan	9.16 ^b	0.20	02(01,13)	15.71 ^c
Ohio	9.00 ^b	0.18	03(01,16)	13.67
Oklahoma	8.89	0.32	04(01,23)	12.24
Maine	8.82	0.50	05(01,36)	11.40
Low Five States				
Delaware	6.87	0.56	47(12,51)	-13.24
Nevada	6.82	0.35	48(23,51)	-13.90
Wyoming	6.32	0.74	49(13,51)	-20.22 ^c
North Dakota	6.31	0.61	50(18,51)	-20.26 ^c
New Mexico	6.05 ^b	0.36	51(42,51)	-23.63 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	7.86	0.28	24(09,44)	-0.80	Montana	8.00	0.55	21(01,49)	0.98
Alaska	7.81	0.92	26(01,51)	-1.34	Nebraska	7.91	0.43	23(02,47)	-0.10
Arizona	7.19	0.22	42(24,49)	-9.17	Nevada	6.82	0.35	48(23,51)	-13.90
Arkansas	8.17	0.35	19(02,42)	3.10	New Hampshire	7.17	0.48	43(09,51)	-9.43
California	7.44 ^b	0.10	35(25,44)	-6.06	New Jersey	7.41	0.19	37(21,47)	-6.46
Colorado	7.02	0.27	46(24,50)	-11.32	New Mexico	6.05 ^b	0.36	51(42,51)	-23.63 ^c
Connecticut	7.13	0.29	44(22,50)	-9.91	New York	7.59	0.13	31(21,43)	-4.12
Delaware	6.87	0.56	47(12,51)	-13.24	North Carolina	7.47	0.20	34(20,47)	-5.67
D.C.	7.05	0.80	45(01,51)	-10.97	North Dakota	6.31	0.61	50(18,51)	-20.26 ^c
Florida	7.49 ^b	0.12	33(23,44)	-5.42	Ohio	9.00 ^b	0.18	03(01,16)	13.67
Georgia	7.32	0.21	39(21,48)	-7.53	Oklahoma	8.89	0.32	04(01,23)	12.24
Hawaii	7.41	0.47	36(07,50)	-6.44	Oregon	8.33	0.30	16(02,36)	5.15
Idaho	8.01	0.49	20(01,48)	1.15	Pennsylvania	8.75 ^b	0.16	08(02,19)	10.46
Illinois	8.20	0.18	18(08,30)	3.50	Rhode Island	7.67	0.55	28(01,50)	-3.17
Indiana	8.64	0.25	12(01,24)	9.11	South Carolina	7.64	0.28	30(12,47)	-3.58
Iowa	8.75	0.34	07(01,27)	10.49	South Dakota	7.51	0.60	32(02,51)	-5.19
Kansas	8.73	0.36	09(01,28)	10.20	Tennessee	8.70	0.25	10(01,24)	9.81
Kentucky	8.81	0.31	06(01,24)	11.27	Texas	7.66	0.13	29(19,40)	-3.23
Louisiana	8.57	0.31	13(01,30)	8.22	Utah	7.19	0.41	41(15,51)	-9.16
Maine	8.82	0.50	05(01,36)	11.40	Vermont	8.25	0.73	17(01,50)	4.15
Maryland	7.29	0.25	40(21,49)	-7.95	Virginia	7.93	0.22	22(11,41)	0.08
Massachusetts	7.38	0.22	38(21,48)	-6.83	Washington	8.35	0.24	15(03,30)	5.41
Michigan	9.16 ^b	0.20	02(01,13)	15.71 ^c	West Virginia	8.44	0.42	14(01,41)	6.58
Minnesota	9.36 ^b	0.28	01(01,14)	18.13 ^c	Wisconsin	8.65	0.25	11(01,25)	9.21
Mississippi	7.85	0.36	25(05,47)	-0.91	Wyoming	6.32	0.74	49(13,51)	-20.22 ^c
Missouri	7.79	0.24	27(12,44)	-1.68					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 19.24
Non-Hodgkin Lymphoma

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	4.82	0.02		
High Five States				
West Virginia	5.78	0.30	01(01,23)	19.93 ^c
Kentucky	5.65 ^b	0.21	02(01,19)	17.32 ^c
Arkansas	5.51	0.25	03(01,26)	14.27
Minnesota	5.42	0.18	04(01,25)	12.36
Michigan	5.41 ^b	0.13	05(01,20)	12.29
Low Five States				
New Hampshire	4.30	0.32	47(09,51)	-10.79
Colorado	4.26	0.18	48(24,51)	-11.66
Rhode Island	4.23	0.34	49(10,51)	-12.31
Georgia	4.20 ^b	0.13	50(32,51)	-12.80
D.C.	3.67	0.47	51(15,51)	-23.91 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	5.12	0.19	13(02,35)	6.31	Montana	4.34	0.37	43(05,51)	-9.94
Alaska	5.02	0.69	17(01,51)	4.21	Nebraska	5.04	0.30	15(01,48)	4.51
Arizona	4.64	0.16	30(15,47)	-3.66	Nevada	4.33	0.26	45(14,51)	-10.12
Arkansas	5.51	0.25	03(01,26)	14.27	New Hampshire	4.30	0.32	47(09,51)	-10.79
California	4.60	0.07	35(23,43)	-4.61	New Jersey	4.67	0.13	28(16,45)	-3.15
Colorado	4.26	0.18	48(24,51)	-11.66	New Mexico	4.30	0.27	46(15,51)	-10.73
Connecticut	4.45	0.19	41(18,51)	-7.66	New York	4.66	0.09	29(20,43)	-3.24
Delaware	4.83	0.41	25(01,51)	0.11	North Carolina	4.69	0.13	27(16,45)	-2.71
D.C.	3.67	0.47	51(15,51)	-23.91 ^c	North Dakota	4.61	0.45	34(01,51)	-4.43
Florida	4.49 ^b	0.08	39(26,47)	-6.93	Ohio	5.28 ^b	0.12	09(02,23)	9.63
Georgia	4.20 ^b	0.13	50(32,51)	-12.80	Oklahoma	5.34	0.22	07(01,29)	10.75
Hawaii	4.47	0.32	40(04,51)	-7.24	Oregon	4.98	0.20	22(03,43)	3.32
Idaho	4.90	0.34	23(01,50)	1.71	Pennsylvania	5.22 ^b	0.11	12(03,24)	8.28
Illinois	5.03	0.11	16(06,30)	4.38	Rhode Island	4.23	0.34	49(10,51)	-12.31
Indiana	5.29	0.17	08(01,26)	9.80	South Carolina	4.50	0.18	37(17,50)	-6.62
Iowa	5.24	0.22	11(01,36)	8.71	South Dakota	5.02	0.43	18(01,51)	4.21
Kansas	4.80	0.24	26(03,49)	-0.37	Tennessee	5.00	0.16	21(04,38)	3.77
Kentucky	5.65 ^b	0.21	02(01,19)	17.32 ^c	Texas	4.64	0.09	31(20,43)	-3.79
Louisiana	5.02	0.20	19(02,39)	4.17	Utah	4.63	0.29	32(04,51)	-3.86
Maine	5.34	0.33	06(01,44)	10.81	Vermont	4.61	0.47	33(01,51)	-4.42
Maryland	4.34	0.16	44(24,51)	-10.01	Virginia	4.58	0.14	36(18,48)	-5.02
Massachusetts	4.50	0.14	38(21,49)	-6.65	Washington	5.09	0.17	14(03,34)	5.62
Michigan	5.41 ^b	0.13	05(01,20)	12.29	West Virginia	5.78	0.30	01(01,23)	19.93 ^c
Minnesota	5.42	0.18	04(01,25)	12.36	Wisconsin	5.25	0.17	10(01,28)	8.89
Mississippi	4.40	0.23	42(15,51)	-8.81	Wyoming	4.83	0.56	24(01,51)	0.23
Missouri	5.01	0.16	20(04,37)	3.89					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 20.24
Cancer of the Oral Cavity and Pharynx (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	2.45	0.01		
High Five States				
Mississippi	3.66 ^b	0.15	01(01,03)	49.18 ^c
D.C.	3.56	0.35	02(01,19)	45.18 ^c
Louisiana	3.22 ^b	0.12	03(01,09)	31.45 ^c
Arkansas	3.05 ^b	0.14	04(02,16)	24.23 ^c
Tennessee	3.01 ^b	0.09	05(02,12)	22.95 ^c
Low Five States				
Kansas	1.98 ^b	0.11	47(31,51)	-19.17 ^c
Minnesota	1.97 ^b	0.08	48(36,50)	-19.45 ^c
Connecticut	1.93 ^b	0.10	49(37,51)	-21.20 ^c
Colorado	1.92 ^b	0.09	50(38,51)	-21.76 ^c
Utah	1.54 ^b	0.12	51(47,51)	-37.06 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	2.70	0.10	12(05,30)	10.05	Montana	2.03	0.18	44(17,51)	-17.25 ^c
Alaska	2.06	0.29	42(06,51)	-15.97 ^c	Nebraska	2.47	0.16	26(07,45)	0.90
Arizona	2.00 ^b	0.08	45(36,50)	-18.62 ^c	Nevada	2.36	0.13	31(10,46)	-3.71
Arkansas	3.05 ^b	0.14	04(02,16)	24.23 ^c	New Hampshire	2.42	0.18	29(05,48)	-1.28
California	2.46	0.04	28(17,34)	0.21	New Jersey	2.06 ^b	0.06	41(34,49)	-15.81 ^c
Colorado	1.92 ^b	0.09	50(38,51)	-21.76 ^c	New Mexico	2.32	0.14	35(10,48)	-5.44
Connecticut	1.93 ^b	0.10	49(37,51)	-21.20 ^c	New York	2.18 ^b	0.04	38(32,45)	-11.07
Delaware	2.74	0.23	10(02,43)	11.66	North Carolina	2.60	0.07	17(08,31)	6.09
D.C.	3.56	0.35	02(01,19)	45.18 ^c	North Dakota	2.04	0.23	43(09,51)	-16.76 ^c
Florida	2.66 ^b	0.05	15(08,23)	8.46	Ohio	2.53	0.06	20(11,33)	3.05
Georgia	2.53	0.08	19(10,35)	3.15	Oklahoma	2.67	0.11	14(05,35)	8.94
Hawaii	2.61	0.18	16(03,43)	6.29	Oregon	2.34	0.10	34(14,44)	-4.61
Idaho	2.50	0.18	22(04,46)	1.92	Pennsylvania	2.20 ^b	0.05	37(30,45)	-10.35
Illinois	2.41	0.06	30(16,38)	-1.58	Rhode Island	2.00	0.18	46(16,51)	-18.62 ^c
Indiana	2.35	0.08	32(16,42)	-4.04	South Carolina	2.88 ^b	0.11	07(03,18)	17.64 ^c
Iowa	2.17	0.11	39(22,50)	-11.67	South Dakota	2.12	0.22	40(09,51)	-13.58
Kansas	1.98 ^b	0.11	47(31,51)	-19.17 ^c	Tennessee	3.01 ^b	0.09	05(02,12)	22.95 ^c
Kentucky	2.80	0.11	08(03,25)	14.36	Texas	2.54	0.05	18(12,31)	3.62
Louisiana	3.22 ^b	0.12	03(01,09)	31.45 ^c	Utah	1.54 ^b	0.12	51(47,51)	-37.06 ^c
Maine	2.69	0.18	13(03,39)	9.70	Vermont	2.52	0.26	21(03,50)	2.67
Maryland	2.35	0.09	33(14,42)	-4.14	Virginia	2.29	0.07	36(20,43)	-6.43
Massachusetts	2.49	0.08	25(10,37)	1.52	Washington	2.92 ^b	0.09	06(02,15)	18.99 ^c
Michigan	2.49	0.07	23(12,36)	1.65	West Virginia	2.70	0.15	11(03,36)	10.08
Minnesota	1.97 ^b	0.08	48(36,50)	-19.45 ^c	Wisconsin	2.49	0.09	24(10,38)	1.64
Mississippi	3.66 ^b	0.15	01(01,03)	49.18 ^c	Wyoming	2.75	0.30	09(01,49)	12.14
Missouri	2.47	0.09	27(11,37)	0.83					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 20.25
Cancer of the Oral Cavity and Pharynx (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	3.77	0.02		
High Five States				
Mississippi	6.09 ^b	0.30	01(01,03)	61.33 ^c
D.C.	5.90	0.69	02(01,22)	56.32 ^c
Louisiana	5.28 ^b	0.23	03(01,08)	39.88 ^c
Tennessee	4.80 ^b	0.18	04(02,12)	27.19 ^c
South Carolina	4.71 ^b	0.20	05(02,15)	24.87 ^c
Low Five States				
Kansas	3.03 ^b	0.21	47(27,50)	-19.63 ^c
Montana	2.92	0.32	48(19,51)	-22.56 ^c
Minnesota	2.79 ^b	0.15	49(39,50)	-26.06 ^c
Colorado	2.65 ^b	0.16	50(42,51)	-29.76 ^c
Utah	2.00 ^b	0.20	51(49,51)	-47.10 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	4.30	0.19	10(04,27)	13.89	Montana	2.92	0.32	48(19,51)	-22.56 ^c
Alaska	3.37	0.55	39(03,51)	-10.65	Nebraska	3.47	0.28	33(10,49)	-7.90
Arizona	3.16 ^b	0.14	43(30,49)	-16.22 ^c	Nevada	3.53	0.24	32(11,48)	-6.47
Arkansas	4.67 ^b	0.25	06(02,20)	23.91 ^c	New Hampshire	3.71	0.33	24(05,49)	-1.74
California	3.71	0.07	26(18,35)	-1.77	New Jersey	3.10 ^b	0.12	46(33,49)	-17.89 ^c
Colorado	2.65 ^b	0.16	50(42,51)	-29.76 ^c	New Mexico	3.46	0.26	35(11,49)	-8.27
Connecticut	3.11 ^b	0.19	45(26,50)	-17.61 ^c	New York	3.40 ^b	0.09	38(27,44)	-9.93
Delaware	4.59	0.45	07(01,41)	21.72 ^c	North Carolina	4.10	0.14	14(07,28)	8.54
D.C.	5.90	0.69	02(01,22)	56.32 ^c	North Dakota	3.41	0.44	37(05,50)	-9.59
Florida	4.11 ^b	0.09	12(08,23)	8.88	Ohio	3.93	0.12	20(10,32)	4.21
Georgia	3.98	0.15	19(08,33)	5.51	Oklahoma	4.02	0.21	16(06,37)	6.67
Hawaii	4.07	0.34	15(03,44)	7.91	Oregon	3.44	0.19	36(16,48)	-8.84
Idaho	3.72	0.32	23(06,48)	-1.53	Pennsylvania	3.36 ^b	0.10	40(27,45)	-10.85
Illinois	3.64	0.11	28(17,40)	-3.57	Rhode Island	3.15	0.34	44(11,51)	-16.59 ^c
Indiana	3.62	0.15	29(14,43)	-4.14	South Carolina	4.71 ^b	0.20	05(02,15)	24.87 ^c
Iowa	3.47	0.21	34(13,48)	-8.01	South Dakota	3.18	0.39	42(09,51)	-15.79 ^c
Kansas	3.03 ^b	0.21	47(27,50)	-19.63 ^c	Tennessee	4.80 ^b	0.18	04(02,12)	27.19 ^c
Kentucky	4.38	0.21	09(03,24)	16.08 ^c	Texas	3.99	0.09	18(09,27)	5.68
Louisiana	5.28 ^b	0.23	03(01,08)	39.88 ^c	Utah	2.00 ^b	0.20	51(49,51)	-47.10 ^c
Maine	4.23	0.34	11(03,41)	12.03	Vermont	3.31	0.45	41(06,51)	-12.29
Maryland	3.57	0.17	31(14,45)	-5.33	Virginia	3.66	0.14	27(14,41)	-2.86
Massachusetts	3.91	0.16	21(08,35)	3.65	Washington	4.10	0.16	13(06,31)	8.66
Michigan	3.89	0.12	22(10,34)	3.20	West Virginia	4.46	0.29	08(02,30)	18.16 ^c
Minnesota	2.79 ^b	0.15	49(39,50)	-26.06 ^c	Wisconsin	3.59	0.16	30(14,44)	-4.76
Mississippi	6.09 ^b	0.30	01(01,03)	61.33 ^c	Wyoming	4.01	0.54	17(02,50)	6.40
Missouri	3.71	0.16	25(12,42)	-1.75					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 20.26
Cancer of the Oral Cavity and Pharynx (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	1.35	0.01		
High Five States				
Washington	1.91 ^b	0.10	01(01,08)	41.31 ^c
Vermont	1.81	0.28	02(01,45)	34.40 ^c
Mississippi	1.75	0.14	03(01,29)	29.46 ^c
D.C.	1.73	0.32	04(01,50)	28.15 ^c
Arkansas	1.69	0.14	05(01,30)	25.12 ^c
Low Five States				
Rhode Island	1.08	0.18	46(05,50)	-19.84 ^c
Iowa	1.07	0.10	47(21,50)	-21.02 ^c
Connecticut	1.01 ^b	0.09	48(30,50)	-25.43 ^c
Arizona	1.00 ^b	0.07	49(36,50)	-26.19 ^c
North Dakota	0.94	0.23	50(04,50)	-30.06 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	1.39	0.10	19(05,44)	2.64	Montana	1.21	0.20	40(02,50)	-10.07
Alaska	-	-	-(-,-)	-	Nebraska	1.61	0.17	07(01,43)	19.15 ^c
Arizona	1.00 ^b	0.07	49(36,50)	-26.19 ^c	Nevada	1.31	0.14	31(04,49)	-3.14
Arkansas	1.69	0.14	05(01,30)	25.12 ^c	New Hampshire	1.30	0.19	32(02,50)	-3.86
California	1.41	0.04	15(08,32)	4.45	New Jersey	1.25	0.07	36(13,46)	-7.09
Colorado	1.28	0.10	35(07,48)	-5.31	New Mexico	1.32	0.15	28(03,50)	-2.17
Connecticut	1.01 ^b	0.09	48(30,50)	-25.43 ^c	New York	1.22	0.04	39(23,46)	-9.73
Delaware	1.24	0.21	38(02,50)	-8.10	North Carolina	1.40	0.07	16(06,40)	3.47
D.C.	1.73	0.32	04(01,50)	28.15 ^c	North Dakota	0.94	0.23	50(04,50)	-30.06 ^c
Florida	1.42	0.05	13(07,34)	5.01	Ohio	1.36	0.06	24(08,42)	0.37
Georgia	1.36	0.07	22(07,42)	0.59	Oklahoma	1.50	0.12	12(02,41)	11.21
Hawaii	1.32	0.17	30(02,50)	-2.57	Oregon	1.39	0.11	17(04,45)	3.05
Idaho	1.36	0.18	21(02,50)	0.97	Pennsylvania	1.25	0.05	37(16,45)	-7.22
Illinois	1.41	0.06	14(06,38)	4.46	Rhode Island	1.08	0.18	46(05,50)	-19.84 ^c
Indiana	1.28	0.08	34(09,47)	-5.16	South Carolina	1.36	0.10	23(05,45)	0.51
Iowa	1.07	0.10	47(21,50)	-21.02 ^c	South Dakota	1.17	0.21	42(03,50)	-13.42
Kansas	1.09	0.11	45(13,50)	-19.42 ^c	Tennessee	1.55	0.09	09(02,32)	14.62
Kentucky	1.51	0.11	11(02,39)	12.05	Texas	1.33	0.05	26(12,41)	-1.37
Louisiana	1.52	0.11	10(02,39)	12.91	Utah	1.12	0.14	44(09,50)	-16.75 ^c
Maine	1.39	0.17	18(02,50)	2.78	Vermont	1.81	0.28	02(01,45)	34.40 ^c
Maryland	1.33	0.09	25(06,46)	-1.35	Virginia	1.21	0.07	41(15,48)	-10.59
Massachusetts	1.33	0.08	27(07,45)	-1.78	Washington	1.91 ^b	0.10	01(01,08)	41.31 ^c
Michigan	1.32	0.07	29(09,44)	-2.24	West Virginia	1.14	0.13	43(08,50)	-15.79 ^c
Minnesota	1.29	0.09	33(07,47)	-4.29	Wisconsin	1.55	0.10	08(02,33)	15.10 ^c
Mississippi	1.75	0.14	03(01,29)	29.46 ^c	Wyoming	1.61	0.32	06(01,50)	19.60 ^c
Missouri	1.37	0.08	20(06,44)	1.66					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 21.14
Cancer of the Ovary (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	7.71	0.03		
High Five States				
Oregon	8.98 ^b	0.28	01(01,12)	16.58 ^c
Washington	8.70 ^b	0.22	02(01,15)	12.91
Montana	8.67	0.52	03(01,41)	12.54
Alabama	8.65 ^b	0.24	04(01,19)	12.30
Iowa	8.44	0.29	05(01,31)	9.53
Low Five States				
Louisiana	6.96	0.23	47(28,50)	-9.71
North Dakota	6.78	0.57	48(05,51)	-11.97
Alaska	6.70	0.74	49(01,51)	-13.07
Rhode Island	6.57	0.44	50(19,51)	-14.70
Hawaii	5.31 ^b	0.36	51(49,51)	-31.13 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	8.65 ^b	0.24	04(01,19)	12.30	Montana	8.67	0.52	03(01,41)	12.54
Alaska	6.70	0.74	49(01,51)	-13.07	Nebraska	7.00	0.36	45(13,50)	-9.11
Arizona	7.35	0.20	36(17,48)	-4.64	Nevada	6.97	0.32	46(17,50)	-9.57
Arkansas	8.16	0.30	10(01,38)	5.93	New Hampshire	7.66	0.43	28(01,49)	-0.60
California	7.67	0.09	27(16,37)	-0.48	New Jersey	8.26	0.17	07(02,25)	7.25
Colorado	8.26	0.25	08(01,31)	7.22	New Mexico	7.84	0.36	20(01,48)	1.78
Connecticut	7.32	0.25	38(14,49)	-5.03	New York	7.74	0.11	24(13,37)	0.41
Delaware	7.67	0.52	26(01,50)	-0.43	North Carolina	7.51	0.16	33(15,44)	-2.50
D.C.	7.17	0.66	40(01,51)	-6.99	North Dakota	6.78	0.57	48(05,51)	-11.97
Florida	7.04 ^b	0.10	43(34,49)	-8.62	Ohio	7.96	0.15	16(06,33)	3.27
Georgia	7.87	0.18	19(06,38)	2.17	Oklahoma	8.06	0.27	13(02,39)	4.55
Hawaii	5.31 ^b	0.36	51(49,51)	-31.13 ^c	Oregon	8.98 ^b	0.28	01(01,12)	16.58 ^c
Idaho	7.58	0.42	32(02,50)	-1.67	Pennsylvania	8.26 ^b	0.14	09(03,22)	7.12
Illinois	7.77	0.15	23(09,38)	0.87	Rhode Island	6.57	0.44	50(19,51)	-14.70
Indiana	8.15	0.21	11(02,33)	5.81	South Carolina	7.81	0.24	22(05,42)	1.35
Iowa	8.44	0.29	05(01,31)	9.53	South Dakota	7.09	0.53	41(03,51)	-8.01
Kansas	7.88	0.31	18(02,44)	2.25	Tennessee	7.72	0.20	25(08,42)	0.16
Kentucky	7.48	0.24	34(10,48)	-2.93	Texas	7.26 ^b	0.11	39(29,46)	-5.80
Louisiana	6.96	0.23	47(28,50)	-9.71	Utah	7.39	0.36	35(05,50)	-4.05
Maine	7.62	0.40	30(02,49)	-1.12	Vermont	7.32	0.60	37(01,51)	-5.01
Maryland	7.64	0.21	29(08,45)	-0.89	Virginia	8.03	0.19	14(04,35)	4.25
Massachusetts	7.61	0.19	31(10,44)	-1.27	Washington	8.70 ^b	0.22	02(01,15)	12.91
Michigan	7.92	0.16	17(06,36)	2.75	West Virginia	8.01	0.36	15(01,45)	3.88
Minnesota	7.84	0.23	21(04,41)	1.68	Wisconsin	8.12	0.22	12(02,34)	5.38
Mississippi	7.00	0.29	44(19,50)	-9.11	Wyoming	8.33	0.75	06(01,50)	8.15
Missouri	7.05	0.19	42(27,50)	-8.50					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 22.18
Cancer of the Pancreas (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	10.94	0.03		
High Five States				
D.C.	13.88 ^b	0.71	01(01,08)	26.94 ^c
Louisiana	12.99 ^b	0.24	02(01,04)	18.76 ^c
Mississippi	12.35 ^b	0.28	03(01,12)	12.90
Vermont	11.75	0.56	04(01,45)	7.45
Michigan	11.62 ^b	0.14	05(03,23)	6.25
Low Five States				
Colorado	9.84 ^b	0.21	47(36,51)	-10.02
Montana	9.61	0.40	48(30,51)	-12.10
Wyoming	9.60	0.58	49(14,51)	-12.23
New Mexico	9.43 ^b	0.29	50(41,51)	-13.81
Utah	9.41 ^b	0.30	51(41,51)	-13.97

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	11.43	0.21	14(04,31)	4.52	Montana	9.61	0.40	48(30,51)	-12.10
Alaska	11.50	0.73	11(01,50)	5.18	Nebraska	10.61	0.33	34(08,49)	-3.01
Arizona	10.08 ^b	0.17	46(34,50)	-7.80	Nevada	10.72	0.29	32(07,48)	-1.95
Arkansas	11.15	0.26	23(04,40)	1.98	New Hampshire	11.32	0.39	21(03,44)	3.54
California	10.43 ^b	0.08	37(32,45)	-4.61	New Jersey	11.62 ^b	0.15	06(03,24)	6.25
Colorado	9.84 ^b	0.21	47(36,51)	-10.02	New Mexico	9.43 ^b	0.29	50(41,51)	-13.81
Connecticut	11.51	0.24	10(03,32)	5.23	New York	11.33 ^b	0.10	20(07,28)	3.56
Delaware	11.36	0.47	17(02,46)	3.83	North Carolina	10.44	0.14	36(28,47)	-4.53
D.C.	13.88 ^b	0.71	01(01,08)	26.94 ^c	North Dakota	10.38	0.51	41(04,51)	-5.06
Florida	10.42 ^b	0.09	39(31,45)	-4.69	Ohio	11.53 ^b	0.13	09(04,23)	5.46
Georgia	10.40	0.16	40(28,47)	-4.86	Oklahoma	10.85	0.23	31(10,44)	-0.77
Hawaii	11.48	0.38	12(03,41)	4.97	Oregon	10.88	0.22	30(09,44)	-0.56
Idaho	10.51	0.37	35(09,50)	-3.88	Pennsylvania	11.55 ^b	0.12	08(04,23)	5.58
Illinois	11.33	0.13	19(06,29)	3.57	Rhode Island	10.22	0.41	43(14,51)	-6.52
Indiana	10.94	0.18	28(12,39)	0.07	South Carolina	11.39	0.21	15(04,33)	4.19
Iowa	10.70	0.24	33(13,47)	-2.20	South Dakota	10.15	0.46	45(10,51)	-7.15
Kansas	11.07	0.27	26(05,42)	1.25	Tennessee	11.07	0.18	25(08,36)	1.25
Kentucky	10.98	0.22	27(08,42)	0.36	Texas	10.26 ^b	0.10	42(33,47)	-6.14
Louisiana	12.99 ^b	0.24	02(01,04)	18.76 ^c	Utah	9.41 ^b	0.30	51(41,51)	-13.97
Maine	11.36	0.36	16(03,43)	3.89	Vermont	11.75	0.56	04(01,45)	7.45
Maryland	11.55	0.20	07(03,28)	5.64	Virginia	10.88	0.16	29(15,40)	-0.49
Massachusetts	11.28	0.17	22(05,33)	3.12	Washington	11.35	0.18	18(04,32)	3.81
Michigan	11.62 ^b	0.14	05(03,23)	6.25	West Virginia	10.16	0.29	44(24,51)	-7.06
Minnesota	10.43	0.19	38(26,47)	-4.64	Wisconsin	11.48	0.19	13(04,30)	4.94
Mississippi	12.35 ^b	0.28	03(01,12)	12.90	Wyoming	9.60	0.58	49(14,51)	-12.23
Missouri	11.14	0.18	24(06,35)	1.87					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 22.19
Cancer of the Pancreas (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	12.56	0.04		
High Five States				
D.C.	15.70	1.17	01(01,35)	24.98 ^c
Louisiana	14.87 ^b	0.39	02(01,07)	18.42 ^c
Mississippi	14.42 ^b	0.48	03(01,17)	14.84 ^c
Alaska	13.70	1.19	04(01,50)	9.04
Delaware	13.65	0.78	05(01,47)	8.65
Low Five States				
New Mexico	11.12	0.48	47(27,51)	-11.47 ^c
Utah	10.94	0.48	48(29,51)	-12.92 ^c
Wyoming	10.87	0.91	49(06,51)	-13.46 ^c
South Dakota	10.79	0.71	50(15,51)	-14.07 ^c
Colorado	10.79 ^b	0.33	51(41,51)	-14.13 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	13.31	0.35	13(03,34)	5.96	Montana	11.21	0.65	46(13,51)	-10.78 ^c
Alaska	13.70	1.19	04(01,50)	9.04	Nebraska	11.98	0.52	40(08,51)	-4.59
Arizona	11.47 ^b	0.27	45(33,51)	-8.69	Nevada	12.48	0.47	31(05,48)	-0.61
Arkansas	13.19	0.43	15(02,40)	5.00	New Hampshire	13.59	0.65	06(01,43)	8.18
California	11.75 ^b	0.12	44(35,48)	-6.43	New Jersey	13.29	0.26	14(04,30)	5.80
Colorado	10.79 ^b	0.33	51(41,51)	-14.13 ^c	New Mexico	11.12	0.48	47(27,51)	-11.47 ^c
Connecticut	13.19	0.39	16(03,39)	4.98	New York	13.05	0.17	18(06,31)	3.89
Delaware	13.65	0.78	05(01,47)	8.65	North Carolina	11.92	0.24	42(26,48)	-5.07
D.C.	15.70	1.17	01(01,35)	24.98 ^c	North Dakota	13.33	0.88	11(01,50)	6.13
Florida	12.13	0.15	37(26,45)	-3.41	Ohio	13.38 ^b	0.22	10(03,25)	6.54
Georgia	12.19	0.27	36(18,47)	-2.94	Oklahoma	12.35	0.37	33(08,47)	-1.65
Hawaii	12.87	0.60	22(02,48)	2.44	Oregon	12.27	0.36	35(10,48)	-2.28
Idaho	12.28	0.58	34(05,50)	-2.27	Pennsylvania	13.31 ^b	0.20	12(04,26)	6.00
Illinois	13.00	0.22	20(06,33)	3.49	Rhode Island	12.67	0.70	27(02,50)	0.85
Indiana	12.76	0.30	24(06,41)	1.54	South Carolina	13.02	0.35	19(04,40)	3.67
Iowa	12.43	0.39	32(07,47)	-1.07	South Dakota	10.79	0.71	50(15,51)	-14.07 ^c
Kansas	12.68	0.44	26(04,46)	0.98	Tennessee	12.92	0.30	21(05,40)	2.87
Kentucky	12.84	0.36	23(04,43)	2.22	Texas	11.84 ^b	0.16	43(31,48)	-5.78
Louisiana	14.87 ^b	0.39	02(01,07)	18.42 ^c	Utah	10.94	0.48	48(29,51)	-12.92 ^c
Maine	12.05	0.57	38(06,51)	-4.07	Vermont	13.39	0.90	09(01,49)	6.63
Maryland	13.41	0.33	08(03,31)	6.78	Virginia	12.54	0.27	30(10,44)	-0.20
Massachusetts	12.63	0.29	29(08,43)	0.57	Washington	12.74	0.29	25(07,42)	1.44
Michigan	13.42 ^b	0.24	07(03,26)	6.81	West Virginia	11.97	0.49	41(09,50)	-4.71
Minnesota	12.05	0.31	39(16,48)	-4.08	Wisconsin	13.11	0.31	17(04,36)	4.35
Mississippi	14.42 ^b	0.48	03(01,17)	14.84 ^c	Wyoming	10.87	0.91	49(06,51)	-13.46 ^c
Missouri	12.66	0.30	28(07,42)	0.83					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 10% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 22.20
Cancer of the Pancreas (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	9.59	0.03		
High Five States				
D.C.	12.20	0.87	01(01,23)	27.20 ^c
Louisiana	11.38 ^b	0.30	02(01,06)	18.65 ^c
Maine	10.63	0.47	03(01,33)	10.78
Mississippi	10.60	0.35	04(01,26)	10.48
New Jersey	10.29 ^b	0.19	05(03,24)	7.23
Low Five States				
Wyoming	8.35	0.73	47(04,51)	-12.99
Montana	8.21	0.51	48(22,51)	-14.41
Utah	8.11 ^b	0.38	49(32,51)	-15.43 ^c
New Mexico	7.97 ^b	0.36	50(37,51)	-16.95 ^c
North Dakota	7.83	0.60	51(24,51)	-18.41 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	9.89	0.26	19(04,37)	3.09	Montana	8.21	0.51	48(22,51)	-14.41
Alaska	9.24	0.88	35(01,51)	-3.69	Nebraska	9.37	0.41	32(04,49)	-2.35
Arizona	8.86	0.22	44(27,49)	-7.68	Nevada	9.20	0.37	37(07,49)	-4.04
Arkansas	9.40	0.32	31(06,47)	-2.04	New Hampshire	9.45	0.48	29(03,49)	-1.51
California	9.34	0.10	33(23,40)	-2.64	New Jersey	10.29 ^b	0.19	05(03,24)	7.23
Colorado	9.00	0.27	39(20,49)	-6.12	New Mexico	7.97 ^b	0.36	50(37,51)	-16.95 ^c
Connecticut	10.20	0.30	09(02,33)	6.29	New York	9.97	0.13	17(06,28)	3.95
Delaware	9.49	0.57	28(02,50)	-1.04	North Carolina	9.20	0.18	36(22,46)	-4.04
D.C.	12.20	0.87	01(01,23)	27.20 ^c	North Dakota	7.83	0.60	51(24,51)	-18.41 ^c
Florida	8.98 ^b	0.11	41(30,46)	-6.42	Ohio	10.05	0.16	16(04,27)	4.81
Georgia	9.00	0.19	40(25,48)	-6.20	Oklahoma	9.56	0.29	25(05,45)	-0.39
Hawaii	10.24	0.49	07(01,44)	6.76	Oregon	9.67	0.29	22(04,43)	0.86
Idaho	8.88	0.46	43(07,51)	-7.38	Pennsylvania	10.09	0.15	14(04,26)	5.22
Illinois	9.95	0.16	18(05,30)	3.76	Rhode Island	8.38	0.49	46(18,51)	-12.63
Indiana	9.51	0.22	27(08,42)	-0.83	South Carolina	10.09	0.27	15(03,33)	5.17
Iowa	9.25	0.30	34(10,48)	-3.57	South Dakota	9.58	0.61	24(02,50)	-0.17
Kansas	9.71	0.34	21(04,45)	1.19	Tennessee	9.64	0.22	23(07,40)	0.55
Kentucky	9.44	0.27	30(09,45)	-1.57	Texas	8.95 ^b	0.12	42(31,47)	-6.73
Louisiana	11.38 ^b	0.30	02(01,06)	18.65 ^c	Utah	8.11 ^b	0.38	49(32,51)	-15.43 ^c
Maine	10.63	0.47	03(01,33)	10.78	Vermont	10.25	0.71	06(01,48)	6.82
Maryland	10.15	0.25	11(03,30)	5.85	Virginia	9.55	0.21	26(10,41)	-0.49
Massachusetts	10.15	0.22	10(03,28)	5.85	Washington	10.21	0.24	08(02,29)	6.42
Michigan	10.12	0.18	12(03,27)	5.54	West Virginia	8.65	0.36	45(22,51)	-9.85
Minnesota	9.04	0.24	38(21,48)	-5.81	Wisconsin	10.12	0.24	13(03,31)	5.48
Mississippi	10.60	0.35	04(01,26)	10.48	Wyoming	8.35	0.73	47(04,51)	-12.99
Missouri	9.85	0.23	20(05,36)	2.64					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 23.14
Cancer of the Prostate (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	21.41	0.06		
High Five States				
D.C.	34.92 ^b	1.84	01(01,01)	63.11 ^c
Mississippi	28.44 ^b	0.73	02(02,03)	32.84 ^c
Alabama	26.37 ^b	0.53	03(02,06)	23.19 ^c
South Carolina	25.46 ^b	0.54	04(03,09)	18.91 ^c
Idaho	24.67 ^b	0.88	05(03,27)	15.23 ^c
Low Five States				
Texas	19.59 ^b	0.22	47(40,49)	-8.49
Arizona	19.44 ^b	0.37	48(37,50)	-9.19
Kansas	19.23 ^b	0.55	49(35,50)	-10.17 ^c
Florida	18.70 ^b	0.18	50(47,50)	-12.64 ^c
Hawaii	14.84 ^b	0.66	51(51,51)	-30.70 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	26.37 ^b	0.53	03(02,06)	23.19 ^c	Montana	23.44	0.97	09(03,38)	9.50
Alaska	21.89	1.72	28(03,50)	2.23	Nebraska	21.99	0.73	26(07,45)	2.72
Arizona	19.44 ^b	0.37	48(37,50)	-9.19	Nevada	21.45	0.67	31(10,47)	0.21
Arkansas	22.75	0.60	13(06,35)	6.26	New Hampshire	20.81	0.85	35(10,50)	-2.81
California	21.12	0.17	32(27,40)	-1.34	New Jersey	20.32	0.33	42(30,48)	-5.10
Colorado	22.56	0.51	17(07,34)	5.37	New Mexico	21.86	0.71	29(07,46)	2.13
Connecticut	20.20	0.49	44(27,50)	-5.65	New York	20.60 ^b	0.22	39(31,45)	-3.79
Delaware	22.55	1.07	18(04,47)	5.34	North Carolina	23.49 ^b	0.37	08(05,21)	9.70
D.C.	34.92 ^b	1.84	01(01,01)	63.11 ^c	North Dakota	22.57	1.15	16(03,48)	5.42
Florida	18.70 ^b	0.18	50(47,50)	-12.64 ^c	Ohio	22.02	0.29	25(13,34)	2.86
Georgia	24.57 ^b	0.42	06(04,12)	14.76 ^c	Oklahoma	22.39	0.54	22(08,36)	4.56
Hawaii	14.84 ^b	0.66	51(51,51)	-30.70 ^c	Oregon	23.21 ^b	0.51	11(05,29)	8.40
Idaho	24.67 ^b	0.88	05(03,27)	15.23 ^c	Pennsylvania	21.08	0.26	33(26,42)	-1.53
Illinois	22.39	0.30	21(10,31)	4.57	Rhode Island	20.80	0.91	36(09,50)	-2.83
Indiana	21.98	0.41	27(11,38)	2.65	South Carolina	25.46 ^b	0.54	04(03,09)	18.91 ^c
Iowa	20.73	0.52	38(20,49)	-3.20	South Dakota	20.75	1.01	37(08,50)	-3.08
Kansas	19.23 ^b	0.55	49(35,50)	-10.17 ^c	Tennessee	22.58	0.43	15(08,32)	5.48
Kentucky	21.52	0.51	30(12,44)	0.49	Texas	19.59 ^b	0.22	47(40,49)	-8.49
Louisiana	24.21 ^b	0.54	07(04,18)	13.08 ^c	Utah	23.31	0.75	10(04,34)	8.88
Maine	21.05	0.79	34(10,50)	-1.66	Vermont	22.35	1.24	23(03,49)	4.41
Maryland	22.51	0.45	19(08,33)	5.12	Virginia	22.65	0.40	14(08,31)	5.79
Massachusetts	20.48	0.37	40(28,48)	-4.33	Washington	22.20	0.40	24(09,35)	3.69
Michigan	20.30 ^b	0.31	43(31,48)	-5.16	West Virginia	20.13	0.67	45(23,50)	-5.97
Minnesota	22.50	0.44	20(08,33)	5.11	Wisconsin	23.13 ^b	0.42	12(06,27)	8.02
Mississippi	28.44 ^b	0.73	02(02,03)	32.84 ^c	Wyoming	20.34	1.34	41(06,50)	-4.99
Missouri	19.89 ^b	0.39	46(32,50)	-7.10					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 10% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 24.18
Cancer of the Stomach (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	3.37	0.01		
High Five States				
Hawaii	5.26 ^b	0.26	01(01,03)	56.00 ^c
Louisiana	4.48 ^b	0.14	02(01,07)	32.83 ^c
D.C.	4.48	0.40	03(01,19)	32.80 ^c
California	4.26 ^b	0.05	04(02,07)	26.48 ^c
Alaska	4.16	0.43	05(01,32)	23.45 ^c
Low Five States				
Wyoming	2.39	0.29	47(19,51)	-29.24 ^c
Maine	2.38 ^b	0.17	48(32,51)	-29.48 ^c
Kansas	2.36 ^b	0.12	49(36,51)	-29.86 ^c
Montana	2.22 ^b	0.20	50(35,51)	-34.04 ^c
Utah	2.03 ^b	0.14	51(43,51)	-39.64 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	3.43	0.12	17(11,25)	1.84	Montana	2.22 ^b	0.20	50(35,51)	-34.04 ^c
Alaska	4.16	0.43	05(01,32)	23.45 ^c	Nebraska	2.40 ^b	0.16	44(32,51)	-28.82 ^c
Arizona	2.92 ^b	0.09	32(21,40)	-13.38	Nevada	2.96	0.16	31(17,43)	-12.32
Arkansas	3.08	0.14	25(16,39)	-8.57	New Hampshire	2.39 ^b	0.18	46(32,51)	-29.16 ^c
California	4.26 ^b	0.05	04(02,07)	26.48 ^c	New Jersey	3.87 ^b	0.09	10(04,15)	14.72
Colorado	2.76 ^b	0.11	36(24,46)	-18.21 ^c	New Mexico	3.95	0.19	08(02,18)	17.07 ^c
Connecticut	3.62	0.13	14(06,21)	7.27	New York	3.93 ^b	0.06	09(04,12)	16.71 ^c
Delaware	3.84	0.28	11(02,28)	13.90	North Carolina	3.16	0.08	21(17,32)	-6.34
D.C.	4.48	0.40	03(01,19)	32.80 ^c	North Dakota	3.01	0.27	27(09,49)	-10.74
Florida	2.97 ^b	0.05	30(23,36)	-11.79	Ohio	2.89 ^b	0.07	34(24,40)	-14.21
Georgia	3.60	0.09	15(08,20)	6.79	Oklahoma	3.02	0.12	26(17,39)	-10.53
Hawaii	5.26 ^b	0.26	01(01,03)	56.00 ^c	Oregon	2.55 ^b	0.11	41(32,50)	-24.49 ^c
Idaho	2.49 ^b	0.18	42(27,51)	-26.18 ^c	Pennsylvania	3.09 ^b	0.06	24(19,33)	-8.32
Illinois	3.74 ^b	0.08	12(07,17)	10.89	Rhode Island	3.44	0.23	16(05,37)	1.93
Indiana	2.70 ^b	0.09	38(29,46)	-19.89 ^c	South Carolina	3.95 ^b	0.13	07(03,15)	17.12 ^c
Iowa	2.39 ^b	0.12	45(37,51)	-29.12 ^c	South Dakota	2.60	0.24	40(19,51)	-22.82 ^c
Kansas	2.36 ^b	0.12	49(36,51)	-29.86 ^c	Tennessee	3.34	0.10	20(13,28)	-0.85
Kentucky	2.86 ^b	0.11	35(21,43)	-15.18 ^c	Texas	3.71 ^b	0.06	13(08,16)	10.14
Louisiana	4.48 ^b	0.14	02(01,07)	32.83 ^c	Utah	2.03 ^b	0.14	51(43,51)	-39.64 ^c
Maine	2.38 ^b	0.17	48(32,51)	-29.48 ^c	Vermont	2.40 ^b	0.26	43(21,51)	-28.68 ^c
Maryland	3.43	0.11	18(11,25)	1.61	Virginia	3.10	0.09	23(18,35)	-8.04
Massachusetts	3.35	0.09	19(13,26)	-0.61	Washington	3.01 ^b	0.09	28(20,38)	-10.81
Michigan	3.16	0.08	22(17,32)	-6.35	West Virginia	2.99	0.16	29(16,43)	-11.36
Minnesota	2.72 ^b	0.10	37(27,46)	-19.32 ^c	Wisconsin	2.91 ^b	0.10	33(21,41)	-13.73
Mississippi	4.03 ^b	0.16	06(02,15)	19.67 ^c	Wyoming	2.39	0.29	47(19,51)	-29.24 ^c
Missouri	2.66 ^b	0.09	39(30,47)	-21.15 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 24.19
Cancer of the Stomach (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	4.58	0.03		
High Five States				
Hawaii	7.12 ^b	0.46	01(01,05)	55.39 ^c
D.C.	6.58	0.77	02(01,22)	43.66 ^c
Louisiana	6.11 ^b	0.25	03(01,10)	33.41 ^c
South Carolina	5.79 ^b	0.23	04(02,13)	26.42 ^c
California	5.58 ^b	0.09	05(03,11)	21.91 ^c
Low Five States				
Idaho	3.24 ^b	0.31	47(26,51)	-29.35 ^c
Maine	3.20 ^b	0.29	48(28,51)	-30.19 ^c
New Hampshire	3.20 ^b	0.32	49(27,51)	-30.19 ^c
Montana	2.94 ^b	0.34	50(30,51)	-35.80 ^c
Utah	2.71 ^b	0.24	51(41,51)	-40.81 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	4.60	0.21	20(10,32)	0.41	Montana	2.94 ^b	0.34	50(30,51)	-35.80 ^c
Alaska	5.27	0.74	10(01,46)	15.01 ^c	Nebraska	3.44 ^b	0.28	41(25,51)	-25.00 ^c
Arizona	3.91 ^b	0.16	34(22,44)	-14.62	Nevada	3.81	0.26	36(18,48)	-16.90 ^c
Arkansas	4.07	0.24	28(16,45)	-11.12	New Hampshire	3.20 ^b	0.32	49(27,51)	-30.19 ^c
California	5.58 ^b	0.09	05(03,11)	21.91 ^c	New Jersey	5.31 ^b	0.16	09(04,17)	15.96 ^c
Colorado	3.76 ^b	0.20	38(23,47)	-17.94 ^c	New Mexico	5.15	0.33	14(03,26)	12.31
Connecticut	5.19	0.24	12(03,22)	13.21	New York	5.46 ^b	0.11	07(03,13)	19.23 ^c
Delaware	5.52	0.50	06(01,28)	20.49 ^c	North Carolina	4.23	0.14	24(18,37)	-7.58
D.C.	6.58	0.77	02(01,22)	43.66 ^c	North Dakota	3.88	0.46	35(08,51)	-15.28 ^c
Florida	3.92 ^b	0.08	33(25,40)	-14.54	Ohio	3.98 ^b	0.12	30(23,41)	-13.04
Georgia	4.83	0.17	16(08,25)	5.38	Oklahoma	4.41	0.23	22(12,37)	-3.80
Hawaii	7.12 ^b	0.46	01(01,05)	55.39 ^c	Oregon	3.32 ^b	0.19	44(33,51)	-27.48 ^c
Idaho	3.24 ^b	0.31	47(26,51)	-29.35 ^c	Pennsylvania	4.40	0.12	23(16,31)	-3.98
Illinois	5.16 ^b	0.14	13(05,18)	12.62	Rhode Island	5.19	0.45	11(02,34)	13.25
Indiana	3.93 ^b	0.17	31(21,44)	-14.20	South Carolina	5.79 ^b	0.23	04(02,13)	26.42 ^c
Iowa	3.35 ^b	0.21	43(30,51)	-26.86 ^c	South Dakota	3.41	0.41	42(17,51)	-25.59 ^c
Kansas	3.31 ^b	0.22	45(31,51)	-27.72 ^c	Tennessee	4.78	0.19	18(09,27)	4.26
Kentucky	3.92	0.20	32(20,45)	-14.45	Texas	4.90	0.10	15(10,20)	6.90
Louisiana	6.11 ^b	0.25	03(01,10)	33.41 ^c	Utah	2.71 ^b	0.24	51(41,51)	-40.81 ^c
Maine	3.20 ^b	0.29	48(28,51)	-30.19 ^c	Vermont	3.61	0.48	39(11,51)	-21.19 ^c
Maryland	4.82	0.20	17(08,27)	5.30	Virginia	4.22	0.16	25(17,38)	-7.85
Massachusetts	4.60	0.17	19(12,29)	0.49	Washington	4.14	0.17	27(18,40)	-9.62
Michigan	4.43	0.14	21(15,32)	-3.35	West Virginia	4.21	0.29	26(12,44)	-8.08
Minnesota	3.79 ^b	0.18	37(23,46)	-17.37 ^c	Wisconsin	4.04	0.17	29(19,42)	-11.76
Mississippi	5.34	0.29	08(02,21)	16.61 ^c	Wyoming	3.29	0.51	46(13,51)	-28.19 ^c
Missouri	3.60 ^b	0.16	40(28,48)	-21.43 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 24.20
Cancer of the Stomach (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	2.43	0.02		
High Five States				
Hawaii	3.77 ^b	0.29	01(01,07)	54.81 ^c
Louisiana	3.25 ^b	0.16	02(01,10)	33.39 ^c
California	3.24 ^b	0.06	03(01,07)	32.99 ^c
Alaska	3.13	0.51	04(01,42)	28.74 ^c
D.C.	3.08	0.44	05(01,41)	26.57 ^c
Low Five States				
Montana	1.62 ^b	0.22	47(20,51)	-33.49 ^c
Wyoming	1.60	0.32	48(11,51)	-34.21 ^c
Vermont	1.56	0.28	49(14,51)	-35.78 ^c
Nebraska	1.55 ^b	0.17	50(31,51)	-36.33 ^c
Utah	1.49 ^b	0.16	51(33,51)	-38.77 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	2.56	0.13	13(06,26)	5.13	Montana	1.62 ^b	0.22	47(20,51)	-33.49 ^c
Alaska	3.13	0.51	04(01,42)	28.74 ^c	Nebraska	1.55 ^b	0.17	50(31,51)	-36.33 ^c
Arizona	2.09	0.11	29(16,44)	-14.07	Nevada	2.25	0.18	23(08,44)	-7.36
Arkansas	2.32	0.16	20(08,40)	-4.62	New Hampshire	1.69 ^b	0.20	43(19,51)	-30.47 ^c
California	3.24 ^b	0.06	03(01,07)	32.99 ^c	New Jersey	2.80 ^b	0.10	09(04,15)	15.05 ^c
Colorado	1.95 ^b	0.13	36(18,48)	-20.03 ^c	New Mexico	2.94	0.23	07(01,22)	20.81 ^c
Connecticut	2.40	0.14	17(08,36)	-1.32	New York	2.83 ^b	0.07	08(04,13)	16.27 ^c
Delaware	2.41	0.29	16(02,49)	-0.93	North Carolina	2.33	0.09	19(12,32)	-4.29
D.C.	3.08	0.44	05(01,41)	26.57 ^c	North Dakota	2.31	0.33	21(02,50)	-5.10
Florida	2.20 ^b	0.06	25(17,34)	-9.67	Ohio	2.05 ^b	0.07	32(21,42)	-15.96 ^c
Georgia	2.68	0.11	11(05,20)	9.99	Oklahoma	1.90 ^b	0.13	40(19,50)	-22.10 ^c
Hawaii	3.77 ^b	0.29	01(01,07)	54.81 ^c	Oregon	1.93 ^b	0.13	37(18,48)	-20.54 ^c
Idaho	1.81	0.20	41(15,51)	-25.78 ^c	Pennsylvania	2.09 ^b	0.07	30(20,40)	-14.22
Illinois	2.67	0.08	12(05,18)	9.74	Rhode Island	2.16	0.24	27(06,49)	-11.07
Indiana	1.78 ^b	0.10	42(29,50)	-26.71 ^c	South Carolina	2.47	0.13	14(07,32)	1.34
Iowa	1.67 ^b	0.13	45(29,51)	-31.51 ^c	South Dakota	1.96	0.27	35(08,51)	-19.40 ^c
Kansas	1.65 ^b	0.14	46(30,51)	-32.30 ^c	Tennessee	2.29	0.11	22(12,37)	-6.02
Kentucky	2.00	0.12	34(17,47)	-17.78 ^c	Texas	2.79 ^b	0.07	10(04,14)	14.57
Louisiana	3.25 ^b	0.16	02(01,10)	33.39 ^c	Utah	1.49 ^b	0.16	51(33,51)	-38.77 ^c
Maine	1.68 ^b	0.19	44(21,51)	-30.83 ^c	Vermont	1.56	0.28	49(14,51)	-35.78 ^c
Maryland	2.38	0.12	18(09,34)	-2.33	Virginia	2.24	0.10	24(13,37)	-7.87
Massachusetts	2.41	0.11	15(10,30)	-0.90	Washington	2.09	0.11	28(16,43)	-14.01
Michigan	2.19	0.08	26(15,37)	-9.89	West Virginia	2.04	0.18	33(13,49)	-16.22 ^c
Minnesota	1.91 ^b	0.11	38(22,48)	-21.39 ^c	Wisconsin	2.05 ^b	0.11	31(17,44)	-15.67 ^c
Mississippi	3.06	0.19	06(01,15)	25.94 ^c	Wyoming	1.60	0.32	48(11,51)	-34.21 ^c
Missouri	1.90 ^b	0.10	39(24,48)	-22.06 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 25.14
Cancer of the Testis (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	0.25	0.01		
High Five States				
Oklahoma	0.40	0.07	01(01,19)	57.59 ^c
Arizona	0.37	0.05	02(01,16)	46.18 ^c
Missouri	0.33	0.05	03(01,22)	32.34 ^c
California	0.33 ^b	0.02	04(01,12)	29.34 ^c
Arkansas	0.32	0.07	05(01,28)	28.77 ^c
Low Five States				
Georgia	0.20	0.03	25(08,29)	-20.04 ^c
South Carolina	0.20	0.04	26(05,29)	-21.91 ^c
Maryland	0.18	0.04	27(09,29)	-29.26 ^c
Virginia	0.15 ^b	0.03	28(17,29)	-40.97 ^c
Massachusetts	0.14 ^b	0.03	29(18,29)	-45.22 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	0.30	0.05	06(01,27)	19.08 ^c	Montana	-	-	-(-,-)	-
Alaska	-	-	-(-,-)	-	Nebraska	-	-	-(-,-)	-
Arizona	0.37	0.05	02(01,16)	46.18 ^c	Nevada	0.23	0.06	18(01,29)	-9.29
Arkansas	0.32	0.07	05(01,28)	28.77 ^c	New Hampshire	-	-	-(-,-)	-
California	0.33 ^b	0.02	04(01,12)	29.34 ^c	New Jersey	0.24	0.03	16(04,28)	-6.60
Colorado	0.27	0.05	13(01,28)	5.29	New Mexico	-	-	-(-,-)	-
Connecticut	-	-	-(-,-)	-	New York	0.22	0.02	19(09,27)	-13.28
Delaware	-	-	-(-,-)	-	North Carolina	0.29	0.04	08(01,25)	14.72
D.C.	-	-	-(-,-)	-	North Dakota	-	-	-(-,-)	-
Florida	0.27	0.02	12(03,23)	6.98	Ohio	0.25	0.03	14(03,26)	0.63
Georgia	0.20	0.03	25(08,29)	-20.04 ^c	Oklahoma	0.40	0.07	01(01,19)	57.59 ^c
Hawaii	-	-	-(-,-)	-	Oregon	-	-	-(-,-)	-
Idaho	-	-	-(-,-)	-	Pennsylvania	0.29	0.03	07(01,22)	15.49 ^c
Illinois	0.22	0.03	21(07,28)	-13.85	Rhode Island	-	-	-(-,-)	-
Indiana	0.28	0.04	11(01,27)	10.88	South Carolina	0.20	0.04	26(05,29)	-21.91 ^c
Iowa	0.21	0.05	22(02,29)	-16.16 ^c	South Dakota	-	-	-(-,-)	-
Kansas	-	-	-(-,-)	-	Tennessee	0.22	0.04	20(04,29)	-13.45
Kentucky	0.28	0.05	09(01,28)	12.94	Texas	0.28	0.02	10(03,20)	11.92
Louisiana	0.21	0.04	23(04,29)	-17.21 ^c	Utah	-	-	-(-,-)	-
Maine	-	-	-(-,-)	-	Vermont	-	-	-(-,-)	-
Maryland	0.18	0.04	27(09,29)	-29.26 ^c	Virginia	0.15 ^b	0.03	28(17,29)	-40.97 ^c
Massachusetts	0.14 ^b	0.03	29(18,29)	-45.22 ^c	Washington	0.20	0.04	24(06,29)	-19.64 ^c
Michigan	0.24	0.03	15(04,27)	-4.06	West Virginia	-	-	-(-,-)	-
Minnesota	-	-	-(-,-)	-	Wisconsin	0.23	0.04	17(03,29)	-7.59
Mississippi	-	-	-(-,-)	-	Wyoming	-	-	-(-,-)	-
Missouri	0.33	0.05	03(01,22)	32.34 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 26.18
Cancer of the Thyroid (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	0.51	0.01		
High Five States				
Hawaii	0.76	0.10	01(01,30)	48.89 ^c
Alaska	0.74	0.18	02(01,50)	46.24 ^c
Idaho	0.70	0.10	03(01,40)	38.64 ^c
D.C.	0.66	0.15	04(01,50)	30.12 ^c
Vermont	0.63	0.13	05(01,50)	25.05 ^c
Low Five States				
Mississippi	0.42	0.05	46(09,50)	-17.29 ^c
Connecticut	0.41	0.05	47(11,50)	-18.73 ^c
South Carolina	0.40	0.04	48(18,50)	-20.57 ^c
Georgia	0.40 ^b	0.03	49(28,50)	-21.86 ^c
Montana	0.37	0.08	50(07,50)	-26.90 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	0.47	0.04	37(07,50)	-7.84	Montana	0.37	0.08	50(07,50)	-26.90 ^c
Alaska	0.74	0.18	02(01,50)	46.24 ^c	Nebraska	0.55	0.07	12(01,49)	8.31
Arizona	0.48	0.04	35(07,49)	-5.33	Nevada	0.46	0.06	40(05,50)	-9.52
Arkansas	0.50	0.06	32(04,50)	-2.40	New Hampshire	0.45	0.08	41(03,50)	-10.47
California	0.60 ^b	0.02	06(02,16)	18.44 ^c	New Jersey	0.47	0.03	36(09,49)	-6.65
Colorado	0.53	0.05	14(03,47)	5.13	New Mexico	0.55	0.07	11(01,49)	8.67
Connecticut	0.41	0.05	47(11,50)	-18.73 ^c	New York	0.53	0.02	15(06,35)	4.78
Delaware	0.56	0.11	09(01,50)	10.20	North Carolina	0.50	0.03	29(06,46)	-1.09
D.C.	0.66	0.15	04(01,50)	30.12 ^c	North Dakota	0.57	0.12	07(01,50)	11.46
Florida	0.43 ^b	0.02	43(27,49)	-14.73	Ohio	0.52	0.03	19(06,41)	2.30
Georgia	0.40 ^b	0.03	49(28,50)	-21.86 ^c	Oklahoma	0.44	0.05	42(07,50)	-12.95
Hawaii	0.76	0.10	01(01,30)	48.89 ^c	Oregon	0.49	0.05	33(04,50)	-3.41
Idaho	0.70	0.10	03(01,40)	38.64 ^c	Pennsylvania	0.52	0.03	16(06,39)	2.68
Illinois	0.50	0.03	28(07,44)	-1.06	Rhode Island	0.55	0.10	10(01,50)	8.86
Indiana	0.51	0.04	26(05,47)	-0.35	South Carolina	0.40	0.04	48(18,50)	-20.57 ^c
Iowa	0.50	0.05	30(03,49)	-1.62	South Dakota	0.50	0.10	27(01,50)	-0.62
Kansas	0.47	0.05	38(04,50)	-8.24	Tennessee	0.46	0.04	39(10,49)	-8.57
Kentucky	0.51	0.05	23(03,48)	1.13	Texas	0.56	0.02	08(04,29)	10.48
Louisiana	0.43	0.04	44(11,50)	-15.15 ^c	Utah	0.52	0.07	21(02,50)	1.66
Maine	0.50	0.08	31(02,50)	-2.20	Vermont	0.63	0.13	05(01,50)	25.05 ^c
Maryland	0.42	0.04	45(17,50)	-17.00 ^c	Virginia	0.52	0.04	20(05,45)	1.90
Massachusetts	0.53	0.04	13(04,43)	5.35	Washington	0.52	0.04	18(04,45)	2.35
Michigan	0.49	0.03	34(08,46)	-3.74	West Virginia	0.52	0.07	17(02,50)	2.45
Minnesota	0.51	0.04	24(04,48)	-0.11	Wisconsin	0.51	0.04	25(04,47)	-0.11
Mississippi	0.42	0.05	46(09,50)	-17.29 ^c	Wyoming	-	-	-(,-)	-
Missouri	0.52	0.04	22(05,47)	1.46					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 26.19
Cancer of the Thyroid (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	0.51	0.01		
High Five States				
Rhode Island	0.81	0.18	01(01,41)	59.47 ^c
Utah	0.66	0.12	02(01,41)	30.58 ^c
Idaho	0.65	0.14	03(01,42)	27.47 ^c
Virginia	0.64	0.06	04(01,27)	25.67 ^c
Kansas	0.64	0.10	05(01,40)	25.41 ^c
Low Five States				
Maryland	0.42	0.06	38(08,42)	-17.61 ^c
Georgia	0.40	0.05	39(15,42)	-21.83 ^c
Connecticut	0.38	0.07	40(07,42)	-25.01 ^c
Louisiana	0.37	0.06	41(11,42)	-26.73 ^c
Mississippi	0.34	0.07	42(12,42)	-33.75 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	0.51	0.07	20(02,41)	1.01	Montana	-	-	-(-,-)	-
Alaska	-	-	-(-,-)	-	Nebraska	0.63	0.12	06(01,41)	23.18 ^c
Arizona	0.56	0.06	10(01,37)	10.69	Nevada	0.44	0.08	33(02,42)	-13.56
Arkansas	0.43	0.08	36(04,42)	-15.27 ^c	New Hampshire	0.44	0.11	32(01,42)	-13.20
California	0.53	0.03	17(06,31)	4.68	New Jersey	0.49	0.05	23(04,40)	-2.65
Colorado	0.59	0.08	07(01,39)	16.27 ^c	New Mexico	0.44	0.09	34(02,42)	-14.16
Connecticut	0.38	0.07	40(07,42)	-25.01 ^c	New York	0.55	0.04	13(03,31)	8.67
Delaware	-	-	-(-,-)	-	North Carolina	0.54	0.05	14(02,38)	6.02
D.C.	-	-	-(-,-)	-	North Dakota	-	-	-(-,-)	-
Florida	0.45	0.03	31(16,40)	-11.83	Ohio	0.56	0.04	12(02,34)	9.42
Georgia	0.40	0.05	39(15,42)	-21.83 ^c	Oklahoma	0.56	0.08	11(01,41)	9.61
Hawaii	0.51	0.12	19(01,42)	1.17	Oregon	0.49	0.07	24(02,42)	-3.31
Idaho	0.65	0.14	03(01,42)	27.47 ^c	Pennsylvania	0.49	0.04	26(06,39)	-3.57
Illinois	0.43	0.04	35(13,42)	-14.87	Rhode Island	0.81	0.18	01(01,41)	59.47 ^c
Indiana	0.52	0.06	18(02,40)	2.44	South Carolina	0.43	0.06	37(05,42)	-15.96 ^c
Iowa	0.53	0.08	16(01,42)	4.72	South Dakota	-	-	-(-,-)	-
Kansas	0.64	0.10	05(01,40)	25.41 ^c	Tennessee	0.48	0.06	29(03,42)	-5.13
Kentucky	0.48	0.07	28(02,42)	-5.00	Texas	0.58	0.04	08(02,28)	14.06
Louisiana	0.37	0.06	41(11,42)	-26.73 ^c	Utah	0.66	0.12	02(01,41)	30.58 ^c
Maine	-	-	-(-,-)	-	Vermont	-	-	-(-,-)	-
Maryland	0.42	0.06	38(08,42)	-17.61 ^c	Virginia	0.64	0.06	04(01,27)	25.67 ^c
Massachusetts	0.57	0.06	09(01,37)	12.20	Washington	0.51	0.06	21(03,40)	0.03
Michigan	0.49	0.05	25(05,40)	-3.55	West Virginia	0.50	0.10	22(01,42)	-2.09
Minnesota	0.54	0.07	15(01,39)	5.95	Wisconsin	0.47	0.06	30(04,42)	-7.23
Mississippi	0.34	0.07	42(12,42)	-33.75 ^c	Wyoming	-	-	-(-,-)	-
Missouri	0.49	0.06	27(03,41)	-4.11					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 26.20
Cancer of the Thyroid (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	0.50	0.01		
High Five States				
Hawaii	0.95	0.15	01(01,13)	90.17 ^c
Vermont	0.86	0.21	02(01,44)	70.93 ^c
North Dakota	0.78	0.18	03(01,44)	56.23 ^c
Idaho	0.77	0.14	04(01,39)	53.27 ^c
California	0.65 ^b	0.03	05(02,11)	29.72 ^c
Low Five States				
Utah	0.40	0.09	41(04,45)	-19.83 ^c
Georgia	0.38	0.04	42(19,45)	-23.83 ^c
South Carolina	0.38	0.05	43(13,45)	-23.93 ^c
Oklahoma	0.36	0.06	44(13,45)	-28.41 ^c
Kansas	0.34	0.06	45(12,45)	-31.96 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	0.43	0.05	34(07,45)	-13.51	Montana	-	-	-(-,-)	-
Alaska	-	-	-(-,-)	-	Nebraska	0.43	0.08	37(04,45)	-14.36
Arizona	0.40	0.05	40(12,45)	-19.33 ^c	Nevada	0.45	0.08	31(04,45)	-9.63
Arkansas	0.54	0.08	12(02,44)	7.14	New Hampshire	0.43	0.10	33(03,45)	-13.45
California	0.65 ^b	0.03	05(02,11)	29.72 ^c	New Jersey	0.46	0.04	30(08,44)	-8.52
Colorado	0.47	0.06	28(05,45)	-5.66	New Mexico	0.62	0.10	06(01,43)	23.14 ^c
Connecticut	0.43	0.06	35(06,45)	-14.22	New York	0.52	0.03	17(06,34)	3.21
Delaware	0.58	0.14	08(01,45)	15.68 ^c	North Carolina	0.48	0.04	26(07,43)	-4.75
D.C.	-	-	-(-,-)	-	North Dakota	0.78	0.18	03(01,44)	56.23 ^c
Florida	0.42 ^b	0.02	39(22,44)	-17.01 ^c	Ohio	0.48	0.04	22(07,41)	-3.53
Georgia	0.38	0.04	42(19,45)	-23.83 ^c	Oklahoma	0.36	0.06	44(13,45)	-28.41 ^c
Hawaii	0.95	0.15	01(01,13)	90.17 ^c	Oregon	0.50	0.07	20(04,45)	-1.02
Idaho	0.77	0.14	04(01,39)	53.27 ^c	Pennsylvania	0.54	0.04	09(05,32)	7.86
Illinois	0.54	0.04	11(05,33)	7.20	Rhode Island	-	-	-(-,-)	-
Indiana	0.48	0.05	27(06,44)	-5.29	South Carolina	0.38	0.05	43(13,45)	-23.93 ^c
Iowa	0.46	0.07	29(04,45)	-7.76	South Dakota	-	-	-(-,-)	-
Kansas	0.34	0.06	45(12,45)	-31.96 ^c	Tennessee	0.45	0.05	32(07,45)	-11.08
Kentucky	0.53	0.06	14(03,42)	6.13	Texas	0.54	0.03	10(05,30)	7.82
Louisiana	0.48	0.06	24(05,45)	-4.36	Utah	0.40	0.09	41(04,45)	-19.83 ^c
Maine	0.58	0.11	07(01,45)	16.57 ^c	Vermont	0.86	0.21	02(01,44)	70.93 ^c
Maryland	0.42	0.05	38(10,45)	-16.53 ^c	Virginia	0.43	0.04	36(10,45)	-14.31
Massachusetts	0.50	0.05	19(05,42)	-0.60	Washington	0.52	0.05	16(04,41)	3.70
Michigan	0.48	0.04	23(07,42)	-4.26	West Virginia	0.51	0.09	18(02,45)	0.79
Minnesota	0.49	0.06	21(05,44)	-3.16	Wisconsin	0.54	0.06	13(03,40)	6.83
Mississippi	0.48	0.07	25(04,45)	-4.71	Wyoming	-	-	-(-,-)	-
Missouri	0.53	0.05	15(04,40)	5.40					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 27.18
Cancer of the Urinary Bladder (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	4.40	0.02		
High Five States				
Maine	6.12 ^b	0.27	01(01,04)	38.87 ^c
Vermont	5.69	0.40	02(01,21)	29.25 ^c
Nevada	5.32 ^b	0.21	03(01,18)	20.80 ^c
Massachusetts	5.16 ^b	0.12	04(02,14)	17.15 ^c
Ohio	5.04 ^b	0.09	05(03,16)	14.36
Low Five States				
South Dakota	3.79	0.28	47(20,50)	-13.97
Alaska	3.76	0.44	48(05,51)	-14.61
Texas	3.70 ^b	0.06	49(43,50)	-16.01 ^c
Mississippi	3.59 ^b	0.15	50(38,50)	-18.41 ^c
Hawaii	2.56 ^b	0.18	51(50,51)	-41.79 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	3.93 ^b	0.12	43(29,50)	-10.87	Montana	4.79	0.28	17(02,42)	8.75
Alaska	3.76	0.44	48(05,51)	-14.61	Nebraska	4.05	0.20	39(18,50)	-8.04
Arizona	4.12	0.11	37(25,47)	-6.44	Nevada	5.32 ^b	0.21	03(01,18)	20.80 ^c
Arkansas	4.53	0.17	21(06,41)	2.85	New Hampshire	5.00	0.26	06(02,34)	13.47
California	3.94 ^b	0.05	42(37,48)	-10.60	New Jersey	4.90 ^b	0.10	08(04,21)	11.15
Colorado	3.85 ^b	0.13	45(32,50)	-12.61	New Mexico	3.81	0.19	46(28,50)	-13.39
Connecticut	4.83	0.15	13(03,28)	9.58	New York	4.43	0.06	27(18,34)	0.68
Delaware	4.69	0.30	18(02,46)	6.47	North Carolina	4.20	0.09	33(23,44)	-4.73
D.C.	4.00	0.38	41(04,50)	-9.06	North Dakota	4.45	0.33	26(02,50)	0.97
Florida	4.47	0.06	25(17,32)	1.45	Ohio	5.04 ^b	0.09	05(03,16)	14.36
Georgia	4.37	0.10	29(17,40)	-0.75	Oklahoma	4.48	0.15	24(09,41)	1.74
Hawaii	2.56 ^b	0.18	51(50,51)	-41.79 ^c	Oregon	4.85	0.15	10(03,27)	10.12
Idaho	4.18	0.23	36(10,50)	-5.14	Pennsylvania	4.81 ^b	0.08	15(05,22)	9.32
Illinois	4.40	0.08	28(18,37)	-0.07	Rhode Island	4.90	0.28	07(02,39)	11.23
Indiana	4.80	0.12	16(04,26)	9.00	South Carolina	4.29	0.13	30(17,44)	-2.60
Iowa	4.52	0.16	23(07,40)	2.56	South Dakota	3.79	0.28	47(20,50)	-13.97
Kansas	4.09	0.16	38(20,49)	-7.21	Tennessee	4.25	0.11	32(19,44)	-3.43
Kentucky	4.82	0.15	14(03,28)	9.42	Texas	3.70 ^b	0.06	49(43,50)	-16.01 ^c
Louisiana	4.18	0.14	34(20,47)	-5.10	Utah	3.91	0.20	44(22,50)	-11.13
Maine	6.12 ^b	0.27	01(01,04)	38.87 ^c	Vermont	5.69	0.40	02(01,21)	29.25 ^c
Maryland	4.89 ^b	0.13	09(03,24)	10.98	Virginia	4.18	0.10	35(22,45)	-5.12
Massachusetts	5.16 ^b	0.12	04(02,14)	17.15 ^c	Washington	4.84 ^b	0.12	11(04,25)	9.88
Michigan	4.84 ^b	0.09	12(04,22)	9.80	West Virginia	4.53	0.20	22(05,42)	2.83
Minnesota	4.04	0.12	40(27,48)	-8.30	Wisconsin	4.65	0.12	20(06,32)	5.63
Mississippi	3.59 ^b	0.15	50(38,50)	-18.41 ^c	Wyoming	4.68	0.41	19(02,50)	6.30
Missouri	4.27	0.11	31(19,43)	-3.08					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 27.19
Cancer of the Urinary Bladder (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	7.65	0.03		
High Five States				
Maine	10.88 ^b	0.56	01(01,03)	42.19 ^c
Vermont	9.61	0.80	02(01,33)	25.49 ^c
New Hampshire	8.91	0.55	03(01,32)	16.44 ^c
Massachusetts	8.88 ^b	0.24	04(02,19)	15.96 ^c
Ohio	8.80 ^b	0.18	05(02,18)	14.92
Low Five States				
Texas	6.40 ^b	0.13	47(41,50)	-16.41 ^c
Mississippi	6.33 ^b	0.34	48(30,50)	-17.27 ^c
D.C.	5.78	0.74	49(18,51)	-24.55 ^c
Alaska	5.74	0.82	50(11,51)	-25.03 ^c
Hawaii	4.73 ^b	0.37	51(49,51)	-38.15 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	7.10	0.27	36(20,48)	-7.29	Montana	8.66	0.59	07(01,41)	13.10
Alaska	5.74	0.82	50(11,51)	-25.03 ^c	Nebraska	7.03	0.41	40(16,49)	-8.15
Arizona	7.08	0.22	38(23,47)	-7.47	Nevada	8.67	0.42	06(02,29)	13.23
Arkansas	7.81	0.35	20(04,42)	2.09	New Hampshire	8.91	0.55	03(01,32)	16.44 ^c
California	6.76 ^b	0.10	42(35,47)	-11.69	New Jersey	8.59 ^b	0.21	09(02,22)	12.23
Colorado	6.60 ^b	0.27	44(29,50)	-13.71	New Mexico	6.60	0.38	45(22,50)	-13.75
Connecticut	8.46	0.32	16(02,29)	10.49	New York	7.81	0.14	21(16,31)	1.99
Delaware	7.49	0.61	30(02,50)	-2.18	North Carolina	7.32	0.20	31(21,44)	-4.41
D.C.	5.78	0.74	49(18,51)	-24.55 ^c	North Dakota	7.71	0.66	24(02,49)	0.67
Florida	7.66	0.12	26(19,33)	0.11	Ohio	8.80 ^b	0.18	05(02,18)	14.92
Georgia	7.66	0.23	27(14,39)	0.08	Oklahoma	7.62	0.31	28(08,43)	-0.51
Hawaii	4.73 ^b	0.37	51(49,51)	-38.15 ^c	Oregon	8.47	0.31	15(02,27)	10.69
Idaho	7.26	0.47	32(08,49)	-5.21	Pennsylvania	8.59 ^b	0.16	08(03,20)	12.28
Illinois	7.73	0.17	23(15,36)	1.06	Rhode Island	8.49	0.58	12(01,41)	10.94
Indiana	8.47	0.25	14(02,25)	10.72	South Carolina	7.07	0.28	39(20,48)	-7.58
Iowa	8.17	0.33	19(03,36)	6.72	South Dakota	6.50	0.56	46(15,51)	-15.08 ^c
Kansas	6.94	0.33	41(20,49)	-9.39	Tennessee	7.68	0.25	25(12,40)	0.29
Kentucky	8.52	0.31	11(02,27)	11.30	Texas	6.40 ^b	0.13	47(41,50)	-16.41 ^c
Louisiana	7.09	0.29	37(20,49)	-7.43	Utah	6.69	0.40	43(20,50)	-12.53
Maine	10.88 ^b	0.56	01(01,03)	42.19 ^c	Vermont	9.61	0.80	02(01,33)	25.49 ^c
Maryland	8.58	0.28	10(02,25)	12.16	Virginia	7.14	0.22	34(22,46)	-6.72
Massachusetts	8.88 ^b	0.24	04(02,19)	15.96 ^c	Washington	8.31	0.25	18(03,27)	8.57
Michigan	8.48 ^b	0.20	13(03,23)	10.81	West Virginia	7.49	0.40	29(07,47)	-2.17
Minnesota	7.13	0.25	35(22,47)	-6.87	Wisconsin	8.32	0.25	17(03,28)	8.69
Mississippi	6.33 ^b	0.34	48(30,50)	-17.27 ^c	Wyoming	7.78	0.80	22(02,50)	1.68
Missouri	7.22	0.23	33(21,46)	-5.72					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 27.20
Cancer of the Urinary Bladder (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	2.19	0.02		
High Five States				
D.C.	3.02	0.43	01(01,46)	37.51 ^c
Delaware	2.84	0.31	02(01,39)	29.45 ^c
Vermont	2.81	0.37	03(01,45)	28.26 ^c
Massachusetts	2.76 ^b	0.11	04(01,15)	25.94 ^c
Maine	2.75	0.24	05(01,35)	25.57 ^c
Low Five States				
Colorado	1.84	0.12	47(28,50)	-16.24 ^c
Texas	1.84 ^b	0.06	48(36,50)	-16.30 ^c
Idaho	1.81	0.21	49(10,50)	-17.64 ^c
New Mexico	1.73	0.17	50(22,50)	-21.11 ^c
Hawaii	0.96 ^b	0.14	51(50,51)	-56.20 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	1.86	0.11	43(26,50)	-15.27 ^c	Montana	1.93	0.24	40(05,50)	-12.06
Alaska	2.06	0.46	35(01,51)	-6.23	Nebraska	2.03	0.19	37(07,50)	-7.59
Arizona	1.84 ^b	0.10	44(30,50)	-15.93 ^c	Nevada	2.70	0.20	06(01,32)	23.35 ^c
Arkansas	2.26	0.15	25(04,46)	3.19	New Hampshire	2.27	0.23	22(02,49)	3.66
California	1.96 ^b	0.04	39(32,46)	-10.50	New Jersey	2.54 ^b	0.09	08(02,27)	15.88 ^c
Colorado	1.84	0.12	47(28,50)	-16.24 ^c	New Mexico	1.73	0.17	50(22,50)	-21.11 ^c
Connecticut	2.44	0.14	13(02,37)	11.19	New York	2.26	0.06	24(12,35)	3.21
Delaware	2.84	0.31	02(01,39)	29.45 ^c	North Carolina	2.20	0.09	30(11,41)	0.46
D.C.	3.02	0.43	01(01,46)	37.51 ^c	North Dakota	2.09	0.30	33(01,50)	-4.64
Florida	2.08	0.05	34(23,42)	-5.06	Ohio	2.54 ^b	0.08	10(03,22)	15.65 ^c
Georgia	2.30	0.10	19(07,37)	4.70	Oklahoma	2.29	0.14	20(04,43)	4.53
Hawaii	0.96 ^b	0.14	51(50,51)	-56.20 ^c	Oregon	2.22	0.13	29(06,45)	1.04
Idaho	1.81	0.21	49(10,50)	-17.64 ^c	Pennsylvania	2.32	0.07	18(08,34)	5.95
Illinois	2.23	0.08	27(11,39)	1.51	Rhode Island	2.60	0.26	07(01,43)	18.62 ^c
Indiana	2.34	0.11	17(05,37)	6.88	South Carolina	2.45	0.13	11(02,35)	11.85
Iowa	2.04	0.14	36(13,50)	-7.17	South Dakota	1.90	0.26	41(04,50)	-13.13
Kansas	2.15	0.16	32(06,49)	-2.06	Tennessee	2.00	0.10	38(20,49)	-8.91
Kentucky	2.37	0.13	16(03,39)	7.95	Texas	1.84 ^b	0.06	48(36,50)	-16.30 ^c
Louisiana	2.29	0.13	21(05,43)	4.22	Utah	1.84	0.18	46(14,50)	-16.06 ^c
Maine	2.75	0.24	05(01,35)	25.57 ^c	Vermont	2.81	0.37	03(01,45)	28.26 ^c
Maryland	2.54	0.12	09(02,29)	15.76 ^c	Virginia	2.22	0.10	28(09,42)	1.11
Massachusetts	2.76 ^b	0.11	04(01,15)	25.94 ^c	Washington	2.37	0.11	15(04,36)	8.27
Michigan	2.40	0.09	14(05,32)	9.33	West Virginia	2.44	0.19	12(01,43)	11.48
Minnesota	1.90	0.11	42(27,50)	-13.28	Wisconsin	2.16	0.11	31(10,45)	-1.44
Mississippi	1.84	0.14	45(21,50)	-15.99 ^c	Wyoming	2.25	0.39	26(01,50)	2.81
Missouri	2.27	0.11	23(06,39)	3.61					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.