

Table 1.20
U.S. and SEER Death Rates by Primary Cancer Site and Race/Ethnicity, 2013-2017

Site		Total United States ^a							SEER 21 Areas ^{ab}						
		Total	White	Black	AI/AN ^c	API ^d	Hisp ^e	W-NHisp ^e	Total	White	Black	AI/AN ^c	API ^d	Hisp ^e	W-NHisp ^e
All Sites	Both Sexes	158.3	159.0	181.7	144.2	98.9	112.3	162.9	151.9	155.4	170.3	123.7	102.4	112.4	159.6
	Male	189.5	189.6	227.3	173.1	117.3	136.2	193.8	180.5	183.6	211.3	149.5	122.7	134.9	188.1
	Female	135.7	136.4	153.4	123.0	85.6	95.5	139.9	131.7	135.1	145.4	104.8	87.7	97.0	138.9
Oral Cavity & Pharynx	Both Sexes	2.5	2.5	2.7	2.2	2.0	1.5	2.6	2.4	2.4	2.5	1.8	2.1	1.6	2.5
	Male	3.9	3.9	4.5	3.5	3.1	2.4	4.0	3.7	3.8	4.0	3.2	3.3	2.5	3.9
	Female	1.3	1.4	1.3	1.1	1.1	0.8	1.4	1.3	1.3	1.3	-	1.2	0.8	1.4
Esophagus	Both Sexes	3.9	4.2	3.2	3.4	1.6	2.0	4.4	3.6	4.0	3.0	3.0	1.6	2.1	4.2
	Male	7.0	7.4	5.3	6.2	2.8	3.7	7.8	6.5	7.0	5.0	4.9	2.9	3.8	7.5
	Female	1.4	1.4	1.6	1.3	0.7	0.7	1.5	1.4	1.4	1.5	1.4	0.6	0.8	1.5
Stomach	Both Sexes	3.1	2.7	5.3	4.8	5.2	5.0	2.3	3.5	3.0	5.2	5.5	5.4	5.4	2.5
	Male	4.1	3.6	7.8	6.3	6.7	6.4	3.2	4.7	4.1	7.7	7.4	7.1	6.8	3.5
	Female	2.2	1.9	3.6	3.6	4.1	4.0	1.6	2.5	2.2	3.5	3.9	4.2	4.4	1.7
Colon & Rectum	Both Sexes	13.9	13.6	18.5	15.8	9.6	11.1	13.8	13.4	13.3	17.2	14.7	10.0	10.8	13.4
	Male	16.6	16.2	23.2	19.4	11.5	14.2	16.3	15.8	15.6	21.3	17.3	12.2	13.7	15.7
	Female	11.8	11.5	15.2	13.0	8.1	8.7	11.7	11.4	11.3	14.3	12.7	8.3	8.7	11.6
Liver & Intrahepatic Bile Duct	Both Sexes	6.6	6.2	8.4	10.6	9.1	9.3	5.8	6.8	6.3	8.0	9.2	9.3	9.5	5.7
	Male	9.6	9.0	13.2	14.8	13.5	13.3	8.4	10.0	9.2	12.7	12.2	13.9	13.6	8.3
	Female	4.0	3.8	4.8	7.1	5.6	6.0	3.5	4.1	3.9	4.6	6.5	5.6	6.2	3.5
Pancreas	Both Sexes	11.0	10.9	13.3	8.9	7.6	8.5	11.1	10.9	11.1	12.3	7.7	8.0	8.6	11.3
	Male	12.7	12.7	14.9	10.3	8.2	9.4	13.0	12.4	12.7	13.9	9.4	8.8	9.4	13.0
	Female	9.6	9.4	12.0	7.8	7.1	7.8	9.6	9.6	9.7	11.2	6.3	7.5	7.9	9.8
Larynx	Both Sexes	1.0	0.9	1.6	0.9	0.3	0.7	1.0	0.9	0.9	1.4	0.6	0.3	0.6	0.9
	Male	1.7	1.7	3.0	1.7	0.6	1.4	1.7	1.5	1.5	2.7	1.3	0.6	1.3	1.5
	Female	0.4	0.4	0.5	-	0.1	0.1	0.4	0.3	0.3	0.5	-	0.1	0.1	0.4
Lung & Bronchus	Both Sexes	40.2	41.0	42.3	33.4	22.2	17.6	43.4	36.1	37.4	37.8	24.1	23.0	16.6	40.2
	Male	49.3	49.4	58.8	40.1	29.3	24.2	51.8	43.6	44.1	52.1	31.0	31.1	22.2	46.8
	Female	33.2	34.5	31.1	28.4	16.9	12.7	36.8	30.4	32.4	28.3	18.9	16.9	12.6	35.1
Melanoma of the Skin	Both Sexes	2.4	2.8	0.3	0.7	0.3	0.7	3.0	2.2	2.7	0.3	0.6	0.3	0.6	3.1
	Male	3.5	4.1	0.4	1.0	0.4	0.9	4.5	3.3	4.0	0.4	-	0.4	0.9	4.5
	Female	1.5	1.7	0.3	0.5	0.3	0.5	1.9	1.4	1.7	0.2	-	0.3	0.5	1.9
Breast	Female	20.3	19.8	27.6	14.6	11.4	14.0	20.3	19.9	19.9	26.7	13.3	12.1	14.1	20.5

^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b The SEER 21 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, Georgia excluding ATL/RG, Idaho, New York and Massachusetts.

^c Rates for American Indian/Alaska Native (AI/AN) are based on the Purchased/Referred Care Delivery Area (PRCDA) counties.

^d Asian/Pacific Islander.

^e Hispanic (Hisp) and White Non-Hispanic (W-NHisp) are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

- Statistic could not be calculated due to less than 16 cases in the time interval.

Table 1.20 - continued
 U.S. and SEER Death Rates by Primary Cancer Site and Race/Ethnicity, 2013-2017

Site		Total United States ^a							SEER 21 Areas ^{ab}						
		Total	White	Black	AI/AN ^c	API ^d	Hisp ^e	W-NHisp ^e	Total	White	Black	AI/AN ^c	API ^d	Hisp ^e	W-NHisp ^e
Cervix	Female	2.3	2.2	3.4	2.5	1.8	2.6	2.1	2.1	2.0	3.2	2.0	1.8	2.5	1.9
Corpus & Uterus, NOS	Female	4.8	4.5	8.6	3.4	3.2	4.0	4.5	5.1	4.8	8.7	3.1	3.3	4.2	4.7
Ovary	Female	6.9	7.1	6.0	6.3	4.4	5.2	7.3	7.0	7.4	5.9	5.1	4.5	5.4	7.6
Prostate	Male	19.1	17.9	37.9	18.7	8.6	15.8	18.0	19.3	18.6	36.9	16.2	9.0	16.6	18.7
Testis	Male	0.3	0.3	0.1	0.4	0.1	0.3	0.3	0.3	0.3	0.1	-	0.1	0.3	0.3
Urinary Bladder	Both Sexes	4.4	4.6	3.5	2.7	1.7	2.3	4.8	4.3	4.6	3.4	2.8	1.8	2.3	4.9
	Male	7.5	8.0	5.3	4.2	2.8	3.9	8.3	7.3	8.0	5.1	4.1	3.0	3.8	8.5
	Female	2.1	2.2	2.4	1.7	0.9	1.3	2.3	2.1	2.2	2.4	1.8	0.9	1.3	2.4
Kidney & Renal Pelvis	Both Sexes	3.7	3.8	3.6	5.5	1.7	3.5	3.8	3.4	3.6	3.1	4.6	1.9	3.2	3.6
	Male	5.4	5.6	5.4	8.2	2.6	5.0	5.6	5.1	5.3	5.0	6.6	2.8	4.6	5.3
	Female	2.3	2.4	2.2	3.4	1.1	2.2	2.4	2.1	2.3	1.8	3.1	1.2	2.1	2.3
Brain & Nervous System	Both Sexes	4.4	4.8	2.6	2.5	2.2	3.0	5.1	4.3	4.8	2.5	2.3	2.3	3.0	5.1
	Male	5.4	5.8	3.2	2.8	2.6	3.5	6.1	5.3	5.9	3.1	2.8	2.6	3.6	6.2
	Female	3.6	3.9	2.2	2.4	1.9	2.6	4.1	3.5	3.9	2.0	1.9	2.0	2.6	4.1
Thyroid	Both Sexes	0.5	0.5	0.5	0.5	0.6	0.6	0.5	0.6	0.6	0.5	-	0.6	0.7	0.5
	Male	0.5	0.5	0.4	0.6	0.5	0.6	0.5	0.6	0.6	0.4	-	0.5	0.7	0.6
	Female	0.5	0.5	0.6	0.4	0.7	0.7	0.5	0.6	0.5	0.5	-	0.7	0.7	0.5
Hodgkin Lymphoma	Both Sexes	0.3	0.3	0.3	0.2	0.1	0.4	0.3	0.3	0.3	0.2	-	0.2	0.4	0.3
	Male	0.4	0.4	0.3	-	0.2	0.5	0.4	0.4	0.4	0.3	-	0.2	0.5	0.4
	Female	0.2	0.2	0.2	-	0.1	0.3	0.2	0.2	0.3	0.2	-	0.1	0.3	0.2
Non-Hodgkin Lymphoma	Both Sexes	5.5	5.7	4.0	4.2	3.8	4.7	5.8	5.4	5.8	3.9	3.9	3.8	4.8	5.8
	Male	7.1	7.4	5.2	5.5	4.9	5.9	7.5	7.0	7.5	5.1	4.8	4.9	6.0	7.5
	Female	4.2	4.4	3.2	3.2	3.0	3.7	4.5	4.2	4.4	3.1	3.2	3.0	3.8	4.4
Myeloma	Both Sexes	3.3	3.0	6.2	3.2	1.6	2.7	3.0	3.2	3.0	5.7	3.0	1.6	2.7	3.0
	Male	4.1	3.9	7.5	3.6	2.0	3.4	3.9	4.0	3.9	6.9	2.9	2.0	3.3	3.9
	Female	2.6	2.4	5.3	3.0	1.2	2.2	2.3	2.5	2.4	5.0	3.2	1.2	2.3	2.3
Leukemia	Both Sexes	6.4	6.7	5.4	4.0	3.6	4.6	6.8	6.2	6.6	5.1	3.3	3.7	4.7	6.7
	Male	8.6	8.9	7.0	5.1	4.8	5.8	9.1	8.2	8.7	6.6	4.7	5.0	5.8	8.8
	Female	4.8	5.0	4.3	3.0	2.6	3.8	5.0	4.7	5.0	4.2	2.2	2.8	3.8	5.0

^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b The SEER 21 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, Georgia excluding ATL/RG, Idaho, New York and Massachusetts.

^c Rates for American Indian/Alaska Native (AI/AN) are based on the Purchased/Referred Care Delivery Area (PRCDA) counties.

^d Asian/Pacific Islander.

^e Hispanic (Hisp) and White Non-Hispanic (W-NHisp) are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

- Statistic could not be calculated due to less than 16 cases in the time interval.