

Table XX-1
OVARY CANCER (Invasive)

TRENDS IN SEER INCIDENCE AND U.S. MORTALITY, BY RACE AND AGE

	All Races, Females			White Females			Black Females		
	All	<50	50+	All	<50	50+	All	<50	50+
<u>INCIDENCE TRENDS (expressed in percents)§:</u>									
<u>1973-93</u>									
All Ages									
Percent Change (PC)	3.1	6.0	2.1	3.6	7.2	2.4	3.8	53.7	-5.8
Est. Annual PC (EAPC)	0.5★	0.4	0.5★	0.6★	0.6	0.5★	0.3	0.3	0.4
Under 65									
Percent Change (PC)	-3.4	-	-	-4.6	-	-	34.7	-	-
Est. Annual PC (EAPC)	0.0	-	-	0.1	-	-	0.2	-	-
65 and over									
Percent Change (PC)	15.8	-	-	20.3	-	-	-25.2	-	-
Est. Annual PC (EAPC)	1.3★	-	-	1.4★	-	-	0.6	-	-
<u>1975-79 - ALL AGES</u>									
Percent Change (PC)	-4.9	-6.8	-4.2	-5.0	-5.2	-5.0	-2.9	-6.6	-1.7
Est. Annual PC (EAPC)	-1.6★	-2.3	-1.4★	-1.7★	-1.6	-1.7	-0.3	-2.2	0.4
<u>1989-93 - ALL AGES</u>									
Percent Change (PC)	-2.5	-1.4	-2.9	-3.8	-1.9	-4.4	2.3	26.8	-3.5
Est. Annual PC (EAPC)	-0.8◇	-1.0	-0.8	-1.3★	-1.3	-1.3	0.7	5.7	-0.7
<u>MORTALITY TRENDS (expressed in percents)δ:</u>									
<u>1973-93</u>									
All Ages									
Percent Change (PC)	-8.7	-45.5	-1.2	-8.1	-44.9	-0.7	-5.7	-48.3	3.1
Est. Annual PC (EAPC)	-0.5★	-3.0★	-0.1	-0.4★	-3.0★	-0.1	-0.2	-3.0★	0.3
Under 65									
Percent Change (PC)	-29.8	-	-	-29.0	-	-	-32.7	-	-
Est. Annual PC (EAPC)	-1.8★	-	-	-1.8★	-	-	-2.0★	-	-
65 and over									
Percent Change (PC)	20.4	-	-	20.6	-	-	32.5	-	-
Est. Annual PC (EAPC)	1.0★	-	-	0.9★	-	-	1.8★	-	-
<u>1975-79 - ALL AGES</u>									
Percent Change (PC)	-5.1	-16.6	-3.0	-5.4	-17.6	-3.1	-2.1	-8.7	-0.9
Est. Annual PC (EAPC)	-1.6★	-5.6★	-0.9	-1.8★	-6.2★	-1.0	0.0	-1.7	0.3
<u>1989-93 - ALL AGES</u>									
Percent Change (PC)	-0.5	-5.1	0.0	-1.1	-5.8	-0.5	4.0	-5.2	5.1
Est. Annual PC (EAPC)	-0.4	-1.4◆	-0.2	-0.5	-1.5◆	-0.4	1.1	-2.6	1.5

The Est. Annual PC is the Estimated Annual Percent Change (EAPC) over the time interval.

§ SEER Program.

δ NCHS public use tape.

★ The EAPC is significantly different from zero ($p < .05$).

◆ The EAPC for 1989-93 is significantly different from the EAPC for 1975-79 ($p < .05$).

◇ The EAPC for 1989-93 is significantly different from the EAPC for 1975-79 ($p < .10$).

- Statistic could not be calculated.

Table XX-2
OVARY CANCER (Invasive)

SEER INCIDENCES AND U.S. MORTALITY^δ RATES, AGE-ADJUSTED AND AGE-SPECIFIC RATES, BY RACE

	<u>Incidences</u>			<u>US Mortality^δ</u>		
	<u>All Races, Females</u>	<u>White Females</u>	<u>Black Females</u>	<u>All Races, Females</u>	<u>White Females</u>	<u>Black Females</u>
<u>AGE-SPECIFIC RATES, 1989-93</u>						
AGE AT DIAGNOSIS/DEATH:						
0-4	0.0	0.0	0.0	0.0	0.0	0.0
5-9	0.2	0.2	0.0	0.0	0.0	0.0
10-14	0.6	0.4	0.9	0.0	0.0	0.1
15-19	2.4	2.2	2.2	0.1	0.1	0.1
20-24	2.9	2.8	2.6	0.2	0.2	0.2
25-29	4.9	5.3	3.9	0.3	0.3	0.3
30-34	6.7	7.0	2.9	0.7	0.7	0.7
35-39	10.1	10.9	5.8	1.6	1.6	1.6
40-44	13.2	14.4	6.8	3.3	3.5	2.8
45-49	21.8	23.9	10.9	7.1	7.5	5.3
50-54	28.8	29.4	23.3	12.4	12.9	9.5
55-59	39.4	41.9	22.4	18.6	19.5	14.0
60-64	45.1	48.3	29.7	26.1	27.2	20.5
65-69	53.7	56.1	46.8	35.0	35.9	31.5
70-74	58.9	62.4	44.2	42.8	43.8	37.6
75-79	65.1	68.7	47.8	48.4	49.6	41.2
80-84	64.8	67.1	43.7	53.5	54.5	47.7
85+	59.0	60.3	54.0	52.8	53.2	51.4
<u>AGE-ADJUSTED RATES, 1989-93</u>						
AGE AT DIAGNOSIS/DEATH:						
All ages	15.1	15.9	10.5	7.8	8.0	6.5
Under 65	10.3	10.9	6.6	3.9	4.0	3.0
65 and over	58.9	61.8	46.5	43.1	44.1	38.4
All ages (world std.) [⊕]	12.9	13.6	8.9	6.2	6.3	5.1

§ SEER Program. Rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population, except where noted.

δ NCHS public use tape. Rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population, except where noted.

⊕ Rates are per 100,000 and are age-adjusted to the world standard population.

Table XX-3
OVARY CANCER (Invasive)

SEER INCIDENCE§ AND U.S. MORTALITYδ RATES, AGE-ADJUSTED RATES, BY RACE AND AGE

	All Races, Females			White Females			Black Females		
	All	<50	50+	All	<50	50+	All	<50	50+
<u>SEER INCIDENCE RATES</u>									
YEAR OF DIAGNOSIS:									
1973	14.1	4.7	43.1	14.7	5.0	44.6	10.5	2.3	36.0
1974	14.7	4.9	44.9	15.3	5.1	46.7	10.2	2.1	35.1
1975	14.1	4.9	42.3	14.4	4.9	43.9	10.1	3.7	29.8
1976	13.7	4.4	42.3	14.3	4.5	44.5	9.4	2.9	29.8
1977	13.6	4.6	41.1	14.1	4.7	42.9	9.6	3.8	27.5
1978	13.2	4.2	41.1	13.8	4.3	43.1	8.5	2.8	26.0
1979	13.2	4.5	40.0	13.5	4.6	40.9	10.5	3.3	32.5
1980	13.3	4.1	41.8	14.0	4.2	44.0	10.0	2.4	33.4
1981	13.2	4.1	41.4	13.7	4.2	43.1	9.9	2.6	32.5
1982	13.4	3.9	42.5	13.8	4.1	43.9	10.8	3.3	33.9
1983	13.7	4.3	42.9	14.1	4.3	44.3	11.5	2.9	38.1
1984	14.0	4.4	43.7	14.8	4.6	46.1	9.4	2.5	30.6
1985	14.3	4.3	45.1	15.0	4.5	47.4	10.1	3.1	31.7
1986	12.9	3.8	41.0	13.5	4.1	42.4	9.3	1.1	34.7
1987	13.9	3.8	44.9	14.7	4.0	47.4	10.1	2.9	32.2
1988	14.9	4.9	45.7	15.6	5.2	47.8	10.8	4.0	31.6
1989	15.3	5.3	46.1	16.2	5.7	48.8	10.7	2.9	35.0
1990	15.2	5.1	46.5	16.1	5.4	49.0	10.3	2.5	34.4
1991	15.3	5.1	46.9	16.2	5.4	49.8	9.9	2.9	31.5
1992	14.9	5.3	44.6	15.7	5.6	46.5	10.6	3.7	31.8
1993	14.8	4.9	45.3	15.4	5.2	47.0	11.0	3.1	35.1
1989-93	15.1	5.1	45.9	15.9	5.4	48.2	10.5	3.0	33.6
<u>U.S. MORTALITY RATES</u>									
YEAR OF DEATH:									
1973	8.4	1.9	28.4	8.5	1.9	29.0	7.1	1.6	24.0
1974	8.5	1.9	28.8	8.7	2.0	29.5	6.8	1.6	23.1
1975	8.4	1.8	28.9	8.7	1.8	29.7	6.3	1.3	21.4
1976	8.5	1.7	29.6	8.7	1.7	30.4	6.9	1.3	24.1
1977	8.2	1.5	28.9	8.4	1.5	29.8	6.3	1.3	21.9
1978	8.2	1.5	28.8	8.4	1.5	29.6	6.3	1.1	22.6
1979	7.9	1.4	27.9	8.1	1.4	28.6	6.5	1.4	22.4
1980	7.8	1.3	27.9	8.0	1.4	28.6	6.7	1.2	23.7
1981	7.8	1.3	27.9	8.0	1.3	28.7	6.2	1.0	22.3
1982	7.8	1.3	27.9	8.0	1.3	28.7	6.3	0.9	22.8
1983	7.8	1.2	28.1	8.0	1.2	28.9	6.5	1.1	22.8
1984	7.7	1.2	27.4	7.9	1.3	28.2	6.3	1.0	22.8
1985	7.7	1.2	27.6	7.9	1.2	28.4	6.3	0.9	22.8
1986	7.8	1.2	27.9	8.0	1.3	28.7	6.3	0.9	22.9
1987	7.7	1.1	28.1	8.0	1.1	29.0	6.3	1.0	22.7
1988	7.8	1.1	28.5	8.0	1.1	29.3	6.6	1.0	24.0
1989	7.7	1.1	28.2	8.0	1.1	29.2	6.1	0.9	22.4
1990	7.8	1.1	28.3	8.0	1.2	29.2	6.5	0.9	23.9
1991	7.9	1.1	29.1	8.1	1.1	30.0	6.9	1.0	24.8
1992	7.9	1.0	28.9	8.1	1.1	29.7	6.8	0.9	25.0
1993	7.5	1.0	27.6	7.7	1.1	28.4	6.3	0.7	23.6
1989-93	7.8	1.1	28.4	8.0	1.1	29.3	6.5	0.9	23.9

§ SEER Program. Rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population.

δ NCHS public use tape. Rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population.

Table XX-4
OVARY CANCER (Invasive)

SURVIVAL RATES, BY RACE, DIAGNOSIS YEAR, STAGE AND AGE

	All Races, Females			White Females			Black Females		
	All	<50	50+	All	<50	50+	All	<50	50+
<u>5-YR RELATIVE SURVIVAL RATES</u>									
YEAR OF DIAGNOSIS:									
1960-63 ^a	-	-	-	32	-	-	32	-	-
1970-73 ^a	-	-	-	36	-	-	32	-	-
1974-76 ^b	36.6	56.1	29.8	36.3	55.9	29.9	40.4	61.0 [†]	29.7
1977-79 ^b	38.3	62.7	29.7	37.7	61.9	29.7	40.0	67.2 [†]	24.3
1980-82 ^b	39.0	64.6	31.2	38.8	64.0	31.5	38.6	69.8 [†]	26.2
1983-85 ^b	40.9	65.0	33.1	40.3	63.4	33.2	41.4	79.7	26.0
1986-92 ^b	45.5 Ω	73.0 Ω	35.3 Ω	45.6 Ω	72.7 Ω	35.8 Ω	40.0	74.4 Ω	25.0
1986-92 ^b									
ALL STAGES	45.5	73.0	35.3	45.6	72.7	35.8	40.0	74.4	25.0
LOCALIZED	91.5	94.5	89.3	91.8	94.6	89.9	83.8	90.8	75.2 [†]
REGIONAL	51.1	73.2	42.8	51.2	73.2	43.4	50.9 [†]	93.5 [†]	30.5 [†]
DISTANT	24.5	47.3	19.3	24.4	46.5	19.5	23.7	54.1 [†]	16.3
UNSTAGED	28.1	67.0 [†]	21.7	27.4	67.7 [†]	21.9	26.8 [†]	-	17.8 [†]

STAGE DISTRIBUTION (%) 1986-92^b

ALL STAGES

Number of cases	11,924	3,038	8,886	10,652	2,642	8,010	730	212	518
Percent	100%	100%	100%	100%	100%	100%	100%	100%	100%
LOCALIZED	24	46	17	24	46	17	21	43	12
REGIONAL	14	14	14	14	14	14	10	12	9
DISTANT	57	37	63	57	38	63	62	40	70
UNSTAGED	6	3	7	6	2	7	7	5	8

5-YR RELATIVE SURVIVAL RATES, 1986-92^b

AGE AT DIAGNOSIS:

<45	77.7	-	-	77.5	-	-	76.7	-	-
45-54	57.4	-	-	58.0	-	-	48.4 [†]	-	-
55-64	40.8	-	-	42.2	-	-	20.1	-	-
65-74	31.6	-	-	32.0	-	-	25.1	-	-
75+	22.4	-	-	22.4	-	-	18.2 [†]	-	-
Under 65	57.5	-	-	57.9	-	-	51.6	-	-
65 and over	28.1	-	-	28.3	-	-	22.8	-	-

^a Rates are based on End Results data from a series of hospital registries and one population-based registry.

^b Rates are from the SEER Program. They are based on data from population-based registries in Connecticut, New Mexico, Utah, Iowa, Hawaii, Atlanta, Detroit, Seattle-Puget Sound and San Francisco-Oakland. Rates are based on follow-up of patients through 1993.

Ω The difference in rates between 1974-76 and 1986-92 is statistically significant ($p < .05$).

[†] The standard error of the survival rate is between 5 and 10 percentage points.

^{††} The standard error of the survival rate is greater than 10 percentage points.

- Statistic could not be calculated.

Table XX-5
OVARY CANCER (Invasive)

INCIDENCE, MORTALITY, AND SURVIVAL RATES

By Year of Diagnosis/Death

All Races, Females

Year of Diagnosis/Death

	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993
Incidence																					
SEER	14.1	14.7	14.1	13.7	13.6	13.2	13.2	13.3	13.2	13.4	13.7	14.0	14.3	12.9	13.9	14.9	15.3	15.2	15.3	14.9	14.8
Mortality																					
U.S.	8.4	8.5	8.4	8.5	8.2	8.2	7.9	7.8	7.8	7.8	7.8	7.7	7.7	7.8	7.7	7.8	7.7	7.8	7.9	7.9	7.5
SEER	8.8	8.7	8.7	9.1	8.1	8.2	8.2	7.9	7.7	7.8	7.9	7.2	7.9	8.1	7.7	8.3	8.0	7.5	7.9	7.8	7.7
Relative Survival Rates (SEER)																					
1-year	59.9	64.4	59.6	63.7	66.2	67.7	67.5	69.8	69.3	70.6	70.1	73.1	72.1	69.1	71.0	73.6	75.5	76.1	76.5	78.3	
2-year	45.7	48.7	45.0	47.8	51.0	52.0	51.5	53.9	52.5	52.7	53.8	56.2	55.5	52.4	53.9	62.0	62.1	64.0	64.3		
3-year	40.8	43.5	39.7	42.2	43.0	44.5	44.1	46.6	45.7	44.6	47.8	48.3	48.0	46.0	46.3	55.5	54.4	56.4			
4-year	38.1	40.8	36.5	39.4	40.1	40.5	40.1	41.9	41.9	40.9	43.7	43.2	42.6	42.2	42.3	51.4	49.4				
5-year	36.0	38.1	34.1	37.8	38.7	38.4	37.7	39.0	40.0	38.1	42.5	40.6	39.6	38.5	38.8	48.3					
6-year	35.2	35.8	33.9	36.6	37.8	36.4	36.2	37.0	38.5	36.8	40.0	39.6	37.5	36.7	37.3						
7-year	34.9	35.4	33.3	35.3	36.4	35.8	34.7	36.6	38.5	35.8	39.0	38.8	36.2	35.6							
8-year	33.7	34.7	32.5	34.4	36.0	35.1	34.0	35.5	37.6	35.2	37.2	37.8	35.7								
9-year	33.1	34.1	32.2	34.4	35.8	34.6	33.6	34.1	36.9	35.0	37.0	36.4									
10-year	32.5	34.1	32.2	34.4	35.8	34.6	33.4	33.5	36.3	34.8	36.3										
11-year	32.5	34.0	32.2	34.0	35.7	34.3	33.4	33.5	36.3	34.5											
12-year	32.5	33.7	32.2	33.7	35.1	33.7	33.4	33.0	36.3												
13-year	32.1	33.3	32.2	33.7	35.1	33.2	33.4	32.2													
14-year	31.9	33.2	32.2	33.4	34.8	33.0	33.4														
15-year	31.6	33.1	32.2	32.2	34.8	32.8															
16-year	30.5	33.1	32.2	31.6	34.8																
17-year	30.2	33.1	32.2	31.1																	
18-year	30.2	33.1	32.2																		
19-year	30.1	33.0																			
20-year	30.1																				

Note: Incidence and mortality rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population.

Survival rates are relative rates expressed as percents.

Table XX-6
OVARY CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
 Given Cancer Free At Current Age By Race

Lifetime Risk (Percent) of Being Diagnosed With Cancer and
 Lifetime Risk (Percent) of Dying From Cancer

SEER Areas, 1991-93

All Races

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.01	0.05	1.81
10	0.01	0.05	0.14	1.82
20	0.04	0.12	0.29	1.82
30	0.09	0.25	0.58	1.79
40	0.17	0.50	0.92	1.72
50	0.34	0.77	1.21	1.58
60	0.46	0.92	1.25	1.31

Lifetime Risk of Being Diagnosed = 1.81%
 Lifetime Risk of Dying = 1.12%

Whites

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.01	0.05	1.92
10	0.01	0.05	0.14	1.93
20	0.04	0.13	0.31	1.92
30	0.09	0.28	0.62	1.89
40	0.19	0.53	0.98	1.82
50	0.35	0.81	1.28	1.66
60	0.48	0.98	1.32	1.38

Lifetime Risk of Being Diagnosed = 1.92%
 Lifetime Risk of Dying = 1.19%

Blacks

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.02	0.05	1.10
10	0.02	0.05	0.09	1.13
20	0.03	0.08	0.17	1.11
30	0.05	0.14	0.35	1.10
40	0.10	0.31	0.60	1.07
50	0.23	0.52	0.78	1.02
60	0.33	0.61	0.80	0.88

Lifetime Risk of Being Diagnosed = 1.10%
 Lifetime Risk of Dying = 0.81%

Table XX-7
OVARY CANCER (Invasive)AVERAGE ANNUAL AGE-ADJUSTED CANCER MORTALITY^δ RATES BY STATE, 1989-93

All Races, Females

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
TOTAL U.S.	7.8	0.03			Montana	7.6	0.56	(29)	-2.6
High Five States					Nebraska	7.6	0.39	(28)	-2.6
North Dakota	9.3 [*]	0.69	(01)	19.2	Nevada	6.4 [*]	0.43	(48)	-17.9
Vermont	9.2 [*]	0.76	(02)	17.9	New Hampshire	8.9	0.54	(04)	14.1
New Jersey	9.1 ^{⊕*}	0.19	(03)	16.7	New Jersey	9.1 ^{⊕*}	0.19	(03)	16.7
New Hampshire	8.9	0.54	(04)	14.1	New Mexico	6.7	0.40	(47)	-14.1
Washington	8.8	0.25	(05)	12.8	New York	8.4	0.12	(09)	7.7
Low Five States					North Carolina	7.1	0.18	(39)	-9.0
New Mexico	6.7	0.40	(47)	-14.1	North Dakota	9.3 [*]	0.69	(01)	19.2
Nevada	6.4 [*]	0.43	(48)	-17.9	Ohio	7.9	0.15	(19)	1.3
Louisiana	6.3 ^{⊕*}	0.23	(49)	-19.2	Oklahoma	7.0	0.26	(42)	-10.3
Mississippi	6.3 ^{⊕*}	0.28	(50)	-19.2	Oregon	8.4	0.30	(10)	7.7
Hawaii	5.8 [*]	0.44	(51)	-25.6	Pennsylvania	8.5	0.14	(07)	9.0
					Rhode Island	7.4	0.47	(34)	-5.1
					South Carolina	7.7	0.27	(25)	-1.3
					South Dakota	7.7	0.60	(24)	-1.3
					Tennessee	6.9	0.21	(45)	-11.5
					Texas	6.9 [⊕]	0.12	(44)	-11.5
					Utah	7.6	0.44	(31)	-2.6
					Vermont	9.2 [*]	0.76	(02)	17.9
					Virginia	7.5	0.21	(32)	-3.8
					Washington	8.8	0.25	(05)	12.8
					West Virginia	7.3	0.35	(37)	-6.4
					Wisconsin	8.4	0.24	(08)	7.7
					Wyoming	8.0	0.81	(16)	2.6
Alabama	7.1	0.23	(38)	-9.0					
Alaska	7.0	0.98	(43)	-10.3					
Arizona	7.1	0.25	(41)	-9.0					
Arkansas	7.8	0.31	(21)	0.0					
California	7.8	0.10	(22)	0.0					
Colorado	7.4	0.29	(36)	-5.1					
Connecticut	7.6	0.27	(27)	-2.6					
Delaware	7.9	0.62	(20)	1.3					
District of Columbia	6.9	0.60	(46)	-11.5					
Florida	7.7	0.13	(26)	-1.3					
Georgia	7.1	0.20	(40)	-9.0					
Hawaii	5.8 [*]	0.44	(51)	-25.6					
Idaho	8.2	0.52	(12)	5.1					
Illinois	8.1	0.15	(14)	3.8					
Indiana	7.9	0.21	(18)	1.3					
Iowa	8.2	0.30	(11)	5.1					
Kansas	7.7	0.32	(23)	-1.3					
Kentucky	7.6	0.25	(30)	-2.6					
Louisiana	6.3 ^{⊕*}	0.23	(49)	-19.2					
Maine	8.6	0.47	(06)	10.3					
Maryland	7.4	0.23	(33)	-5.1					
Massachusetts	8.2	0.21	(13)	5.1					
Michigan	8.0	0.17	(15)	2.6					
Minnesota	8.0	0.25	(17)	2.6					
Mississippi	6.3 ^{⊕*}	0.28	(50)	-19.2					
Missouri	7.4	0.21	(35)	-5.1					

^δ NCHS public use tape. Rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

^{*} Absolute difference between state rate and total U.S. rate is 15% or more.[⊕] Difference between state rate and total U.S. rate is statistically significant (p<=.0002).



Table XX-8
OVARY CANCER (Invasive)
 AGE-ADJUSTED SEER INCIDENCE RATES \S , SEER MORTALITY RATES δ , AND 5-YEAR RELATIVE SURVIVAL RATES ϕ
 By Registry, Race and Age

	All Races, Females			White Females			Black Females		
	All	<50	50+	All	<50	50+	All	<50	50+
<u>SEER INCIDENCE RATES\S, 1989-93</u>									
REGISTRY									
Standard 9 registries	15.1	5.1	45.9	15.9	5.4	48.2	10.5	3.0	33.6
San Francisco-Oakland	15.8	5.2	48.6	17.3	5.6	53.4	12.2	3.6	38.7
Connecticut	14.9	5.0	45.5	15.4	5.3	46.6	9.7	3.1	30.2
Metropolitan Detroit	14.6	5.0	44.2	15.9	5.7	47.3	9.9	3.0	31.4
Hawaii	12.6	5.4	35.0	18.4	7.1	52.9	-	-	-
Iowa	15.8	5.8	46.7	16.0	6.0	47.0	-	-	-
New Mexico	13.4	4.6	40.5	13.2	4.3	40.6	-	-	-
Seattle-Puget Sound	17.1	5.6	52.4	17.4	5.7	53.5	12.1	5.3	33.2
Utah	14.1	4.8	42.9	14.2	4.8	43.1	-	-	-
Metropolitan Atlanta	14.4	4.4	45.0	15.6	5.1	47.8	10.5	2.6	34.7
San Jose-Monterey	14.9	5.5	43.8	15.6	5.5	46.6	8.4	3.3	24.2
Los Angeles	15.0	5.3	45.0	16.1	5.6	48.8	10.3	3.5	31.4
<u>SEER MORTALITY RATESδ, 1989-93</u>									
REGISTRY									
Standard 9 registries	7.8	1.1	28.5	8.3	1.1	30.2	5.9	0.6	22.0
San Francisco-Oakland	8.1	1.0	29.9	9.3	1.1	34.5	6.2	0.6	23.7
Connecticut	7.6	1.0	28.0	7.8	1.1	28.5	5.7	0.6	21.1
Metropolitan Detroit	7.7	1.2	27.9	8.1	1.2	29.4	6.1	0.8	22.5
Hawaii	5.8	1.1	20.4	8.9	1.2	32.7	-	-	-
Iowa	8.2	1.1	30.0	8.2	1.1	30.2	-	-	-
New Mexico	6.7	1.0	24.4	6.7	1.0	24.6	-	-	-
Seattle-Puget Sound	8.9	1.2	32.7	9.1	1.2	33.6	6.1	1.6	19.9
Utah	7.6	1.2	27.1	7.6	1.2	27.2	-	-	-
Metropolitan Atlanta	7.1	0.8	26.5	7.9	1.0	29.1	4.8	0.3	18.6
San Jose-Monterey	7.5	1.1	27.0	7.9	1.3	28.3	8.2	0.0	33.4
Los Angeles	7.9	1.0	29.2	8.6	1.1	31.8	5.8	0.9	21.2
<u>SURVIVAL RATESϕ, 1986-92</u>									
REGISTRY									
Standard 9 registries	45.5	73.0	35.3	45.6	72.7	35.8	40.0	74.4	25.0
San Francisco-Oakland	44.1	72.4	34.0	42.9	70.0	34.1	35.2 \uparrow	76.5 \uparrow	19.3 \uparrow
Connecticut	43.7	71.9	34.4	43.7	71.9	34.5	44.6 \uparrow	-	29.1 \uparrow
Metropolitan Detroit	44.0	75.3	33.5	45.2	75.6	35.4	37.1	72.3 \uparrow	23.1
Hawaii	50.4	74.6 \uparrow	39.8	52.6 \uparrow	74.6 \uparrow	41.3 \uparrow	-	-	-
Iowa	45.2	74.0	35.7	45.4	74.0	35.9	-	-	-
New Mexico	45.1	73.1	33.3	45.7	74.8	34.3	-	-	-
Seattle-Puget Sound	47.5	71.4	38.1	47.6	71.7	38.1	-	-	-
Utah	46.3	71.4	34.6	46.4	71.4	34.8	-	-	-
Metropolitan Atlanta	49.7	71.0	38.6	50.0	70.4	39.9	49.0 \uparrow	74.2 \uparrow	33.7 \uparrow

Note: San Jose-Monterey and Los Angeles are new SEER areas as of 1992 and their data are only used in this table and not other tables in this section.

- \S SEER Program. Incidence rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population.
 δ NCHS public use tape. Mortality rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population.
 ϕ Survival rates are relative rates expressed as percents and are based on follow-up of patients through 1993.
 \uparrow The standard error of the survival rate is between 5 and 10 percentage points.
 \uparrow The standard error of the survival rate is greater than 10 percentage points.
 - Statistic could not be calculated or populations are inadequate.



SEER Incidence and U.S. Mortality Ovary, All Ages By Race



The Incidence trends represented by the thin line excludes histologies 8442, 8451, 8462, 8472, and 8473 which only became reportable as malignancies after 1987.


Figure XX-1

SEER Incidence and U.S. Mortality Ovary, Under 65 Years of Age By Race



The Incidence trends represented by the thin line excludes histologies 8442, 8451, 8462, 8472, and 8473 which only became reportable as malignancies after 1987.

SEER Incidence and U.S. Mortality Ovary, Ages 65 and Over By Race



The Incidence trends represented by the thin line excludes histologies 8442, 8451, 8462, 8472, and 8473 which only became reportable as malignancies after 1987.

Figure XX-3