

Table VII-1  
CORPUS AND UTERUS, NOS CANCER (Invasive)

TRENDS IN SEER INCIDENCE AND U.S. MORTALITY USING THE JOINPOINT REGRESSION PROGRAM, 1973-1998  
WITH UP TO THREE JOINPOINTS BY RACE AND AGE

	All Races		Whites		Blacks	
	Year	APC	Year	APC	Year	APC
<u>SEER Cancer Incidence</u>						
Age - All Ages	1973-75	6.8★	1973-75	7.5★	1973-98	0.1
	1975-79	-6.5★	1975-79	-6.6★		
	1979-88	-1.9★	1979-88	-1.8★		
	1988-98	0.5★	1988-98	0.4★		
Age - Under 50	1973-84	-3.5★	1973-82	-4.0★	1973-98	-0.7
	1984-98	0.2	1982-98	-0.4		
Age - 50 and Over	1973-75	8.4★	1973-75	9.0★	1973-98	0.2
	1975-79	-6.7★	1975-79	-6.8★		
	1979-88	-1.9★	1979-88	-1.9★		
	1988-98	0.5★	1988-98	0.5★		
<u>U.S. Cancer Mortality</u>						
Age - All Ages	1973-89	-1.7★	1973-91	-1.7★	1973-75	-7.5
	1989-98	-0.6★	1991-98	-0.4	1975-98	-0.9★
Age - Under 50	1973-84	-5.1★	1973-84	-4.8★	1973-98	-4.6★
	1984-98	-0.9★	1984-98	-0.6		
Age - 50 and Over	1973-91	-1.5★	1973-83	-1.3★	1973-98	-0.8★
	1991-98	-0.4	1983-89	-2.3★		
			1989-98	-0.5★		

The APC is the Annual Percent Change based on rates age-adjusted to the 1970 U.S. standard population using the Joinpoint Regression Program.

\* The APC is significantly different from zero ( $p < .05$ ).

Table VII-2  
CORPUS AND UTERUS, NOS CANCER (Invasive)

SEER INCIDENCES AND U.S. MORTALITY<sup>§</sup> RATES, AGE-ADJUSTED AND AGE-SPECIFIC RATES, BY RACE

	Incidence§			US Mortality <sup>§</sup>		
	All Races, Females	White Females	Black Females	All Races, Females	White Females	Black Females
<b>AGE-SPECIFIC RATES, 1994-98</b>						
AGE AT DIAGNOSIS/DEATH:						
0-4	-	-	-	-	-	-
5-9	-	-	-	-	-	-
10-14	-	-	-	-	-	-
15-19	-	-	-	-	-	-
20-24	-	-	-	-	-	-
25-29	0.8	-	-	0.1	-	-
30-34	2.5	2.0	-	0.2	0.2	-
35-39	5.7	5.5	-	0.4	0.3	0.5
40-44	12.2	11.9	9.5	0.7	0.7	0.9
45-49	23.9	24.3	9.7	1.7	1.7	1.8
50-54	48.0	49.8	26.9	3.3	3.1	4.5
55-59	69.7	74.1	41.0	6.0	5.6	9.4
60-64	85.0	89.6	69.6	10.9	9.8	21.2
65-69	91.7	96.6	81.0	15.4	14.0	30.3
70-74	105.4	111.1	80.7	21.6	20.1	40.8
75-79	107.7	114.2	69.0	25.2	24.0	42.7
80-84	95.4	100.1	85.6	30.3	29.2	46.8
85+	70.3	72.1	72.0	36.7	35.9	50.8
<b>AGE-ADJUSTED RATES, 1994-98</b>						
AGE AT DIAGNOSIS/DEATH:						
All ages	21.9	22.9	15.7	3.3	3.1	5.7
Under 65	13.7	14.2	8.8	1.2	1.1	2.0
65 and over	97.3	102.5	78.5	22.2	20.9	38.9
All ages (world std.) <sup>⊕</sup>	18.1	18.9	12.9	2.5	2.3	4.3

SEER Program. Rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population, unless noted.  
NCHS public use tape. Rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population, unless noted.  
Rates are per 100,000 and are age-adjusted to the world standard population.  
Statistic not shown. Rate based on less than 25 cases for the time interval.

Table VII-3  
CORPUS AND UTERUS, NOS CANCER (Invasive)

SEER INCIDENCES AND U.S. MORTALITY<sup>D</sup> RATES, AGE-ADJUSTED RATES, BY RACE AND AGE

	All Races, Females			White Females			Black Females		
	All	<50	50+	All	<50	50+	All	<50	50+
<b>SEER INCIDENCE RATES</b>									
<b>YEAR OF DIAGNOSIS:</b>									
1973	28.4	5.3	99.5	29.5	5.4	103.8	15.0	-	54.3
1974	30.8	4.9	110.8	32.4	5.1	116.8	13.0	-	46.1
1975	32.1	4.9	116.1	33.8	5.1	122.3	17.0	-	61.3
1976	31.0	4.7	112.1	32.7	4.7	118.9	14.5	-	51.7
1977	28.5	4.3	103.1	29.8	4.4	108.2	17.1	-	61.9
1978	26.5	4.5	94.3	28.0	4.7	99.8	16.0	-	56.6
1979	24.9	4.0	89.4	26.1	4.0	94.3	14.4	-	49.9
1980	24.2	3.7	87.5	25.3	3.7	92.0	14.2	-	52.4
1981	24.1	4.1	85.6	25.1	4.0	90.2	14.3	-	52.4
1982	23.6	3.7	85.0	24.9	3.8	89.9	14.6	-	54.0
1983	23.4	3.6	84.4	24.6	3.6	89.2	16.0	-	57.6
1984	22.6	3.4	81.8	24.0	3.6	86.8	14.8	-	56.3
1985	22.0	3.4	79.5	23.2	3.6	83.7	15.2	-	56.9
1986	21.3	3.6	75.8	22.4	3.7	80.2	14.2	-	52.0
1987	21.6	3.7	76.7	22.8	3.9	81.0	13.5	-	51.2
1988	20.5	3.4	73.3	21.5	3.3	77.4	14.3	2.1	51.7
1989	21.3	3.4	76.4	22.2	3.6	79.6	16.5	-	63.2
1990	21.8	3.4	78.6	23.2	3.6	83.7	14.7	-	54.8
1991	21.3	3.3	77.0	22.5	3.3	81.9	14.9	2.4	53.2
1992	21.5	3.6	76.6	22.7	3.6	81.5	14.7	2.1	53.4
1993	21.0	3.9	73.7	22.2	3.8	78.8	14.6	2.7	51.1
1994	21.9	3.8	77.7	22.8	3.8	81.6	15.5	2.0	57.3
1995	22.0	3.5	79.1	22.9	3.4	83.0	16.4	1.8	61.2
1996	21.5	3.4	77.4	22.4	3.2	81.4	16.5	2.4	59.8
1997	22.4	3.7	80.0	23.4	3.7	84.4	14.1	1.9	51.8
1998	21.8	3.4	78.7	22.9	3.3	83.3	16.1	2.0	59.5
1994-98	21.9	3.6	78.6	22.9	3.5	82.8	15.7	2.0	57.9
<b>U.S. MORTALITY RATES</b>									
<b>YEAR OF DEATH:</b>									
1973	4.6	0.5	17.4	4.3	0.4	16.4	8.0	1.1	29.5
1974	4.5	0.5	16.8	4.2	0.4	15.9	7.0	0.7	26.6
1975	4.3	0.4	16.4	4.1	0.4	15.7	6.6	0.7	24.8
1976	4.3	0.4	16.6	4.1	0.4	15.8	6.8	0.6	25.6
1977	4.3	0.4	16.2	4.0	0.4	15.4	7.0	0.6	26.5
1978	4.2	0.4	15.8	3.9	0.4	15.0	6.8	0.6	25.8
1979	4.1	0.3	15.7	3.8	0.3	14.9	6.6	0.6	25.4
1980	4.2	0.3	16.0	3.9	0.3	15.1	6.9	0.6	26.3
1981	4.0	0.3	15.3	3.8	0.3	14.6	6.1	0.4	23.8
1982	3.9	0.3	15.2	3.7	0.3	14.3	6.3	0.3	24.7
1983	3.9	0.3	15.1	3.7	0.2	14.4	6.0	0.5	23.1
1984	3.8	0.3	14.7	3.6	0.2	13.9	6.3	0.5	23.9
1985	3.7	0.3	14.5	3.6	0.3	13.7	6.1	0.4	23.6
1986	3.6	0.3	13.9	3.4	0.2	13.1	6.0	0.4	23.3
1987	3.6	0.3	13.7	3.4	0.2	13.0	5.9	0.4	22.9
1988	3.5	0.2	13.6	3.3	0.2	12.7	6.1	0.3	23.8
1989	3.4	0.3	13.2	3.2	0.3	12.3	5.9	0.4	23.0
1990	3.5	0.2	13.5	3.3	0.2	12.6	6.0	0.3	23.4
1991	3.4	0.2	12.9	3.1	0.2	12.0	6.0	0.4	23.3
1992	3.4	0.2	13.1	3.2	0.2	12.3	5.7	0.3	22.4
1993	3.3	0.2	12.9	3.1	0.2	12.0	6.0	0.3	23.4
1994	3.3	0.2	13.0	3.1	0.2	12.1	5.9	0.2	23.2
1995	3.3	0.3	12.8	3.2	0.3	12.1	5.3	0.3	20.9
1996	3.3	0.2	12.8	3.1	0.2	11.9	5.9	0.3	23.2
1997	3.2	0.2	12.5	3.0	0.2	11.6	5.5	0.2	22.0
1998	3.3	0.2	12.8	3.1	0.2	12.0	5.7	0.3	22.1
1994-98	3.3	0.2	12.8	3.1	0.2	12.0	5.7	0.3	22.3

SEER Program. Rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population.  
NCHS public use tape. Rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population.  
Statistic not shown. Rate based on less than 25 cases for the time interval.

Table VII-4  
CORPUS AND UTERUS, NOS CANCER (Invasive)

SURVIVAL RATES, BY RACE, DIAGNOSIS YEAR, STAGE AND AGE

	All Races, Females			White Females			Black Females		
	All	<50	50+	All	<50	50+	All	<50	50+
<b>5-YR RELATIVE SURVIVAL RATES</b>									
YEAR OF DIAGNOSIS:									
1960-63 <sup>a</sup>	-	-	-	73	-	-	31	-	-
1970-73 <sup>a</sup>	-	-	-	81	-	-	44	-	-
1974-76 <sup>b</sup>	87.9	92.1	87.4	88.8	92.9	88.3	61.2	84.3 <sup>†</sup>	56.3
1977-79 <sup>b</sup>	85.0	92.1	84.0	86.3	92.9	85.4	57.8	80.7 <sup>†</sup>	52.6
1980-82 <sup>b</sup>	81.6	91.0	80.4	82.8	92.3	81.7	54.8	84.8 <sup>†</sup>	50.2
1983-85 <sup>b</sup>	82.8	92.5	81.6	84.5	93.3	83.4	54.2	81.4 <sup>†</sup>	50.1
1986-88 <sup>b</sup>	82.8	92.3	81.4	84.4	92.5	83.2	57.0	86.5	51.7
1989-91 <sup>b</sup>	83.7	88.5	82.9	85.6	90.8	84.8	57.3	71.7 <sup>†</sup>	54.8
1992-97 <sup>b</sup>	84.0 <sup>Ω</sup>	90.5	82.8 <sup>Ω</sup>	85.8 <sup>Ω</sup>	91.8	84.8 <sup>Ω</sup>	58.9	81.1	53.8
1992-97 <sup>b</sup>	ALL STAGES	84.0	90.5	82.8	85.8	91.8	84.8	58.9	81.1
LOCALIZED	96.1	96.6	96.1	96.9	97.8	96.8	82.9	88.5	81.2
REGIONAL	62.7	80.4	59.7	65.1	80.0	62.8	42.7	81.9 <sup>†</sup>	35.6 <sup>†</sup>
DISTANT	25.8	46.3	22.6	27.7	50.2	24.4	13.1	-	8.5
UNSTAGED	49.1	77.7 <sup>†</sup>	44.7	47.6	73.6 <sup>†</sup>	44.1	48.9 <sup>†</sup>	-	44.8 <sup>†</sup>
<b>STAGE DISTRIBUTION (%) 1992-97<sup>b</sup></b>									
ALL STAGES									
Number of cases	16,472	2,358	14,114	14,369	1,840	12,529	1,019	165	854
Percent	100%	100%	100%	100%	100%	100%	100%	100%	100%
LOCALIZED	73	77	73	75	78	74	52	67	49
REGIONAL	14	13	14	13	13	13	22	17	23
DISTANT	8	7	9	8	7	8	18	10	19
UNSTAGED	5	3	5	4	3	5	9	6	9
<b>5-YR RELATIVE SURVIVAL RATES, 1992-97<sup>b</sup></b>									
AGE AT DIAGNOSIS:									
<45	90.7	-	-	91.5	-	-	85.0	-	-
45-54	90.5	-	-	91.8	-	-	72.3	-	-
55-64	86.7	-	-	89.4	-	-	57.3	-	-
65-74	82.9	-	-	85.2	-	-	53.5	-	-
75+	73.6	-	-	75.5	-	-	40.0 <sup>†</sup>	-	-
Under 65	88.6	-	-	90.5	-	-	67.7	-	-
65 and over	79.4	-	-	81.5	-	-	49.2	-	-

<sup>a</sup> Rates are based on End Results data from a series of hospital registries and one population-based registry.  
<sup>b</sup> Rates are from the SEER Program. They are based on data from population-based registries in Connecticut, New Mexico, Utah, Iowa, Hawaii, Atlanta, Detroit, Seattle-Puget Sound and San Francisco-Oakland. Rates are based on follow-up of patients through 1997.

The difference in rates between 1974-76 and 1992-97 is statistically significant ( $p < .05$ ).  
The standard error of the survival rate is between 5 and 10 percentage points.  
The standard error of the survival rate is greater than 10 percentage points.  
Statistic could not be calculated.

Ω ↑ -

Table VII-5  
CORPUS AND UTERUS, NOS CANCER (Invasive)

INCIDENCE, MORTALITY, AND SURVIVAL RATES

By Year of Diagnosis/Death

All Races, Females

Year of Diagnosis/Death

	1975	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998
<b>Incidence</b>																				
SEER	32.1	24.2	24.1	23.6	23.4	22.6	22.0	21.3	21.6	20.5	21.3	21.8	21.3	21.5	21.0	21.9	22.0	21.5	22.4	21.8
<b>Mortality</b>																				
U.S.	4.3	4.2	4.0	3.9	3.9	3.8	3.7	3.6	3.6	3.5	3.4	3.5	3.4	3.4	3.4	3.3	3.3	3.3	3.2	3.3
SEER	3.9	4.2	4.1	4.3	4.2	4.0	4.1	3.4	3.3	3.3	3.5	3.4	3.4	3.5	3.2	3.1	3.3	3.3	2.9	3.3
<b>Relative Survival Rates (SEER)</b>																				
1-year	94.5	92.4	91.1	91.6	91.9	91.6	93.2	93.0	92.1	92.3	92.3	92.7	92.1	93.1	92.5	91.9	93.2	91.9	93.2	
2-year	91.9	86.3	86.4	86.4	86.9	87.5	88.2	87.0	87.4	88.4	88.0	88.1	88.1	89.8	88.6	87.4	88.6	88.2		
3-year	90.0	83.4	84.0	83.9	83.8	85.2	85.5	85.3	84.3	85.4	85.9	85.3	85.8	87.4	85.9	85.1	86.2			
4-year	88.6	81.8	82.7	83.2	82.5	84.3	83.7	83.9	83.5	83.9	84.9	84.0	84.2	86.0	84.6	83.5				
5-year	88.3	80.9	81.9	81.9	81.5	83.6	83.4	82.9	82.4	83.2	84.1	83.2	83.6	85.9	83.8					
6-year	88.1	80.7	81.4	81.5	80.9	83.4	82.9	82.5	81.6	82.9	83.7	82.1	82.4	85.2						
7-year	87.9	80.7	80.9	81.5	80.1	82.7	82.3	82.4	81.6	82.6	83.7	82.1	82.3							
8-year	87.8	80.0	80.4	81.0	79.7	82.6	82.1	82.4	81.4	82.6	82.6	81.8								
9-year	87.8	80.0	80.0	81.0	79.3	82.2	81.7	82.1	81.4	82.1	82.6									
10-year	87.8	80.0	79.8	81.0	79.2	81.7	81.6	82.1	81.4	81.6										
11-year	87.8	79.8	79.8	81.0	79.0	81.3	81.6	82.0	81.2											
12-year	87.8	79.4	79.8	81.0	78.9	81.1	81.6	81.7												
13-year	87.8	79.1	79.5	81.0	78.7	81.1	81.1													
14-year	87.8	79.1	79.5	81.0	77.4	81.1														
15-year	87.8	79.1	79.5	80.4	77.2															
16-year	87.8	78.7	79.0	80.4																
17-year	87.8	78.7	78.5																	
18-year	87.8	78.7																		
19-year	87.8																			
20-year	87.8																			
21-year	87.8																			
22-year	87.8																			
23-year	87.8																			

Note: Incidence and mortality rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population.

Survival rates are relative rates expressed as percents.

Table VII-6  
CORPUS AND UTERUS, NOS CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,  
Given Cancer Free At Current Age By Race

Lifetime Risk (Percent) of Being Diagnosed (11 SEER Areas) With Cancer and  
Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

1996-1998

All Races

Current Age	<u>+10 yrs</u>	<u>+20 yrs</u>	<u>+30 yrs</u>	<u>Eventually</u>
0	0.00	0.00	0.01	2.69
10	0.00	0.01	0.05	2.72
20	0.01	0.05	0.21	2.72
30	0.04	0.21	0.76	2.73
40	0.17	0.72	1.48	2.71
50	0.57	1.34	2.11	2.60
60	0.81	1.63	2.06	2.14

Lifetime Risk of Being Diagnosed =      2.69%  
Lifetime Risk of Dying =      0.51%

Whites

Current Age	<u>+10 yrs</u>	<u>+20 yrs</u>	<u>+30 yrs</u>	<u>Eventually</u>
0	0.00	0.00	0.01	2.87
10	0.00	0.01	0.05	2.89
20	0.01	0.05	0.21	2.90
30	0.04	0.21	0.80	2.90
40	0.17	0.76	1.58	2.89
50	0.60	1.43	2.26	2.77
60	0.87	1.75	2.19	2.27

Lifetime Risk of Being Diagnosed =      2.87%  
Lifetime Risk of Dying =      0.49%

Blacks

Current Age	<u>+10 yrs</u>	<u>+20 yrs</u>	<u>+30 yrs</u>	<u>Eventually</u>
0	0.00	0.00	0.01	1.75
10	0.00	0.01	0.03	1.79
20	0.01	0.03	0.12	1.79
30	0.02	0.12	0.44	1.80
40	0.10	0.42	1.02	1.81
50	0.34	0.96	1.45	1.79
60	0.69	1.22	1.52	1.59

Lifetime Risk of Being Diagnosed =      1.75%  
Lifetime Risk of Dying =      0.73%

Table VII-7  
CORPUS AND UTERUS, NOS CANCER (Invasive)

AGE-ADJUSTED SEER INCIDENCE AND U.S. MORTALITY RATES

By Race/Ethnicity and Sex

Incidence

	Rate 1992-1998 Rate per 100,000 persons	Trend 1992-1998 EAPC (%)
	Females	Females
<b>RACE/ETHNICITY</b>		
All Races	21.2	0.4
White	22.5	0.2
White Hispanic	14.1	1.3
White Non-Hispanic	23.4	0.2
Black	15.3	1.5
Asian/Pacific Islander	14.6	2.8
Amer Ind/Alask Nat	8.8	-
Hispanic <sup>®</sup>	13.2	1.3

Mortality<sup>♦</sup>

	Rate 1992-1998 Rate per 100,000 persons	Trend 1992-1998 EAPC (%)
	Females	Females
<b>RACE/ETHNICITY</b>		
All Races	3.3	-0.5
White	3.1	-0.5
White Hispanic	2.6	0.0
White Non-Hispanic	3.1	-0.4
Black	5.7	-0.6
Asian/Pacific Islander	1.8	2.6
Amer Ind/Alask Nat	2.0	-5.3*
Hispanic <sup>®</sup>	2.5	-0.2

- The EAPC is the Estimated Annual Percent Change over the time interval.  
 - Statistic not shown. Rate based on less than 25 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.  
 x Incidence data are from the 11 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, and Los Angeles) and Alaska. Unless specified, other tables use data from 9 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

<sup>®</sup> Hispanic is not mutually exclusive from whites, blacks, Asian Pacific Islanders, and American Indians/Native Alaskans. For incidence, all 11 SEER areas are included. For mortality, information is included for all states except Connecticut, Oklahoma, Louisiana, and New Hampshire.

<sup>♦</sup> Mortality data are analyzed from a public-use file provided by the National Center for Health Statistics (NCHS).

\* The EAPC is significantly different from zero ( $p < .05$ ).

Table VII-8  
CORPUS AND UTERUS, NOS CANCER (Invasive)

AGE-ADJUSTED SEER INCIDENCE RATES\$ AND SEER MORTALITY RATES§  
By Registry, Race and Age

	All Races, Females			White Females			Black Females		
	All	<50	50+	All	<50	50+	All	<50	50+
<b>SEER INCIDENCE RATES\$, 1994-98</b>									
REGISTRY									
11 registries	21.4	3.5	76.6	22.6	3.5	81.6	15.6	2.1	57.5
9 registries	21.9	3.6	78.6	22.9	3.5	82.8	15.7	2.0	57.9
San Francisco-Oakland	22.0	3.3	79.7	24.3	3.4	88.6	14.7	-	54.8
Connecticut	25.2	3.8	91.1	25.7	3.8	93.1	17.2	-	61.1
Metropolitan Detroit	21.0	3.7	74.3	22.4	4.0	78.9	15.6	2.4	56.2
Hawaii	20.8	7.0	63.5	21.8	-	80.5	-	-	-
Iowa	24.2	3.6	87.7	24.2	3.5	87.9	-	-	-
New Mexico	16.4	3.0	57.5	16.6	2.7	59.6	-	-	-
Seattle-Puget Sound	23.3	3.3	85.2	24.1	3.3	87.9	15.2	-	61.5
Utah	20.2	3.7	71.0	20.3	3.6	71.9	-	-	-
Metropolitan Atlanta	19.1	2.7	69.5	19.6	3.1	70.7	17.5	1.6	66.4
San Jose-Monterey	21.3	2.6	78.9	23.1	2.6	86.3	-	-	-
Los Angeles	19.6	3.4	69.6	21.3	3.6	76.2	15.7	2.3	57.2
<b>SEER MORTALITY RATES§, 1994-98</b>									
REGISTRY									
11 registries	3.2	0.2	12.4	3.2	0.2	12.3	4.8	0.4	18.6
9 registries	3.2	0.2	12.4	3.1	0.2	12.1	4.7	-	18.2
San Francisco-Oakland	3.0	-	11.4	3.0	-	11.7	4.4	-	17.1
Connecticut	3.2	-	12.2	3.1	-	11.9	4.6	-	17.0
Metropolitan Detroit	3.4	0.3	12.9	3.1	-	12.0	4.4	-	16.7
Hawaii	2.7	-	10.3	3.0	-	-	-	-	-
Iowa	3.8	-	14.8	3.7	-	14.6	-	-	-
New Mexico	2.8	-	10.8	2.8	-	11.0	-	-	-
Seattle-Puget Sound	2.9	-	11.6	3.0	-	11.8	-	-	-
Utah	3.2	-	12.8	3.2	-	13.0	-	-	-
Metropolitan Atlanta	3.2	-	12.6	2.4	-	9.3	6.4	-	25.5
San Jose-Monterey	2.8	-	10.7	3.0	-	11.8	-	-	-
Los Angeles	3.4	0.3	12.9	3.4	0.3	12.9	5.2	-	19.9

Note: San Jose-Monterey and Los Angeles are new SEER areas as of 1992 and their data are only used in tables where indicated.  
 §/○ - SEER Program. Incidence rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population.  
 NCHS public use tape. Mortality rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population.  
 Statistic could not be calculated or populations are inadequate.

Table VII-9  
CORPUS AND UTERUS, NOS CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER MORTALITY<sup>d</sup> RATES BY STATE, 1994-98

All Races, Females

State	Rate	SE	Rank	PD
TOTAL U.S.	3.3	0.02		
High Five States				
District of Columbia	5.5 <sup>3/4</sup> <sup>e</sup>	0.54	(01)	66.7
New Jersey	4.1 <sup>3/4</sup> <sup>e</sup>	0.12	(02)	24.2
Vermont	4.0 <sup>f</sup>	0.46	(03)	21.2
New York	4.0 <sup>3/4</sup> <sup>e</sup>	0.08	(04)	21.2
Ohio	3.9 <sup>3/4</sup> <sup>e</sup>	0.10	(05)	18.2
Low Five States				
Colorado	2.7 <sup>3/4</sup> <sup>e</sup>	0.16	(47)	-18.2
Mississippi	2.6 <sup>3/4</sup> <sup>e</sup>	0.17	(48)	-21.2
Arizona	2.5 <sup>3/4</sup> <sup>e</sup>	0.13	(49)	-24.2
Alaska	2.1 <sup>f</sup>	0.49	(50)	-36.4
North Dakota	2.0 <sup>3/4</sup> <sup>e</sup>	0.29	(51)	-39.4

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	2.8 <sup>f</sup>	0.14	(41)	-15.2	Montana	2.9	0.31	(39)	-12.1
Alaska	2.1 <sup>f</sup>	0.49	(50)	-36.4	Nebraska	2.9	0.23	(38)	-12.1
Arizona	2.5 <sup>3/4</sup> <sup>e</sup>	0.13	(49)	-24.2	Nevada	2.8 <sup>f</sup>	0.25	(42)	-15.2
Arkansas	3.2	0.19	(24)	-3.0	New Hampshire	3.8 <sup>f</sup>	0.32	(07)	15.2
California	3.1 <sup>3/4</sup>	0.06	(32)	-6.1	New Jersey	4.1 <sup>3/4</sup> <sup>e</sup>	0.12	(02)	24.2
Colorado	2.7 <sup>3/4</sup> <sup>e</sup>	0.16	(47)	-18.2	New Mexico	2.8 <sup>f</sup>	0.24	(44)	-15.2
Connecticut	3.2	0.16	(27)	-3.0	New York	4.0 <sup>3/4</sup> <sup>e</sup>	0.08	(04)	21.2
Delaware	3.6	0.39	(11)	9.1	North Carolina	3.3	0.12	(16)	0.0
District of Columbia	5.5 <sup>3/4</sup> <sup>e</sup>	0.54	(01)	66.7	North Dakota	2.0 <sup>3/4</sup> <sup>e</sup>	0.29	(51)	-39.4
Florida	2.9 <sup>3/4</sup>	0.07	(40)	-12.1	Ohio	3.9 <sup>3/4</sup> <sup>e</sup>	0.10	(05)	18.2
Georgia	3.1	0.12	(30)	-6.1	Oklahoma	2.7 <sup>3/4</sup> <sup>e</sup>	0.16	(45)	-18.2
Hawaii	2.7 <sup>f</sup>	0.28	(46)	-18.2	Oregon	3.3	0.17	(18)	0.0
Idaho	3.0	0.30	(37)	-9.1	Pennsylvania	3.7 <sup>3/4</sup>	0.09	(08)	12.1
Illinois	3.6	0.10	(10)	9.1	Rhode Island	3.6	0.33	(09)	9.1
Indiana	3.5	0.14	(13)	6.1	South Carolina	3.3	0.17	(17)	0.0
Iowa	3.8 <sup>f</sup>	0.19	(06)	15.2	South Dakota	3.1	0.36	(31)	-6.1
Kansas	3.0	0.18	(36)	-9.1	Tennessee	3.2	0.13	(25)	-3.0
Kentucky	3.0	0.15	(35)	-9.1	Texas	3.0	0.08	(34)	-9.1
Louisiana	3.2	0.15	(26)	-3.0	Utah	3.2	0.26	(23)	-3.0
Maine	3.2	0.27	(28)	-3.0	Vermont	4.0 <sup>f</sup>	0.46	(03)	21.2
Maryland	3.2	0.15	(21)	-3.0	Virginia	3.4	0.13	(14)	3.0
Massachusetts	3.2	0.13	(20)	-3.0	Washington	2.8 <sup>3/4</sup> <sup>e</sup>	0.13	(43)	-15.2
Michigan	3.5	0.11	(12)	6.1	West Virginia	3.2	0.21	(22)	-3.0
Minnesota	3.0	0.15	(33)	-9.1	Wisconsin	3.3	0.14	(19)	0.0
Mississippi	2.6 <sup>3/4</sup> <sup>e</sup>	0.17	(48)	-21.2	Wyoming	3.1	0.47	(29)	-6.1
Missouri	3.4	0.13	(15)	3.0					

<sup>d</sup> NCHS public use tape. Rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population.  
<sup>SE</sup> Standard error of the rate.


<sup>PD</sup> Percent difference between state rate and total U.S. rate.

<sup>f</sup> Absolute difference between state rate and total U.S. rate is 15% or more.

<sup>3/4</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).

Figure VII-1

# SEER Incidence and U.S. Mortality Rates<sup>+</sup> Corpus Uteri By Race




<sup>+</sup> Rates are age-adjusted to the 1970 U.S. standard million population. Regression lines are calculated using the Joinpoint Regression Program.


Figure VII-2

# SEER Incidence and U.S. Mortality Rates<sup>+</sup> Corpus Uteri, By Age and Race


## White - Under 50 Years of Age




## Black - Under 50 Years of Age



## White - Ages 50 and Over



## Black - Ages 50 and Over



• Incidence

× Mortality

<sup>+</sup> Rates are age-adjusted to the 1970 U.S. standard million population. Regression lines are calculated using the Joinpoint Regression Program.