

SEER Incidence* and US Death Rates,# 1996-2000 5-Year Relative Survival Rates,^ 1992-1999 All Cancer Combined, by Race and Sex




Figure 1-9


* Incidence rates are from the SEER 12 areas and are age-adjusted to the 2000 US standard population by 5-year age groups.

Death rates are from the NCHS public use data file and are age-adjusted to the 2000 US standard population by 5-year age groups.

^ Survival rates are from the SEER 9 areas. Relative survival rates are expressed as percents.

Figure I-11

5-Year Relative Survival Rates SEER Program, 1992-1999 Both Sexes, by Race



Source: SEER 9 areas.


SEER Cancer Incidence and US Death Rates, 1996-2000 By Cancer Site and Race

Incidence Cancer Site Mortality


Lung and Bronchus




Colon and Rectum



Female Breast



Prostate



Source: SEER 12 areas and NCHS public use data file.

^^ Hispanic is not mutually exclusive from Whites, Blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics excludes Detroit and Hawaii. Mortality data for Hispanics excludes cases from Connecticut, Oklahoma, New York, and New Hampshire.

Rates are per 100,000 and age-adjusted to the 2000 U.S. standard population by 5-year age groups.