

Prostate EOD Revision, Effective for 1995 Cases  
Clinical evaluation - extension of tumor

CODE	CATEGORIES - Clinical evaluation excluding information from prostatectomy
<b>NO EXTENSION BEYOND PROSTATE</b>	
00	IN SITU; Noninvasive; intraepithelial
<b>Clinically inapparent tumor not palpable or visible by imaging; incidentally found microscopic carcinoma (latent, occult), in one or both lobes</b>	
10	No. of foci or % of involved tissue not specified (A, NOS) Clinically inapparent tumor confined to prostate, NOS (T1, NOS)
11	<= 3 microscopic foci (A1 focal)
12	> 3 microscopic foci (A2 diffuse)
13	Incidental histologic finding in 5% or less of tissue resected (T1a)
14	Incidental histologic finding in more than 5% of tissue resected (T1b)
15	Tumor identified by needle bx, e.g., for elevated PSA (T1c)
<b>Clinically/radiographically apparent</b>	
20	Involvement of one lobe, NOS (B)
21	1/2 or less of one lobe involved (T2a)
22	More than 1/2 of 1 lobe involved, not both lobes (T2b)
23	More than one lobe involved (B); (T2b)
24	Clinically apparent tumor confined to prostate, NOS; (Stage B, NOS), (T2, NOS)
30	Not stated if clinically apparent or inapparent but Localized, NOS; Confined to prostate, NOS; Intracapsular involvement only Not stated if Stage A or B, T1 or T2
31	Into prostatic apex/arising in prostatic apex
<b>EXTENSION BEYOND PROSTATE</b>	
40	Invasion of prostatic capsule (C1)
41	Extension to periprostatic tissue (C1) Extracapsular extension (beyond prostatic capsule), NOS Through capsule, NOS
42	Unilateral extracapsular extension (T3a)
43	Bilateral extracapsular extension (T3b)
44	Extraprostatic urethra

CODE	CATEGORIES - Clinical evaluation excluding information from prostatectomy
45	Extension to seminal vesicle(s) (C2); (T3c)
49	Periprostatic extension, NOS (C, NOS; T3, NOS; unknown if seminal vesicles(s) involved)
50	Extension to or fixation to adjacent structures other than seminal vesicles: rectovesical (Denovilliers') fascia; Bladder, NOS; Ureter(s); Fixation, NOS; (T4, NOS)
51	Extension to/fixation to: Bladder neck (T4a)
52	Rectum; external sphincter (T4a)
53	Levator muscles (T4b)
60	Extension to or fixation to pelvic wall or pelvic bone (T4b)
61	Extension to or fixation to other skeletal muscle
70	Further extension to bone, soft tissue, or other organs (D2)
80	Metastasis (D2); D, not further specified
90	Unknown if extension or metastasis

Use all information except the prostatectomy to code this field based on the above codes.

Limit extent of disease information to 4 months after diagnosis in the absence of disease progression.

Give priority to codes 13-14 over codes 11-12.

Use code 30 when there is insufficient information as to whether the tumor is clinically apparent or inapparent but the tumor is confined to the prostate.

In parentheses some of the AUA stages and AJCC T numbers are given as guides in coding this field in the absence of information in the medical record. Note that some stages/T codes are in more than one category such as T2b can be either code 22 or 23. Note do not code using the T number if metastases are present.

Terms to Distinguish Apparent from Inapparent

CLINICALLY APPARENT

YES	MAYBE	NO
Nodule	Asymmetrical	1+, 2+, 3+
Hard nodule	Significant asym.	enlarged
Suspicious	Firm	30 gm size
Positive nodule	Slightly irregular	60 gm size
Hard	Nodular	Slightly enlarged
Fixed	Firm ridge	Large
? nodule	Diffusely firm	Firm without
Firm, Irregular	Abnormal	nodule
Induration		Very large
		Moderately large
		Median lobe

RADIOGRAPHICALLY APPARENT

YES	MAYBE	NO
Suspicious	Streaky densities	Mottled-appearing
Hypoechoic	in prostate	Prominent S.V.
Suggesting	Irregular	Negative
invasion	indentions	Prominent prostate
Streaky densities	(bladder)	Ultrasound
in periprostatic		negative
fat		

Pathologic evaluation of extent of disease  
Prostate, Effective for 1995 Cases  
Pathologic evaluation including prostatectomy

CODE	CATEGORIES - Pathologic evaluation including prostatectomy
<b>NO EXTENSION BEYOND PROSTATE</b>	
00	IN SITU; Noninvasive; intraepithelial
20	Involvement of one lobe, NOS (B)
21	1/2 or less of one lobe involved (pT2a)
22	More than 1/2 of 1 lobe involved, not both lobes (pT2b)
23	More than one lobe involved (B); (pT2b)
30	Localized, NOS; Confined to prostate, NOS Intracapsular involvement only; (Stage B, NOS; pT2, NOS)
31	Into prostatic apex/arising in prostatic apex
<b>EXTENSION BEYOND PROSTATE</b>	
40	Invasion of prostatic capsule (C1)
41	Extension to periprostatic tissue (C1); Extracapsular extension (beyond prostatic capsule), NOS Through capsule, NOS
42	Unilateral extracapsular extension (pT3a)
43	Bilateral extracapsular extension (pT3b)
44	Extraprostatic urethra
45	Extension to seminal vesicle(s) (C2); (pT3c)
49	Periprostatic extension, NOS Extends to posterior margins; margins involved (except urethral) (C, NOS; pT3, NOS; unknown if seminal vesicles(s) involved)
50	Extension to or fixation to adjacent structures other than seminal vesicles: rectovesical (Denovilliers') fascia; Bladder, NOS; Ureter(s); Fixation, NOS; (pT4, NOS)
51	Extension to/fixation to: Bladder neck (pT4a)
52	Rectum; external sphincter (pT4b)
53	Levator muscles (pT4b)
60	Extension to or fixation to pelvic wall or pelvic bone (pT4b)

CODE	CATEGORIES - Pathologic evaluation including prostatectomy
61	Extension to or fixation to other skeletal muscle
70	Further extension to bone, soft tissue, or other organs (D2)
80	Metastasis (D2); D, not further specified
90	Unknown if extension or metastasis
99	No prostatectomy done within 4 months after diagnosis

This field is based on all histologic information including the prostatectomy if done within 4 months of diagnosis. Use code '99' if there was no prostatectomy within 4 months after diagnosis. This would include evaluation of other pathologic tissue such as a biopsy of the rectum.

Limit extent of disease information to 4 months after diagnosis in the absence of disease progression.

Give priority to codes 41-45 over code 49.